

THE KNASTER–KURATOWSKI–MAZURKIEWICZ THEOREM AND ALMOST FIXED POINTS

SEHIE PARK

ABSTRACT. From the KKM theorem for the “closed” and “open” valued cases, we deduce a generalization of the Alexandroff–Pasynkoff theorem, existence theorems for almost fixed points of lower semicontinuous multimap, and a partial solution of the Ben-El-Mechaiekh conjecture.

1. Introduction

It is well-known that the three classical results – the Brouwer fixed point theorem, the Sperner lemma, and the Knaster–Kuratowski–Mazurkiewicz (simply, KKM) theorem – are mutually equivalent in the sense that each one can be deduced from another with or without aid of some minor results. Earlier applications of the Sperner lemma and the KKM theorem to various results closely related to Euclidean spaces or n -simplexes or n -balls were presented in [1] and [2]; see also [13].

Especially, one of the earlier proofs of the Brouwer theorem was given by the KKM theorem in [10]. Recently, the present author and Do Hong Tan [14] gave a simple proof of a generalization of the Schauder–Tychonoff type fixed

2000 *Mathematics Subject Classification*. Primary 47H04, 47H10; Secondary 52A07, 54C60, 54H25.

Key words and phrases. Topological vector space (t.v.s.), KKM theorem, almost convex subset, almost fixed point, lower semicontinuous (l.s.c.) map, upper semi-continuous (u.s.c.) map, fixed point.

The author wishes to acknowledge the financial support of the Korea Research Foundation made in the program year of 1997.

point theorem for compact maps in locally convex Hausdorff topological vector spaces, by directly using the KKM theorem. Subsequently, in [15], elementary proofs of generalizations of the Himmelberg–Idzik type fixed point theorem for convex-valued upper semicontinuous multimaps were obtained from the “open” valued version of the KKM theorem due to Kim [9] and Shih–Tan [16].

On the other hand, in 1957, Alexandroff–Pasynkoff [3] gave an elementary proof of the essentiality of the identity map of the boundary of a simplex by using a variant of the KKM theorem. Consequently, a simple proof of the Brouwer theorem could be given by using their theorem; see [13]. Moreover, in 1990, Lassonde [11] suggested an “open” version of the Alexandroff–Pasynkoff theorem.

In the present paper, we first deduce a generalization of the Alexandroff–Pasynkoff theorem by using the KKM theorem for the “closed” and “open” valued cases and note that these two theorems are actually equivalent. And then, the KKM theorem is applied to our main result, which concerns with the existence of almost fixed points of lower (respectively, upper) semicontinuous multimaps. Our new result is general enough to include properly previous results on almost fixed points due to Ky Fan [7] and Lassonde [11], and fixed point theorems due to Park and Tan [14], [15], Himmelberg [8], and many others. Finally, applying our main result, we give a partial solution to a conjecture raised by Ben-El-Mechaiekh [4], [5].

2. The KKM and Alexandroff–Pasynkoff theorems

From the KKM theorem and its open version, we have immediately the following form as by Fan in [6]:

THEOREM 1. *Let X be a subset of a topological vector space, D a nonempty subset of X such that $\text{co } D \subset X$, and $F : D \multimap X$ a multimap with closed (respectively, open) values in X . If*

$$(1) \quad \text{co } A \subset F(A)$$

for every nonempty finite subset A of D , then the family $\{F(x)\}_{x \in D}$ has the finite intersection property.

The open version of the KKM theorem was due to Kim [9] and Shih–Tan [16], and later, Lassonde [11] showed that the closed and open versions of the KKM theorem can be derived from each other.

From Theorem 1, we have the following generalization of the Alexandroff–Pasynkoff theorem [3]:

THEOREM 2. *Let X be a subset of a topological vector space, $\{A_i\}_{i=0}^n$ a family of $(n+1)$ closed (respectively, open) subsets covering X , and $\{x_i\}_{i=0}^n$ a family of*

$(n + 1)$ points of X such that $\text{co}\{x_i\}_{i=0}^n \subset X$ and $\text{co}\{x_0, \dots, \widehat{x_i}, \dots, x_n\} \subset A_i$ for each $i = 0, \dots, n$. Then $\bigcap_{i=0}^n A_i \neq \emptyset$.

PROOF. Let $D := \{x_i\}_{i=0}^n$ and let $C_0 := \text{co}\{x_0, \dots, x_{n-1}\} \subset A_n$ and $C_i := \text{co}\{x_0, \dots, \widehat{x_{i-1}}, \dots, x_n\} \subset A_{i-1}$ for $1 \leq i \leq n$. Let $F : D \multimap X$ be a map defined by $F(x_0) = A_n$ and $F(x_i) = A_{i-1}$ for $1 \leq i \leq n$. Now we show that F satisfies the requirement of Theorem 1. Note that

$$\text{co}\{x_0, \dots, x_n\} \subset X = \bigcup_{i=0}^n A_i = F(D).$$

Moreover, for any proper subset $\{x_{i_0}, \dots, x_{i_k}\}$, ($0 \leq k < n$, $0 \leq i_0 < \dots < i_k \leq n$) of D , we immediately have $\text{co}\{x_{i_0}, \dots, x_{i_k}\} \subset C_{i_j} \subset A_{i_j-1} = F(x_{i_j})$ for some j , $0 \leq j \leq k$, (with the convention $i_j = 0$ if and only if $i_j - 1 \equiv n$) and hence

$$\text{co}\{x_{i_0}, \dots, x_{i_k}\} \subset \bigcup_{j=0}^k F(x_{i_j}).$$

Consequently, condition (1) is satisfied. Now, the conclusion follows from Theorem 1. □

It is well-known that the Alexandroff–Pasyukoff theorem implies the Brouwer theorem (e.g. see [13]). Therefore, Theorem 2 is also equivalent to the KKM theorem.

3. Almost fixed point theorems

From Theorem 1, in this section, we deduce a very general almost fixed point theorem and some of its direct applications.

A nonempty subset Y of a topological vector space E is said to be almost convex [8] if for any neighbourhood V of the origin 0 of E and for any finite subset $\{y_1, \dots, y_n\}$ of Y , there exists a finite subset $\{z_1, \dots, z_n\}$ of Y , such that $z_i - y_i \in V$ for each $i = 1, \dots, n$, and $\text{co}\{z_1, \dots, z_n\} \subset Y$.

The following almost fixed point theorem is our main result in this paper:

THEOREM 3. *Let X be a subset of a Hausdorff topological vector space E and Y an almost convex dense subset of X . Let $T : X \multimap E$ be a lower (respectively, upper) semicontinuous multimap such that $T(y)$ is convex for all $y \in Y$. If there is a precompact subset K of X such that $T(y) \cap K \neq \emptyset$ for each $y \in Y$, then for any convex neighbourhood U of the origin 0 of E , there exists a point $x_U \in Y$ such that $T(x_U) \cap (x_U + U) \neq \emptyset$.*

PROOF. There exists a symmetric open neighbourhood V of 0 such that $\overline{V} + \overline{V} \subset U$. Since K is precompact in E , there exists a finite subset $\{x_0, \dots, x_n\} \subset K$ such that $K \subset \bigcup_{i=0}^n (x_i + V)$. Moreover, since Y is almost convex and dense

in X , there exists a finite subset $D = \{y_0, \dots, y_n\}$ of Y such that $y_i - x_i \in V$ for each $i = 0, \dots, n$, and $Z := \text{co}\{y_0, \dots, y_n\} \subset Y$.

If T is lower semicontinuous, for each i , let

$$F(y_i) := \{z \in Z : T(z) \cap (x_i + V) = \emptyset\},$$

which is closed in Z . Moreover, we have

$$\bigcap_{i=0}^n F(y_i) = \left\{ z \in Z : T(z) \cap \bigcup_{i=0}^n (x_i + V) = \emptyset \right\} = \emptyset$$

since $\emptyset \neq T(z) \cap K \subset T(z) \cap \bigcup_{i=0}^n (x_i + V)$ for each $z \in Y$.

If T is upper semicontinuous, for each i , let

$$F(y_i) := \{z \in Z : T(z) \cap (x_i + \bar{V}) = \emptyset\},$$

which is open in Z . Moreover, we have $\bigcap_{i=0}^n F(y_i) = \emptyset$ as in the above.

Now we apply Theorem 1 replacing (X, D) by $(Z, \{y_i\}_{i=0}^n)$. Since the conclusion of Theorem 1 does not hold, in any case, condition (1) is violated. Hence, there exist a subset $N := \{y_{i_0}, \dots, y_{i_k}\} \in \langle D \rangle$ and an $x_U \in \text{co} N \subset Y$ such that $x_U \notin F(N)$ or $T(x_U) \cap (x_{i_j} + \bar{V}) \neq \emptyset$ for all $j = 0, \dots, k$. Note that

$$(2) \quad x_{i_j} + \bar{V} = x_{i_j} - y_{i_j} + y_{i_j} + \bar{V} \subset y_{i_j} + V + \bar{V} \subset y_{i_j} + U.$$

Let L be the subspace of E generated by D and

$$M := \{y \in L : T(x_U) \cap (y + U) \neq \emptyset\}.$$

From (2) we get $T(x_U) \cap (y_{i_j} + U) \neq \emptyset$ and hence $y_{i_j} \in M$ for all $j = 0, \dots, k$. Since $L, T(x_U)$, and U are all convex, it is easily checked that M is convex. Therefore, $x_U \in M$ and, by definition of M , we get $T(x_U) \cap (x_U + U) \neq \emptyset$. \square

In case $X = Y$, Theorem 3 reduces to the following:

COROLLARY 4. *Let X be a convex subset of a Hausdorff topological vector space E . Let $T : X \multimap E$ be a lower (respectively, upper) semicontinuous multimap such that $T(x)$ is convex for each $x \in X$. If there is a precompact subset K of X such that $T(x) \cap K \neq \emptyset$ for each $x \in X$, then for every convex neighbourhood U of the origin 0 of E , there exists a point $x_U \in X$ such that $T(x_U) \cap (x_U + U) \neq \emptyset$.*

Ky Fan [7, Theorem 7] obtained Corollary 4 for a locally convex Hausdorff topological vector space E and for a lower semicontinuous multimap $T : X \multimap E$. For a single-valued map $f : X \rightarrow X$, Fan noted that Corollary 4 might be regarded as a generalization of the Tychonoff fixed point theorem to noncompact (or precompact) convex sets; see Corollary 5 below.

Lassonde [5, Théorème 4] obtained Corollary 4 for a compact upper semicontinuous map $T : X \multimap X$ having nonempty convex values.

From Theorem 3, we have the following fixed point theorem:

COROLLARY 5. *Let X be a subset of a locally convex Hausdorff topological vector space E and Y an almost convex dense subset of X . Let $T : X \multimap X$ be a compact upper semicontinuous multimap with closed values such that $T(y)$ is nonempty convex for all $y \in Y$. Then T has a fixed point $x_0 \in X$; that is, $x_0 \in T(x_0)$.*

PROOF. By Theorem 3, for each neighbourhood U of 0, there exist $x_U, y_U \in X$ such that $y_U \in T(x_U)$ and $y_U \in x_U + U$. Since $T(X)$ is relatively compact, we may assume that the net $\{y_U\}$ converges to some $x_0 \in \overline{T(X)} \subset X$. Since E is Hausdorff, the net $\{x_U\}$ also converges to x_0 . Because T is upper semicontinuous with closed values, the graph of T is closed in $X \times T(X)$ and hence we have $x_0 \in T(x_0)$. \square

Corollary 5 is recently due to the author and Do Hong Tan [15] and extends the Himmelberg-Idzik theorem and many other fixed point results in the analytical fixed point theory.

From Theorem 3, we have the following almost fixed point result:

COROLLARY 6. *Let X be a subset of a Hausdorff topological vector space E and Y an almost convex dense subset of X . Let $T : X \multimap E$ be a multimap such that*

- (1) $T^-(z)$ is open for each $z \in E$; and
- (2) $T(y)$ is convex for for each $y \in Y$.

If there is a precompact subset K of X such that $T(y) \cap K \neq \emptyset$ for each $y \in Y$, then for any convex neighbourhood U of the origin 0 of E , there exists a point $x_U \in Y$ such that $T(x_U) \cap (x_U + U) \neq \emptyset$.

PROOF. Since T is lower semicontinuous, Corollary 6 follows immediately from Theorem 3. \square

In case $X = Y$, Corollary 6 reduces to the following:

COROLLARY 7. *Let X be a convex subset of a Hausdorff topological vector space E , and $T : X \multimap X$ be a multimap such that*

- (1) $T(x)$ is nonempty and convex for each $x \in X$,
- (2) $T^{-1}(y)$ is open for each $y \in X$, and
- (3) $T(X)$ is contained in a compact subset K of X .

Then for any convex neighbourhood U of the origin 0 of E , there exists a point $x_U \in X$ such that $T(x_U) \cap (x_U + U) \neq \emptyset$.

Ben-El-Mechaiekh [4], [5] obtained that, if E is further assumed to be locally convex in Corollary 7, then T has a fixed point; and conjectured that, under the

hypotheses of Corollary 7, T would have a fixed point. This conjecture is not resolved yet; for partial solutions, see [12]. However, Corollary 7 is a new partial solution.

REFERENCES

- [1] P. S. ALEXANDROFF, *Combinatorial Topology*, OGIZ, Moscow–Leningrad, 1947. (Russian)
- [2] P. ALEXANDROFF UND H. HOPF, *Topologie I*, Springer, Berlin–Heidelberg–New York, 1935.
- [3] P. ALEXANDROFF AND B. PASYNKOFF, *Elementary proof of the essentiality of the identity mapping of a simplex*, Uspekhi Mat. Nauk **12** (1957), 175–179. (Russian)
- [4] H. BEN-EL-MECHAIEKH, *The coincidence problem for compositions of set-valued maps*, Bull. Austral. Math. Soc. **41** (1990), 421–434.
- [5] ———, *Fixed points for compact set-valued maps*, Questions Answers Gen. Topology **10** (1992), 153–156.
- [6] KY FAN, *A generalization of Tychonoff's fixed point theorem*, Math. Ann. **142** (1961), 305–310.
- [7] ———, *Covering properties of convex sets and fixed point theorems in topological vector spaces*, Symposium on Infinite Dimensional Topology, Ann. of Math. Studies, No. 69, Princeton Univ. Press, 1972, pp. 79–92.
- [8] C. J. HIMMELBERG, *Fixed points of compact multifunctions*, J. Math. Anal. Appl. **38** (1972), 205–207.
- [9] W. K. KIM, *Some applications of the Kakutani fixed point theorem*, J. Math. Anal. Appl. **121** (1987), 119–122.
- [10] B. KNASTER, K. KURATOWSKI UND S. MAZURKIEWICZ, *Ein Beweis des Fixpunktsatzes für n -Dimensionale Simplexe*, Fund. Math. **14** (1929), 132–137.
- [11] M. LASSONDE, *Sur le principe KKM*, C. R. Acad. Sci. Paris Sér. I **310** (1990), 573–576.
- [12] S. PARK, *Remarks on fixed point problems of Bel-El-Mechaiekh*, Nonlinear Analysis and Convex Analysis, (Proc. NACA'98, Niigata, Japan, July 28-31, 1998) (W. Takahashi and T. Tanaka, eds.), World Sci., Singapore, 1999, pp. 79–86.
- [13] S. PARK AND K. S. JEONG, *Fixed point and non-retract theorems – classical circular tours*, Taiwan. J. Math. **5** (to appear).
- [14] S. PARK AND D. H. TAN, *Remarks on the Schauder-Tychonoff fixed point theorem*, Vietnam J. Math. **28**, 127–132.
- [15] ———, *Remarks on the Himmelberg-Idzik fixed point theorem*, Acta Math. Vietnam. (to appear).
- [16] M.-H. SHIH AND K.-K. TAN, *Covering theorems of convex sets related to fixed-point theorems*, Nonlinear and Convex Analysis (Proc. in Honor of Ky Fan), Marcel Dekker, Inc., New York–Basel, 1987, pp. 235–244.

Manuscript received November 10, 1999

SEHIE PARK
 Department of Mathematics
 Seoul National University
 Seoul 151-742, KOREA

E-mail address: shpark@math.snu.ac.kr

TMNA : VOLUME 16 – 2000 – N° 1