# Allelopathic effects of sunflower water extract integrated with affinity herbicide on weed control and wheat yield

# Muhammad Tahir Jatoi<sup>1,\*</sup>, Samreen Riaz Ahmed<sup>2</sup>, Altaf Hussain Lahori<sup>3</sup>, Volodymyr Sydorenko<sup>4,\*</sup>, Ambreen Afzal<sup>5</sup>, Muhammad Nawaz Kandhro<sup>1</sup>, Sarwat Haider Janveri<sup>1</sup>, Haider Bux Jatoi<sup>6</sup>, Zaheer Ahmed Jatoi<sup>6,7</sup>, Sajjad Ali Panhwar<sup>7,8</sup>, Andrii Pruskyi<sup>4</sup>

<sup>1</sup>Department of Agronomy, Sindh Agriculture University, Tandojam, 70060 Pakistan
<sup>2</sup>Department of English, Sindh Madressatul Islam University, Karachi, 74000 Pakistan
<sup>3</sup>Department of Environmental Science, Sindh Madressatul Islam University, Karachi, 74000 Pakistan
<sup>4</sup>Institute of Public Administration and Research in Civil Protection, Kyiv, Ukraine
<sup>5</sup>National Institute of Maritime Affairs, Bahria University, Karachi Campus, Pakistan
<sup>6</sup>Department of Soil Science, Sindh Agriculture University, Tandojam, 70060 Pakistan
<sup>7</sup>Department of Plant Breeding and Genetic, Sindh Agriculture University, Tandojam, 70060 Pakistan

\*corresponding authors e-mail: Volodymyr Sydorenko (V.S.), 2022.12.gl@gmail.com Muhammad Tahir Jatoi (M.T.J.), jatoitahir@gmail.com

Received: 3 May 2022 / Accepted: 8 June 2022

**Abstract.** The aim of present study was to investigate the allelopathic impact of sunflower water extract integrated with affinity herbicide on weed management and production of wheat variety TD-1 under field condition. The trial was carried out in a three replicated and the average plot size was kept at  $3x5m (15 m^2)$ . The obtained data showed that the all the treatments considerably (P<0.05) minimized weed density, weed fresh weight, weed dry weight and weed control up to 90.33, 89.31, 89.61, 90.33% with application of sunflower water extract (1:10) @ 20 L ha<sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha<sup>-1</sup> (One spray at 25 DAS) than control treatment. The application of sunflower water extract (1:10) @ 20 L ha<sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha<sup>-1</sup> (One spray at 25 DAS) significantly increased number of tillers (32.34%), plant height (14.59%), spike length (27.89%), grains spike<sup>-1</sup> (16.80%), grain weight plant<sup>-1</sup> (47%), seed index (19.60%) and grain yield (51%) of wheat as compared with control. Overall results indicated that the sunflower water extract integrated with herbicide can be successfully used to control the narrow and broad leaves in wheat crop.

Key words: allelopathic, sunflower water extract, affinity, agronomic parameters, wheat, productivity.

**List of abbreviations and designations:** DAS – days after sowing, WP – wettable power, grain weight  $plant^{-1}$  – grain weight per plant, grains spike<sup>-1</sup> – grains per spike, @ – Mean at the rate or application rate.

# 1. Introduction

Wheat is a staple food crop and contributing over one-third of the globe population. Approximately, 36% population

of Pakistan depends on this stable food crop and its total cropping area is occupied 30% as compared to other major crops (Abdul et al., 2012; Li et al., 2022). More than 50% of the agricultural land throughout the country is cultivated with

wheat crop during its cropping season (Hussain et al., 2013; Matloob et al., 2020). However, impure seed, irrigation water, wind are the main factors of increasing unwanted weeds cause massive yield losses in wheat and other field crops (Singh et al., 2005; Akbar et al., 2011). On the one side, weed are the main source of animal feed, soil cover, biochar preparation and compost technology, but on the other hand increasing insect population, crop disease and minimize over crop production. In the former practice, the farmers were control weed by using physical, mechanical methods, but these methods were found to time consuming and not applicable for large scale. Since few decades, the application of herbicide had been used at large scale to control unwanted plant from wheat crops (Rashid et al., 2022). The excess application of herbicide for controlling of unwanted weed may cause harm to soil health, water pollution and also affect major crop growth and development (Farooq et al., 2011; Vambol et al., 2020). To tackle this unfriendly approach, the best solution is that to use of organic allelopathy manipulation for crop improvement and ecofriendly safety (Singh et al., 2003). Furthermore, Memon and Bhatti (2003) reported that twenty-four weed species infesting the wheat crop in the wheat fields. Fahad et al. (2015) revealed that the dense population of unwanted weed plants may possibly reduce 30-35% major crop growth parameters and yield, because of dense population unwanted plant include narrow leaf and/or broad leaf may equal share with major crops in water, nutrients, sunlight, density etc. Weeds are responsible for the decline in crop yield losses caused by weeds can be as high as 24% of yield compared with 16.4 and 11.2% for disease and pest, respectively (An et al., 2001). The existing weed control methods are either expensive or hazardous. Sorghum allelopathy for weed management is the safe use of agro-chemicals thus, bioherbicides (allelochemicals) seem to be environment friendly (Naeem et al., 2022). Therefore, alternative strategies against weed must be developed. Allelopathy can be considered as environmental friendly approach for mitigating unwanted weeds and helpful to reduce cost of herbicides and save environmental pollution (Albuquerque et al., 2011). In addition, Bhadoria (2011) stated that the manufacturing of allele-chemicals from several crops biomass has been well known and their potential for weed management. Most of the synthesis herbicides products contents mono or multi toxic chemicals that can be used for control narrow or broad leaves control. The researchers developed some organic herbicides that not only use to control unwanted plants, but also has great importance in environmental protection and socioeconomic point of view. Xuan et al. (2005) reported that the integrated weed management by application of allelopathy could be used as an eco-friendly tool in weed management. Studies have shown an excessive probable of allelopathy for weed management especially in wheat crop (Macias et al.,

2007). It can be considered as a best alternative option as compared to synthetic herbicides to control weeds (Naeem et al., 2018). Allelopathy is a phenomenon in which one plant hinders growth of other plants through discharge of allelechemicals as result due to insufficient food and chemical hang-up (Farooq et al., 2020). Sunflower is recognized to dynamically influence the growth of surrounding plants due to its great allelopathic potential and > 200 natural allelopathic complexes have been isolated from sunflower cultivars. Most of these known allelochemicals affect seed germination (Kamal & Bano, 2008). In Pakistan, many plant species have shown allelopathic influences that could be effectively applied for management of unwanted plant for safe and maximum production (Khan et al., 2011). The negative and positive effects of chemical compounds produced from microorganisms, fungi, viruses and plants manipulate agricultural and biological ecosystems (Martines et al., 2009). The main objectives of this research work were: 1) to examine the allelopathic impact of dissimilar proportion of sunflower water extract on weed management and yield enhancement of wheat; 2) evaluate the effect of studied treatments on narrow leaf and broad leaf herbicides on weed management and yield enhancement of wheat; and 3) find out the most optimum integration of sunflower water extract and herbicides for effective weed management and maximum yield of wheat.

# 2. Materials and methods

**Experimental design:** The field experiment was conducted at "Students Farm", Department of Agronomy Sindh Agriculture University Tandojam aiming to evaluate the allelopathic effect of sunflower water extract and herbicides on weed management and yield enhancement of wheat. The map location of study is indicated in (Fig. 1). The seed of wheat variety TD-1 was used throughout the experiment. The experiment was laid out in a three replicated randomized complete block design (RCBD). The net plot size kept was  $3x5m (15 m^2)$ . The treatment design and dosage is represented in Table 1.

## Preparation of sunflower water extract

The mature dry plants of sunflower were chopped in small pieces and chopped material was soaked in distilled water for 24 hours at 30°C in a ration of 1:10 (w/v). Furthermore, the extract was filtered by using sieve at finally boiled at 100°C up to 20 minutes.

## Application herbicide and sunflower water extract

Except control block the application herbicides and sunflower water extract was sprayed on studied blocks to



Figure 1. The area in which the research was carried out

Code	Treatments description
T1	Weedy check (Untreated).
T2	Sunflower water extract (1:10) @ 20 L ha <sup>-1</sup> (One spray at 25 DAS).
T3	Sunflower water extract (1:10) @ 20 L ha <sup>-1</sup> (Two sprays at 25+50 DAS).
T4	Sunflower water extract (1:10) @ 30 L ha <sup>-1</sup> (One spray at 25 DAS).
T5	Sunflower water extract (1:10) @ 30 L ha <sup>-1</sup> (Two sprays at 25+50 DAS)
T6	Affinity 50 WP @ 1.5 kg ha <sup>-1</sup> (One spray at 25 DAS).
T7	Sunflower water extract (1:10) @ 20 L ha <sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha <sup>-1</sup> (One spray at 25 DAS)
T8	Sunflower water extract (1:10) @ 30 L ha <sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha <sup>-1</sup> (One spray at 25 DAS)

assess the efficacy of herbicide and sunflower water extract for weed management.

**Land preparation:** The land was prepared by two dry plowings followed by precision land levelling. After soaking doze, when soil reached proper moisture level, two plowings with cultivator plow were done to achieve the fine seedbed.

**Sowing time and method:** In the present work, TD-1 variety was chosen and the recommended dose 125 kg ha<sup>-1</sup> was sown by hand drilling method, and the distance among rows to row was kept at 22.5cm.

**Irrigation and fertilizer application:** The 1<sup>st</sup> irrigation was applied to wheat plants 21 days after sowing,  $2^{nd}$  irrigation at ear head formation and  $3^{rd}$  I irrigation was applied at milky stage respectively. The recommended dose of nitrogen 120 kg ha<sup>-1</sup> in the form of urea was applied with  $1^{st}$  and  $2^{nd}$  irrigation, and phosphorus 75 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in the form of DAP was applied at soil preparation.

## **Crop harvesting**

The matured 3 wheat plants were randomly selected from each plot as per 3 replications. After harvesting ear head/spike was separated from straw during threshing by hand, the studied parameters were measured as per experimental design.

## Observations recorded (Weed observations)

Weed flora: In the present study weed density  $(m^{-2})$ , weed fresh weight (g m<sup>-2</sup>), weed dry weight (g m<sup>-2</sup>), and weed control (%) were measured.

**Wheat observations:** The number of tillers (m<sup>-2</sup>), plant height (cm), spike length (cm), grains spike<sup>-1</sup>, grain weight plant<sup>-1</sup> (g), seed index (1000 grain weight, g), and grain yield (kg ha<sup>-1</sup>) were measured as per standard way.

#### Method for recording observations

Weed flora of wheat: All the weed species infesting the experimental area were recorded and their local names,

English names, botanical names and families are mentioned in (Table 2).

Weed density ( $m^{-2}$ ): The weeds were counted at 60 DAS by using wooden frame of one square meter in all plots of each treatment at one location and calculated as ( $m^{-2}$ ). The following formula was applied:

Weed 
$$\frac{1}{\text{density}(m^2)} = \frac{\text{Total number of weeds for given treatment}}{\text{Number of replications}}$$
 (1)

**Weed control (%):** Weed control (%) was calculated by using the following formula.

$$\frac{\text{Weed}_{\text{control}}}{(\%)} = \frac{\text{Weedy Check density} - \text{Given treatment density}}{\text{Weedy Check density}} \cdot 100 \ (2)$$

Weed fresh weight (g m<sup>-2</sup>): Weeds fresh weight was taken from randomly selected areas of one-meter square at one stage at 60 DAS and recording the fresh weight.

**Weed dry weight (g m<sup>-2</sup>):** weeds dry weight the samples taken for fresh weight were dried at 70°C for 72 hours.

**Number of tillers (m<sup>-2</sup>):** Total tillers (m<sup>-2</sup>) in each plot from three different locations was counted at the time of maturity and averaged.

**Plant height (cm):** Plant height was recorded at maturity of the crop in randomly selected three plants using measuring tape from bottom to tip of the spike in centimetres.

**Spike length (cm):** The length of all the spikes in randomly selected plants will be measured in centimetres with measuring tape and average will be worked out.

Number of grains spike<sup>-1</sup>: The number of grains in each spike of the randomly selected plants was counted at the crop maturity and average will be worked out.

**Grain weight plant**<sup>-1</sup> (g): The all spikes of the plant were threshed and measured the yield in grams through electric weight balance in laboratory.

Table 2. Diversity of weed flora infesting the experiment	tal field of wheat (WFO, 2022)
---	--------------------------------

Local Name	English Name	Botanical Name	Family					
Narrow leaf weeds								
Dumbi siti	Phalaris minor Retz.	Poaceae						
Jangli javi	Wild oat	Avena fatua L.	Poaceae					
Kabah	Purple nutsedge	Cyperus rotundus L.	Cyperaceae					
Chhabar	Bermuda grass	<i>Cynodon dactylon</i> (L.)Pers.	Poaceae					
Broadleaf weeds								
Sufaid Sinjh	White sweet clover	Melilotus alba Medik.	Fabaceae					
Peeli Sinjgh	Indian clover	Melilotus indica (L.)All.	Fabaceae					
Jhil	Common Lamb'squarter	Chenopodium album L.	Amaranthaceae					
Chhati kherol	Sunspurge	Euphorbia helioscopia L.	Euphorbiaceae					
Naro	Field Bind weed	Covolvulas arvensis L.	Convolvulaceae					
Jangli palak	Dock	Rumex dentatus L.	Polygonaceae					
Bili booti	Blue pimpernel	Anagallis arvensis L.	Primulaceae					

Seed index (1000 grain weight, g): For recording seed index value, 1000 grains from all the randomly selected plants in each plot was collected at random and weighed in grams.

**Grain yield (kg ha<sup>-1</sup>):** The grain yield (kg ha<sup>-1</sup>) was calculated by using the following formula.

Grain yield (kg·ha<sup>-1</sup>) = 
$$\frac{\text{Grain yield plot}^{-1}(\text{kg}) \cdot 10\ 000}{\text{Area of plot}(\text{m}^2)}$$
(3)

## **Geospatial techniques**

Geospatial techniques are notable in the ex-situ condition for managing, monitoring the huge data set easily and resourcefully. In the present work, high-resolution satellite imaginary data have been extracted from Google earth then geo-referencing techniques have been used for image restructuring. For extraction of the area of interest, a mask tool has been used. The editor tool has been used for the digitization of image data.

#### Statistical analysis

The obtained data was subjected to ANOVA techniqne using MSTAT-C computer software. The LSD test was applied to compare treatment means superiroity, where necessary. All graphs were prepared by using Origin Pro. 16. The pearson correlation coefficients between weed density, weed fresh weight, weed dry weight, weed control, tillers, plant height, spike length, and grains per spike, grain weight, seed index and grain yield were also performed using IBM SPSS Statistics for Windows, Version 21.0.

# 3. Results and discussion

As shown in (Fig. 2a) the obtained results revealed that the all experimental treatments were found to be highly effective to reduce the weed density (m<sup>-2</sup>) as compared to control treatment. While, the maximum reduction of weed density was 90.33% with application of Sunflower water extract



Figure 2. Weed density (a), weed fresh weight (b), weed dry weight (c), and weed control (d) as affected by allelopathic effect of sunflower water extract and herbicides in wheat



**Figure 3.** Tillers (a), plant height (b), spike length (c), and grains per spike (d) as affected by allelopathic effect of sunflower water extract and herbicides in wheat

(1:10) @ 20 L ha<sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha<sup>-1</sup> (One spray at 25 DAS), rather than control treatment. Furthermore, the greatest weed fresh weight (g m<sup>-2</sup>) was observed by 89.31% with application of Sunflower water extract (1:10) @ 20 L ha<sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha<sup>-1</sup> (One spray at 25 DAS), as compared to control treatment (Fig. 2b). As compared the control, the weed dry weight (g m<sup>-2</sup>) was evidently reduced up to 89.61% with application of Sunflower water extract (1:10) @ 20 L ha<sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha<sup>-1</sup> (One spray at 25 DAS) (Fig. 2c). The maximum weed control (%) was found to be 90.33% with application of Sunflower water extract (1:10) @ 20 L ha<sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha<sup>-1</sup> (One spray at 25 DAS) rather than control treatment (Fig. 2d). Above results indicated the application of different concentrations of sunflower water extract and herbicide have highly potential to reduce weed density, weed fresh weight, weed dry weight, and weed control in wheat crop. Indeed, allelopathy is an environment friendly technique in controlling weeds

and helpful in reducing cost of herbicide. Furthermore, allelopathic plants may also be considered a potential source of new molecules with herbicidal action for the chemical industry, the necessity of which is due to the emergence of resistant weeds to older synthetic molecules. Moreover, allelopathic potential in crop plants can be successfully utilized for weed management. Certain crops restrain growth of some weed species while phytotoxins releasing from their residues inhibit seed germination of weeds. Indeed, allelechemicals from several crops have been identified and their activity for weed management has also been established (Bhadoria, 2011). Research in allelopathy focuses on isolating, identifying and quantifying specific active allelochemicals. Once these substances are identified and characterized they can be used either as natural herbicides or as models for developing new and environment friendly herbicides. Studies have shown a great potential of allelopathy for weeds control in wheat. It is the best alternatives to the synthetic herbicides to control weeds (Bhowmik & Inderjit, 2003; Jabran et al., 2008). These outcomes are supported by Naeem et al. (2016), who stated that the lessening in weed dry weight with sorghum and sunflower extracts by foliar spraying.

The obtained experimental data in showed that the studied treatments evidently increased the number of tillers, plant height, spike length, and grains per spike of wheat plant as compared to control treatment. The maximum tillers  $(m^{-2})$  of wheat was received by 32.34% with application of Sunflower water extract (1:10) @ 20 L ha<sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha<sup>-1</sup> (One spray at 25 DAS) rather than control treatment (Fig. 3a). Furthermore, the greatest plant height (cm) of wheat was 14.59% with Sunflower water extract (1:10) @ 20 L ha<sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha<sup>-1</sup> (One spray at 25 DAS) as compared to control treatment (Fig. 3b). Similarly, the highest Spike length (cm) of wheat crop was noted 27.89% with application of Sunflower water extract (1:10) @ 20 L ha<sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha<sup>-1</sup> (One spray at 25 DAS) than control (Fig. 3c). Also, the maximum grains spike<sup>-1</sup> of wheat was 16.80% with



**Figure 4.** Grain weight (a), seed index (b), and grain yield (c) as affected by allelopathic effect of sunflower water extract and herbicides in wheat

application of Sunflower water extract (1:10) @ 20 L ha<sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha<sup>-1</sup> (One spray at 25 DAS) as compared with control treatment (Fig. 3d).The results of present experiment showed that all the levels of sunflower water extract caused significant inhibition on weed management and yield enhancement of wheat. Yields of some crops following sunflower are lower than normal, possibly because of inadequate nutrition and chemical inhibition. Sunflower is often grown when rainfall is marginal and depletion of soil moisture by sunflower may be a factor, although, this remains unproven (Kamal & Bano, 2008).

The experiment results revealed that the grain weight per plant, seed index and grain yield of wheat crop was potentially increased with application of different concentrations of sunflower water extract and herbicide. The maximum grain weight plant<sup>-1</sup> (g), seed index 1000 grain weight (g) and grain yield (kg ha<sup>-1</sup>) was recorded up to 47%, 19.60% and 51% with application of Sunflower water extract (1:10) @ 20 L ha<sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha<sup>-1</sup> (One spray at 25 DAS) than control treatment (Fig. 4a, b, c). The effectiveness of bioherbicide depends on many factors as soil type, moisture content, organic matter, prevailing temperature, etc., (Matloob et al., 2015). Our results are in line with Naeem et al. (2018), who observed that the yield of wheat was significantly increased and weeds were reduced with application of sorghum, sunflower and mulberry extract. Appropriate weed controlling is likely to raise wheat yield by more than one million ton in Pakistan (Marwat et al., 2011).

#### Pearson correlation analysis

Pearson correlation analysis was performed to assess the relationship between different weed density, weed fresh weight, weed dry weight, weed control, tillers, plant height, spike length, and grains per spike, grain weight, seed index and grain yield (Table 3). The weed fresh weight had highly significant correlation with weed density.

The weed dry weight was highly significant correlation with weed density and weed fresh weight. Weed control had negatively highly significant correlation with weed density, weed fresh weight and weed dry weight. Tillers were negatively correlated with weed density, weed fresh weight, weed dry weight, but positive correlation with weed control. Plant height had negative correlation with weed density, weed fresh weight, weed dry weight, but positive correlation with weed control and tillers. Spike length had negative correlation with weed density, weed fresh weight, weed dry weight, but positive correlation with weed control, tillers and plant height. Grains per spike were negatively correlated with weed density, weed fresh weight, weed dry weight, whereas positive correlation with weed control,

Traits	WD	WFW	WDW	WC	Т	PH	SL	GS	GW	SI	GY
WD	1										
WFW	.977**	1									
WDW	.966**	.988**	1								
WC	-1.00**	977**	966**	1							
Т	977**	988**	982**	.977**	1						
PH	956**	973**	983**	.956**	.985**	1					
SL	989**	983**	984**	.989**	.983**	.980**	1				
GS	975**	962**	980**	.975**	.976**	.978**	.988**	1			
GW	977**	990**	988**	.977**	.970**	.962**	.989**	.970**	1		
SI	969**	961**	983**	.969**	.968**	.981**	.987**	.997**	.972**	1	
GY	995**	989**	983**	.994**	.990**	.969**	.993**	.983**	.986**	.976**	1

Table 3. Pearson correlation coefficient between studied parameters

Note: Correlation is significant at \*\*p<0.01; WD = weed density; WFW = weed fresh weight; WDW= weed dry weight; WC= weed control; T= tillers; PH= plant height; SL= spike length; GS= grains spike<sup>-1</sup>; GW= grain weight; SI= seed index and GY= grain yield.

tillers, plant height and spike length. Grain weight found to be negative correlated with weed density, weed fresh weight, weed dry weight, however positive correlation with weed control, tillers, plant height, spike length and grains spike<sup>-1</sup>. Seed index was negatively correlated with weed density, weed fresh weight, weed dry weight, but positively correlated with weed control, tillers, plant height, spike length, grains spike<sup>-1</sup>and grain weight. Grain yield had positive correlation with weed density, weed fresh weight, weed dry weight, but positively correlated with weed control, tillers, plant height, spike length, grains spike<sup>-1</sup>, grain weight and seed index.

# 4. Conclusions

The results concluded that the combination of sunflower water extract (1:10) @ 30 L ha<sup>-1</sup> (One spray at 25 DAS) + Affinity 50 WP @ 0.75 kg ha<sup>-1</sup> soil was found highly allelopathic and resulted in lowest growth and related traits of wheat. However, allelopathic potential of this weed could be exploited as effective, cheap and eco-friendly strategy for weed management. In future, long-term ex-situ studied are needed to understand the mechanism underlying soil health, microbial biomass, soil enzymatic activity, physiology of crop, crop production and weed management under. Furthermore, the herbicides contaminated plants can be convert for energy purpose and to made smart biochar as soil amendment for waste management and environmental clean-up.

### References

Abdul K., Amar M. & Yasir R., 2012, Bio-economic and qualitative impact of reduced herbicide use in direct seeded fine rice through multipurpose tree water extracts. Chilean Journal of Agricultural Research 72(3): 350–357.

- Akbar N., Ullah, E. Jabran K. & Ali M.A., 2011, Weed management improves yield and quality of direct seeded rice. Australian Journal of Crop Science 5(6): 688–694.
- Albuquerque M., Santos R., Lima L., Filho P., Nogueira R., Camara C. & Ramos, A., 2011, Allelopathy, an alternative tool to improve cropping systems. A review. Agronomy for Sustainable Development 31: 379–395.
- An M., Pratley J.E. & Higa T., 2001, Phytotoxicity of Vulpia Residues: III. Biological Activity of Identified Allelochemicals from Vulpia myuros. Journal of Chemical Ecology 27: 383–394.
- Bhadoria P.B.S., 2011, Allelopathy a natural way towards weed management. American Journal of Agricultural Economics 1: 7–20.
- Bhowmik P.C. & Inderjit, 2003, Challenges and opportunities in implementing allelopathy for natural weed management. Crop Protection 22: 661–671.
- Fahad S., Hussain S., Chauhan B.S., Saud S., Wu C., Hassan S., Tanveer M., Jan A., Huang J.L., 2015, Weed growth and crop yield loss in wheat as influenced by row spacing and weed emergence times. Crop Protection 71: 101–108.
- Farooq M., Jabran K. Cheema Z.A., Wahid A. & Siddique K.H.M., 2011, The role of allelopathy in agricultural pest management. Pest Management Science 67(5): 493–506.
- Farooq M., Khan I., Nawaz A., Cheema M.A., Siddique K.H.M., 2020, Using sorghum to suppress weeds in autumn planted maize. Crop Protection 133: 105162.
- Hussain Z., Marwat K.B., Samad F.M.A. & Ali K., 2013, Evaluation of various herbicides and their combinations for weed control in wheat crop. Pakistan Journal of Botany 45(1): 55–59.

- Jabran K., Cheema Z.A., Farooq M., Basra S.M.A., Hussain M. & Rehman H., 2008, Tank mixing of allelopathic crop water extracts with pendimethalin helps in the management of weeds in canola (*Brassica napus*) field. International Journal of Agricultural and Biological 10(3): 293–296.
- Kamal J. & A. Bano., 2008, Effects of sunflower (*Helianthus annuus* L.) extracts on wheat (*Triticum aestivum* L.) and physicochemical characteristics of soil. African Journal of Biotechnology 7 (22): 4130–4135.
- Khan A.A., Awan I.U., Mansoor M., Khan E.A., Khakwani A.A. & Baloch M.S., 2011, Efficacy and economic analysis of plant aqueous exudates on physiological parameters of wheat. Pakistan Journal of Weed Science Research 17(4): 417–424.
- Li L., Wang B., Feng P., Li Liu D., He Q., Zhang Y. & Yu Q., 2022, Developing machine learning models with multi-source environmental data to predict wheat yield in China. Computers and Electronics in Agriculture 194: 106790.
- Macias F.A., Molinillo J.M., Varela R.M. & Galindo, J.C., 2007. Allelopathy – a natural alternative for weed control. Pest Management Science: Formerly Pesticide Science, 63(4): 327–348.
- Martines I.P., Kojouharov H.V. & Grover J.P., 2009. A chemostat model of resource competition and allelopathy. Applied Mathematics and Computation 215: 573582.
- Marwat K.B., Khan M.A., Hashim S., Nawab K. & Khattak A.M., 2011, Integrated weed management in wheat. Pakistan Journal of Botany (43): 625–633.
- Matloob A., Khaliq A., Tanveer A., Hussain S., Aslam F. & Chauhan B.S., 2015, Weed dynamics as influenced by tillage system, sowing time and weed competition duration in dry-seeded rice. Crop Protection 71: 25–38.
- Matloob A., Safdar M.E., Abbas T., Aslam F., Khaliq A., Tanveer A., Rehman A. & Chadhar A.R., 2020, Challenges and prospects for weed management in Pakistan: A review. Crop Protection 134: 104724.
- Memon R.A. & Bhatti G.R., 2003, Weed diversity of wheat crop in Khaipur district, Sindh (Pakistan). Pakistan Journal of Weed Science Research 9: 99–103.

- Naeem M., Cheema Z.A, Ihsan M.Z, Hussain Y, Mazari A. & Abbas H.T., 2018, Allelopathic effects of different plant water extracts on yield and weeds of wheat. Planta Daninha 36: 018177840.
- Naeem M., Farooq S. & Hussain M., 2022, The Impact of Different Weed Management Systems on Weed Flora and Dry Biomass Production of Barley Grown under Various Barley-Based Cropping Systems. Plants 11(6): 718.
- Naeem M., Mahmood A., Ihsan M.Z., Daur I., Hussain S., Aslam Z. & Zamanan S.A., 2016, *Trianthema portulacastrum* and *Cyperus rotundus* interference in maize and application of allelopathic crop extracts for their effective management. Planta Daninha 34(2): 209–218.
- Rashid H.U., Khan A., Hassan G., Munsif F., Tahir N., Zamin M. & Shehzad N., 2022, Integration of some allelopathic species for weed management in spring planted hybrid maize under different tillage regimes. Journal of Animal and Plant Sciences – JAPS 32(1): 114–126.
- Singh H.P., Batish D.R. & Kohli R.K., 2003, Allelopathic interactions and allelochemicals: new possibilities for sustainable weed management. Critical Reviews in Plant Sciences 22(3–4): 239–311.
- Singh H.P., Batish D.R., Setia N. & Kohli R.K., 2005, Herbicidal activity of volatile oils from *Eucalyptus citriodora* against *Parthenium hysterophorus*. Annals of Applied Biology 146(1): 89–94.
- Vambol S., Khan N.A., Khan A.H., Kiriyenko M. & Bilotserkivska N., 2020, Developed jet-centrifugal spray devices: experimental testing to establish the possibility of their application in plants spraying technologies. Journal of Achievements in Materials and Manufacturing Engineering 102(1): 30–41. DOI: 10.5604/01.3001.0014.6326
- WFO, 2022, World Flora Online. Published on the Internet [Accessed on: 5 Jun 2022].
- Xuan T.D., Shinkichi T., Khanh T.D. & Chung I.M., 2005, Biological control of weeds and plant pathogens in paddy rice by exploiting plant allelopathy: An Overview. Crop Protection 24(3): 197–206.