

Lichen diversity in the managed forests of the Karnieszewice Forest Division and its surroundings (N Poland)

Wiesław Fałtynowicz¹, Agnieszka Kowalewska², Rafał Szymczyk³, Martin Kukwa⁴, Edyta Adamska⁵, Paweł Czarnota⁶, Dariusz Kubiak⁷, Katarzyna Pietrzykowska-Urban¹

¹ Department of Botany, Institute of Environmental Biology, Wrocław University,
Kanonia 6/8, 50-328 Wrocław, Poland; e-mail: wieslaw.faltynowicz@uni.wroc.pl

² Pomorski Group of Landscape Parks in Słupsk, Gdańsk Division – Trójmiejski Landscape Park,
Polanki 51, 80-308 Gdańsk, Poland

³ Environmental Survey Laboratory Ekoprojekt,
Nowica 24, 14-405 Wilczęta, Poland

⁴ Department of Plant Taxonomy and Nature Protection, Gdańsk University,
Wita Stwosza 59, 80-308 Gdańsk, Poland

⁵ Nicolaus Copernicus University, Faculty of Biology and Environment Protection,
Chair of Geobotany and Landscape Planning
Lwowska 1, 87-100 Toruń, Poland

⁶ Department of Agroecology, Rzeszów University,
Ćwiklińskiej 1A, 35-601 Rzeszów, Poland

⁷ Department of Mycology, Warmia and Mazury University in Olsztyn,
Oczapowskiego 1A, 10-719 Olsztyn, Poland

Abstract. The lichen biota of the Karnieszewice Forest Division (N Poland) is presented. Despite it is predominantly a strongly managed woodland area, 270 lichen species were found there including many rare species for Poland, as well as for European Lowland. Near 20% of the whole lichen biota are considered to be threatened in the country (categories CR, EN, VU), and 34 species are protected by law in Poland. *Agonimia flabelliformis* is reported for the second time from Polish lowlands.

Key words: biodiversity, lichenized fungi, endangered species, Pomerania, Ascomycota, Basidiomycota, rare species.

1. Introduction

Sensitivity of lichens against changes in habitat conditions is known and widely described in literature, beginning with the classical work of Nylander (1866) and thousands books and articles, (e.g. Hawksworth 1971; Herzig & Urech 1991; Fałtynowicz 1995; van Herk 2001; Zalewska 2012; Adamska 2014; Jóźwiak 2014). Also impact of forest management on lichen biota have been mentioned in many publications. In 1934 dr. F. Krawiec wrote: „...Flora

on trees in some cases better reflect forest state than vascular plants...in lichen flora distinctly is seen impact of cutting down of forest” (Krawiec 1934), and prof. J. Motyka observed that „List of lichen species is one of the best features of plant communities primarity” (Motyka 1934). Problems of the negative impact of forestry culture on lichens has been widely discussed, e.g. by Czyżewska (1976), Cieśliński (2003), Łubek (2007) and Zalewska (2012), however, despite of the antropogenic changes in forests, in many such areas, particularly in northern Po-

land, numerous stenotopic or forest relict lichens have been more and more often found (e.g., Cieśliński 2003; Zalewska et al. 2004a, b; Wieczorek 2005; Zalewska 2012; Kubiak et al. 2014). It may be a proof for the improvement of habitat conditions in forests, but it should be further investigated in the future.

The aim of this work was to know a lichen diversity in managed lowland forests and to compare the list of species with results of similar investigations made in other lowland woodlands in Poland including the old and long-term preserved forest areas.

2. Study area

The area of Karnieszewice Forest Division has very rich and diversified geomorphology. Ground moraines dominate, but almost all geomorphological forms, typical for the zone of the last glaciation in northern Poland (e.g., frontal moraines, pravalleys, narrow erosion valleys with steep slopes, numerous spring areas) are present here. Landscape is enriched by seashore with mosaics of dunes, sandbars, shore of large lakes and peatbogs.

Such geomorphology gives a great diversity of habitats for plant communities. Coniferous habitats dominate in the forests (nearly 45%), however other forest associations are also known from this area: *Melico-Fagetum*, *Luzulo pilosae-Fagetum*, *Stellario-Carpinetum*, *Ribeso nigri-Alnetum*, and small areas covered by phytocoenoses of *Betuletum pubescantis*, *Circaeо-Alnetum* and *Betulo-Quercetum*. Non-forest associations important for the occurrence of lichens are mainly *Helichryso-Jasionetum* and heaths in coniferous complexes.

Tree stands in the Karnieszewice Forest Division possess relatively high average of age (65 years). The dominant tree species is *Pinus sylvestris* (52%), and a great share has also *Picea abies* (9%). Deciduous trees are almost 40% of forests, with dominating *Fagus sylvatica* (14%), *Betula* spp. (10,5%), *Alnus glutinosa* (6%) and *Quercus* spp. (5,5%). Trees and shrubs growing along roadsides are also very important for lichens, with the most frequent: *Acer platanoides*, *A. pseudoplatanus*, *Carpinus betulus*, *Fraxinus excelsior*, *Corylus avellana*, *Larix* spp., *Padus avium*, *Populus* spp., *Sorbus aucuparia* and *Tilia cordata*.

More than a half of the forests of Karnieszewice Forest Division (10 560 ha) is protected in different ways: Nature 2000 areas, nature reserves, areas of protected landscape, etc. Since 2011 the whole Karnieszewice Forest Division is the part of Forest Promotion Complex „Lasy Środkowopomorskie”.

The study area includes also areas not belonging to the Karnieszewice Forest Division – agriculture and rural areas. They are places with rich and completely different lichen biota. Particularly interesting habitats were roadside

trees (most often maple, sycamore and lime trees) and trees around old churches.

3. Materials and methods

The article is based on observations and collections made during field studies in 2013 (localities nos 1–23; see Kukwa et al. 2013) and 2014 (localities nos 24–53 and 60–69), and on published data (Erichsen 1936, 1940; Tobolewski 1964, 1966, 1979, 1980, 1981; Fałtynowicz 1992; Izydorek 2010 – localities nos 54–59). Specimens are deposited in the herbaria UGDA, GPN, OLS-L, OLTC and WRSL. Lichen substances were analyzed by standard technique of thin-layer chromatography in solvent systems A, B, C and G (Orange et al. 2001). The nomenclature follows mostly Fałtynowicz and Kukwa (2006), excluding lichens from Teloschistaceae family (Arup et al. 2013) and genera: *Alyxoria* (Ertz & Tehler 2011), *Caeruleum* (Arcadia & Knudsen 2012), *Circinaria* (Nordin et al. 2010), *Graphis* (Neuwirth & Aptroot 2011), *Varicellaria* (Schmitt et al. 2012), *Violella* (Spribile et al. 2011) and *Zwackhia* (Ertz & Tehler 2011), and species: *Brianaria sylvicola* (Ekman & Svensson 2014), *Cladonia floerkeana* (Santesson et al. 2004), *Lecanora filamentosa* (Palice et al. 2011), *L. panonica* (Brodo et al. 1994), *L. saxicola* (Laundon 2010), *Lepraria finkii* (Lendemer 2011), *Melanelia glabratula* (Arup & Sandler Berlin 2011), *Punctelia jeckeri* (Kalb 2007), *Rinodina oleae* (Santesson et al. 2004) and *Usnea dasopoga* (Arcadia 2013).

Categories of threat follow Cieśliński et al. (2006) and are included in text and Table 2: CR – critically endangered, EN – endangered, VU – vulnerable. Following abbreviations are used: for. – forest district; fs – forest section; Ah – *Aesculus hippocastanum*, Al – *Alnus glutinosa*; Apl – *Acer platanoides*; Aps – *Acer pseudoplatanus*; Bet – *Betula* spp.; Cav – *Corylus avellana*; Cb – *Carpinus betulus*; dw – decaying wood; Fr – *Fraxinus excelsior*; Fs – *Fagus sylvatica*; La – *Larix* spp.; Pa – *Picea abies*; Pav – *Padus avium*; Pm – *Pseudotsuga menziesii*; Ps – *Pinus sylvestris*; Pt – *Populus tremula*; Ra – *Ribes alpinum*, Rr – *Rosa rugosa*, Q – *Quercus* spp.; Sal – *Salix* spp.; Sor – *Sorbus aucuparia*; Tc – *Tilia cordata*; gl – soil.

List of localities:

- 1 – 54°13.690'N, 16°20.130'E, Kacza Island;
- 2 – 54°13.274'N, 16°25.874'E, for. Sieciemin, fs 421b, pine forest;
- 3 – 54°12.817'N, 16°26.971'E, for. Sieciemin, fs 421c, pine forest;
- 4 – 54°14.490'N, 16°29.401'E, for. Sieciemin, fs 221f, beech forest;

- 5 – 54°14.333'N, 16°29.587'E, for. Sieciemin, fs 246, beech forest;
- 6 – 54°10.002'N, 16°29.444'E, for. Sieciemin, fs 411c, mixed forest;
- 7 – 54°15.262'N, 16°34.701'E, for. Lejkowo, trees along road, buildings;
- 8 – 54°15.564'N, 16°34.347'E, for. Lejkowo, fs 115a and 111a, water-headwith marshy meadow, beech forest;
- 9 – 54°12.345'N, 16°34.070'E, for. Lejkowo, fs 152, mixed forest;
- 10 – 54°11.333'N, 16°27.156'E, for. Szczeglino, fs 563f, beech forest and stones on cemetary;
- 11 – 54°11.561'N, 16°30.862'E, trees along road between villages Ratajki and Sowno;
- 12 – 54°12.585'N, 16°27.774 'E, Ratajki village, church and trees;
- 13 – 54°14.626'N, 16°22.221'E, for. Sianów, fs 192a;
- 14 – 54°13.354'N, 16°20.864'E, for. Sianów, fs 240c;
- 15 – 54°17.772'N, 16°16.589'E, Iwięcino, church;
- 16 – 54°18.252'N, 16°13.488'E, for. Iwięcino, "Lazy" nature reserve, fs 673f, peatbog with pine forest;
- 17 – 54°18.333'N, 16°13.688'E, for. Iwięcino, fs 673b, beech forest;
- 18 – 54°13.713'N, 16°09.068'E, for. Chełmoniewo, fs 383b, mixed forest;
- 19 – 54°15.710'N, 16°25.919'E, for. Niemica, fs 99c, peatbog and pine forest;
- 20 – 54°11.541'N, 16°16.377'E, for. Koszalin, fs 332f, 343b,c,d, 333h,d; beech forest;
- 21 – 54°16.809'N, 16°22.043'E, for. Kamionka, fs 19b;
- 22 – 54°16.215'N, 16°19.693 ,E, Arboretum;
- 54°14.474'N, 16°20.837'E, Trawica, builgings of the Karnieszewice upper-forestry;
- 24 – 54°12.070'N, 16°12.447'E; 2014-06-05; Koszalin, beech forest;
- 25 – 54°12.272'N, 16°13.313'E; beech forest and walls;
- 26 – 54°11.420'N, 16°17.465'E; for. Chełmoniewo, fs 401A/C, mixed forest;
- 27 – 54°11.408'N, 16°17.333'E; for. Chełmoniewo, fs 392g, mixed forest;
- 28 – 54°11.406'N, 16°17.540'E; for. Chełmoniewo, mixed forest;
- 29 – 54°11.415'N, 16°17.506'E; for. Chełmoniewo, mixed forest;
- 30 – 54°12.287'N, 16°16.226'E; beech forest;
- 31 – 54°12.270'N, 16°15.488'E; peatbog, trees and stones;
- 32 – 54°12.542'N, 16°16.251'E; trees along road;
- 33 – 54°12.159'N, 16°30.160'E; mixed forest;
- 34 – 54°12.145'N, 16°30.039'E; Dolina Białki, mixed forest and beech forest;
- 35 – 54°12.180'N, 16°29.508'N; Dolina Białki, mixed forest, stones in the stream;
- 36 – 54°12.344'N, 16°29.473'E; oak forest;
- 37 – 54°13.066'N, 16°29.467'E; mixed forest and erratic;
- 38 – 54°13.030'N, 16°29.222'E;mixed forest;
- 39 – 54°13.161'N, 16°29.462'E; pine forest;
- 40 – 54°13.240'N, 16°30.490'E; trees along road;
- 41 – 54°14.313'N, 16°32.553'E; trees along road and stones;
- 42 – 54°12.038'N, 16°27.447'E; Kościernica village, trees along road and stones;
- 43 – 54°11.094'N, 16°20.428'E; peatbog;
- 44 – 54°11.059'N, 16°20.361'E; peatbog;
- 45 – 54°10.266'N, 16°23.385'E; Maszkowo village, trees along road;
- 46 – 54°08.372'N, 16°29.434'E; pine forest;
- 47 – 54°08.578'N, 16°30.019'E; pine forest and wall;
- 48 – 54°11.132'N, 16°31.056'E; trees along road;
- 49 – 54°11.515'N, 16°34.134'E; trees along road;
- 50 – 54°11.024'N, 16°33.394'E; Sierakowo Słowieńskie village, church, walls, stones;
- 51 – 54°13.302'N, 16°31.246'E; trees, wall, stones;
- 52 – 54°12.112'N, 16°30.157'E; buildings, trees,
- 53 – 54°18.585'N, 16°12.315'E; Spit of Kopań Lake;
- 54 – 54°15.331'N, 16°02.317'E; Mielno town;
- 55 – 54°15.175'N, 16°00.452'E; between Mielno and Chłopy;
- 56 – 54°18.579'N, 16°12.351'E; Lazy village, between Jamno Lake and Bukowo Lake;
- 57 – 54°16.384'N, 16°06.559'E; Unieście village, on concrete and *Quercus*;
- 58 – 54°17.044'N, 16°16.153'E; Iwięcino village, ca. 1 km on W, *Acer pseudoplatanus* trees along road;
- 59 – 54°14.562'N, 15°57.121'E; Sarbinowo village, lg. Flössner 1935 (Erichsen 1940).
- 60 – 54°16.243'N, 16°32.258'E; between Bartolino and Sulechowo villages, *Acer platanoides* trees along road;
- 61 – 54°16.157'N, 16°33.031'E; Sulechówko village, near church; stones and different species of trees;
- 62 – 54°15.304'N, 16°34.269'E; between Kukułczyn and Lejkowo villages, wood and different species of trees;
- 63 – 54°16.446'N, 16°18.407'E; Wierciszewo village, limes along road and wall;
- 64 – 54°22.284'N, 16°18.236'E; Spit of Bukowo Lake, Dąbki village, ca. 1 km on W; different species of trees;
- 65 – 54°21.495'N, 16°17.083'E; Spit of Bukowo Lake, Dąbki village, ca. 2 km on W; different species of trees;
- 66 – 54°16.363'N, 16°13.222'E; between Kleszcze and Osieki villages; maples along road and concrete;
- 67 – 54°17.010'N, 16°12.390'E; Osieki village, church, stones, wall, ashes and beeches;
- 68 – 54°18.019'N, 16°10.428'E; Spit of Jamno Lake, Łazy village, ca. 1,5 km on W.

4. Results and discussion

Lichenological data from the Karnieszewice Forest Division and adjacent areas are relatively insufficient. The oldest, single information is comprised in publications of Erichsen (1936, 1940), Tobolewski (1964, 1966, 1979, 1980, 1981) and Fałtynowicz (1992). Several species from south-east part of Karnieszewice Forest Division have been reported by Izydorek (2010), and a brief information about Koszalin City lichens can be found in the conference abstract by Bezmian (1998). The most complete data have been published recently by Kukwa et al. (2013), however those records represented only preliminary results.

Altogether 270 lichen species were found in the explored area. Such a large number of species is apparently a result related to the great diversity of substrates and habitats in this area. This number of species is relatively high also for the Polish Lowland (Table 1). There is no doubts it is also a result of the relatively good condition of forest associations, which despite intensive forest cultivation (from 19th century), preserved their values and specific phytoclimate.

In comparision to other protected areas in northern Poland with similar acreage and partly also habitats, the Karnieszewice Forest Division is one of the richest; also numerous red-listed species (Table 2) and taxa protected by law (Table 3) occur there. Epiphytic lichens dominate in the lichen biota (165 species), but the group of wood-inhabiting species is also rich (75 species). Epilithic lichens are also numerous (69 species), however, on the natural substrate (boulders and stones) only 47 species have been found. On antropogenic substrate (concrete, bricks, tiles, eternite) 38 further species have been recorded. The biota of terricolous lichens includes 45 species (most of them belong to *Cladonia* genus); this group of lichens does not have suitable habitats in the study area. In addition seven species were also found on metal substrate.

The importance of the Karnieszewice Forest Division for the lichen diversity protection can be also proved by the very long list of lichen taxa with the highest categories of threat in Poland. 53 such species have been found (19,6% of the whole local biota) (Table 2), five of which with CR category and 21 species with EN category. Among them there are taxa particularly rare in the country, such as *Cladonia parasitica* and *Ramalina baltica*, but some of them occur very frequently in northern Poland thus in studied area their category of threat is surely lower. It is applicable among the others for *Cladonia caespitica*, *Flavoparmelia caperata*, *Pleurosticta acetabulum*, *Ramalina fastigiata*, *R. fraxinea* and *Usnea florida*. *Agonimia flabelliformis* is reported for the second time from Polish lowland (Kowalewska unpubl. data).

In the Karnieszewice Forest Division 34 species protected by law were found (Table 3).

Table 1. The numer of lichen species known from selected protected areas in N Poland with reference to the Karnieszewice Forest Division

Name of protected area	Number of species
Białowieski National Park (Cieśliński 2003)	ca. 360
Wigierski National Park (Fałtynowicz 2014)	303
Wdzydzki Landscape Park (Kukwa et al. 2012)	276
Karnieszewice Forest Division	270
Słowiński National Park (Fałtynowicz 2015, unpubl.)	270
Woliński National Park (Wieczorek 2014)	266
Drawieński National Park (Schiefelbein et al. 2012)	262
Suwalski Landscape Park (Zalewska et al. 2004a)	231
Landscape Park of Puszcza Romincka (Zalewska et al. 2004b)	222
Szczeciński Landscape Park (Wieczorek 2005)	207

List of species

In the list numbers of sites and substrate type were noted for each species; + – non-lichenized fungus

- Absconditella lignicola* Věžda & Pišút – 3, 34, 35, 38, 39; Al, Ps, dw;
A. sphagnorum Věžda & Poelt – 43, 44; on *Sphagnum* spp.
Acarospora fuscata (Nyl.) Th. Fr. – 12, 15, 31, 50, 51; stones, wall;
A. moenium (Vain.) Räs. – 47; wall;
Acrocordia gemmata (Ach.) A. Massal. – 3; Pt;
Agonimia flabelliformis Halda, Czarnota & Guzow-Krzemińska – 35; Al;
Alyxoria varia (Pers.) Ertz & Tehler – 23; Apl; (Izydorek 2010);
Amandinea punctata (Hoffm.) Coppins & Scheid. – 9, 11, 40–42, 45, 51, 52, 56, 58, 60, 61, 66; Apl, Aps, Bet, Fr, Pa, Q; (Izydorek 2010);
Anaptychia ciliaris (L.) Körb. – 61; Apl;
Anisomeridium polypori (M.B. Ellis & Everh.) M.E. Barr – 3, 32, 48; Apl, Fr, Pt;
Arthonia didyma Körb. – 33; Cb;
A. mediella Nyl. – 20, 29; Q; (Izydorek 2010);
A. radiata (Pers.) Ach. – 9, 32, 33, 35, 37, 39, 54, 64; Apl, Bet, Cav, Cb, Fr, Fs, Ps, Q, Sal, Sor;
A. spadicea Leigh. – 3, 8, 10, 26, 27, 30–33, 36–38, 54, 64, 65; Al, Cav, Cb, Fr, Pav, Ps, Q, Tc;

Table 2. Species from Polish Red List of Lichens (Cieśliński et al. 2006) found in the Karnieszewice Forest Division

Categories of threats	Name of species	Number of species/ Total number of localities	% of lichen biota
CR	<i>Bryoria cf. capillaris</i> , <i>Chrysotrichia candelaris</i> , <i>Melanohalea olivacea</i> , <i>Pertusaria hymenea</i> , <i>Pyrrhospora quernea</i>	5/9	1.8
EN	<i>Anaptychia ciliaris</i> , <i>Arthonia didyma</i> , <i>Calicium adspersum</i> , <i>Caloplaca chlorina</i> , <i>Cetraria sepincola</i> , <i>Chaenotheca brunneola</i> , <i>Ch. stemonea</i> , <i>Cladonia caespiticia</i> , <i>C. parasitica</i> , <i>Fellhanera bouteillei</i> , <i>Flavoparmelia caperata</i> , <i>Lecanora intumescens</i> , <i>Opegrapha atra</i> , <i>Pertusaria flavidula</i> , <i>Physconia distorta</i> , <i>Ph. perisidiosa</i> , <i>Pleurosticta acetabulum</i> , <i>Ramalina baltica</i> , <i>R. fastigiata</i> , <i>R. fraxinea</i> , <i>Usnea florida</i>	21/90	7.8
VU	<i>Acrocordia gemmata</i> , <i>Arthonia mediella</i> , <i>Bacidia rubella</i> , <i>Biatora efflorescens</i> , <i>Bryoria fuscescens</i> , <i>Calicium glauceum</i> , <i>C. salicinum</i> , <i>C. viride</i> , <i>Cetraria chlorophylla</i> , <i>C. islandica</i> , <i>Cliostomum griffithii</i> , <i>Melanohalea elegantula</i> , <i>Ochrolechia androgyna</i> , <i>Opegrapha rufescens</i> , <i>O. vulgata</i> , <i>Parmelia submontana</i> , <i>Parmelina tiliacea</i> , <i>Peltigera canina</i> , <i>P. praetextata</i> , <i>Pertusaria pertusa</i> , <i>Punctelia subrudecta</i> , <i>Pyrenula nitida</i> , <i>Ramalina farinacea</i> , <i>Usnea dasopoga</i> , <i>U. hirta</i> , <i>Varicellaria hemisphaerica</i> , <i>Zwackhia viridis</i>	27/136	10
Total		53/235	19.6

Table 3. Lichen species protected by Polish law (Rozporządzenie Ministra Środowiska 2014) in the Karnieszewice Forest Division

Protection forms	Name of species	Number of species/ Total number of localities	% of lichen biota
Total protection	<i>Anaptychia ciliaris</i> , <i>Bryoria cf. capillaris</i> , <i>Cetraria sepincola</i> , <i>Chrysotrichia candelaris</i> , <i>Melanohalea elegantula</i> , <i>M. olivacea</i> , <i>Parmelia submontana</i> , <i>Parmelina tiliacea</i> , <i>Peltigera hymenina</i> , <i>Punctelia jeckeri</i> , <i>P. subrudecta</i> , <i>Ramalina baltica</i> , <i>R. fastigiata</i> , <i>R. fraxinea</i> , <i>Usnea florida</i> , <i>Xanthoparmelia delisei</i>	15/70	5.7
Partly protected	<i>Bryoria fuscescens</i> , <i>Cetraria islandica</i> , <i>Cladonia arbuscula</i> , <i>C. ciliata</i> , <i>C. portentosa</i> , <i>C. rangiferina</i> , <i>Flavoparmelia caperata</i> , <i>Hypogymnia tubulosa</i> , <i>Imshaugia aleurites</i> , <i>Melanelia subaurifera</i> , <i>Peltigera canina</i> , <i>P. membranacea</i> , <i>P. polydactylon</i> , <i>Pleurosticta acetabulum</i> , <i>Ramalina farinacea</i> , <i>Usnea dasopoga</i> , <i>U. hirta</i> , <i>Vulpicida pinastri</i>	18/133	7.0
Total		33/203	12.7

- A. vinosa* Leight. – 29, 36–38; Q;
Arlothelium ruanum (A. Massal.) Körb. – 8, 35, 54; Cav, Pav;
Aspicilia cinerea (L.) Körb. – 15, 50; stones;
Athallia holocarpa (Hoffm.) Arup, Frödén & Söchting – 12, 15, 51, 52, 55, 67, 68; walls, stone;
A. pyracea (Ach.) Arup, Frödén & Söchting – 1; Pt;
Bacidia rubella (Hoffm.) A. Massal. – (Izydorek 2010);
Bacidina sulphurella (Samp.) M. Hauck & V. Wirth – 24, 26–28, 32–35; Cav, Cb, Fs, Q;
Baeomyces rufus (Huds.) Rebent. – 31, 50; gl, stones; (Izydorek 2010);
Biatora efflorescens (Hedl.) Erichsen – 5, 9, 18, 24, 26, 30, 37; Fr, Fs, Q;
B. globulosa (Flörke) Fr. – 22, 36, 38, 40, 55; Apl, Aps, Fr;
Bilimbia sabuletorum (Schreb.) Arnold – 54; on bryophytes over old wall;
Brianaria sylvicola (Flot. ex Körb.) S. Ekman & M. Svensson – (Izydorek 2010);
Bryoria cf. capillaris (Ach.) Brodo & D. Hawksw. – 36; Q;
B. fuscescens (Gyelnik) Brodo & D. Hawksw. – 6, 36, 37, 39, 40, 52, 54, 55; Bet, La, Q, dw;
Buellia alboatra s.l. – 15, 50, 51, 67; stone, walls;
B. griseovirens (Turner & Borrer ex Sm.) Almb. – 3, 4, 9, 16, 18, 24, 26, 27, 29, 30, 33–35, 38, 42, 48, 51, 52, 54, 67; Al, Bet, Cav, Cb, Fr, Fs, Pt, Q, Sor, dw;
Calicium adspersum Pers. – 28, 37, 38; Q;
C. glauceum Ach. – 39; dw;

- C. parvum* Tibell – 39, 43, 47; Ps;
C. pinastri Tibell – 43, 53; Ps;
C. salicinum Pers. – 20, 29, 35, 36, 38, 40, 53; Q, dw; (Izydorek 2010);
C. viride Pers. – 18, 29, 31, 35, 36, 38; Fr, Q;
Calogaya decipiens (Arnold) Arup, Frödén & Söchting – 3, 15, 51, 55, 63; walls; (Izydorek 2010);
C. pusilla (A. Massal.) Arup, Frödén & Söchting – 15, 63, 67, 69; walls;
Caloplaca cerina (Ehrh. ex Hedwig) Th. Fr. – 36; dw;
C. chlorina (Flot.) H. Olivier – 15; Fr;
C. obscurella (J. Lahm ex Koerb.) Th. Fr. – 11, 41, 45; Apl, Aps;
C. saxicola (Hoffm.) Nordin – 51, 55, 67; walls;
Candelaria concolor (Dickson) B. Stein – 9, 45, 52; Apl, Aps, Fr, dw;
C. pacifica Westberg – 48, 61, 63; Fr, Tc;
Candelariella aurella (Hoffm.) A. Zahlbr. – 3, 11, 15, 25, 51, 52, 55, 61, 66, 69; Aps, walls; (Izydorek 2010);
C. efflorescens R.C. Harris & W.R. Buck – 9, 24, 26, 36, 61; Apl, Cav, Fr, Pt;
C. vitellina (Hoffm.) Müll. Arg. – 12, 15, 48, 51; Fr, stones; (Izydorek 2010);
C. xanthostigma (Ach.) Lettau – 11; 15, 40–42, 48, 56; Aps, Apl, Fr, Q; (Izydorek 2010);
Catillaria nigroclavata (Nyl.) Schuler – 24, 26, 36; Pt, Fs, dw;
Cetraria aculeata Ach. – 14; gl; (Izydorek 2010);
C. chlorophylla (Willd.) Vain. – 6, 7, 11, 22, 40, 42, 52, 54, 55, 62; Aps, Bet, La, Pa, Q, metal, dw;
C. islandica (L.) Ach. – 2; gl; (Izydorek 2010);
C. sepincola (Ehrh.) Ach. – 31, 52; Bet; (Izydorek 2010);
Chaenotheca brunneola (Ach.) Müll. Arg. – 29; Q;
Ch. chrysocephala (Ach.) Th. Fr. – 3, 17, 18, 20, 22, 28, 31, 33, 35–38; Al, Bet, Fr, Q, dw;
Ch. ferruginea (Turner ex Sm.) Mig. – 1, 10, 16, 18–20, 22, 26, 28–31, 33–40, 43, 44, 46, 47, 52–55, 61, 64, 67; Al, Aps, Bet, Fr, La, Ps, Q, dw; (Izydorek 2010);
Ch. stemonea (Ach.) Müll. Arg. – 26, 35; dw, Al;
Ch. trichialis (Ach.) Th. Fr. – 36; Bet; (Izydorek 2010);
Chrysotrichia candelaris (L.) J.R. Laundon – 17, 28, 29, 37, 38, 61; Fr, Q; (Izydorek 2010);
Circinaria caesiocinerea (Nyl. ex Malbr.) A. Nordin, S. Savić & Tibell – 51; stones;
C. calcarea (L.) A. Nordin, S. Savić & Tibell – 52; wall;
C. contorta (Hoffm.) A. Nordin, S. Savić & Tibell – 3, 25, 51, 52; walls;
Cladonia arbuscula s.l. – 2, 14, 19, 39, 44, 55, 67; gl; (Izydorek 2010);
C. caespiticia (Pers.) Flk. – 29; gl; (Izydorek 2010);
C. cenotea (Ach.) Schaer. – 2, 3, 31, 35, 37, 43; Bet, Q, dw; (Izydorek 2010);
C. cervicornis (Ach.) Flot. ssp. *verticillata* (Hoffm.) Ahti – 14, 37, 67; gl;
C. chlorophaea (Flörke ex Sommerf.) Spreng. s.str. – 14, 54, 55; Bet, gl; (Izydorek 2010);
C. ciliata Stirt. – 2, 14, 39, 44, 55; gl; between *Sphagnum* on peatbog;
C. coniocraea (Flörke) Vain. – 1–3, 8–10, 13, 16, 18–22, 24, 25, 27–31, 33–38, 40, 42–44, 46, 47, 52–55, 62, 64, 65, 67; Al, Bet, Fr, Fs, Pa, Ps, Q, gl, dw; (Izydorek 2010);
C. cornuta (L.) Hoffm. – 3, 14, 55; Bet, gl; (Izydorek 2010);
C. deformis (L.) Hoffm. – 31, 46, 47; gl, dw; (Izydorek 2010);
C. digitata (L.) Hoffm. – 1–3, 13, 16, 19, 21, 22, 26, 27, 29, 31, 34–37, 39, 42, 46, 47, 52, 54, 55, 64; Al, Bet, Fs, La, Ps, Q, dw, gl; (Izydorek 2010);
C. fimbriata (L.) Fr. – 3, 16, 19, 27, 29, 31, 32, 40, 45, 52, 54, 55, 62, 67; Apl, Aps, Bet, Ps, dw, gl; (Izydorek 2010);
C. floerkeana (Fr.) Flörke – 3, 14, 27, 46; Bet, gl, dw; (Izydorek 2010);
C. foliacea (Huds.) Willd. – 55; gl;
C. furcata (Huds.) Schrad. – 2, 14, 31, 51–53, 55, 67; gl, dw; (Izydorek 2010);
C. glauca Flörke – 2, 14, 19, 55; gl; (Izydorek 2010);
C. gracilis (L.) Willd. – 14, 39, 55; gl; (Izydorek 2010);
C. grayi G. Merr. ex Sandst. – 2, 3, 26, 31; gl, dw;
C. macilenta Hoffm. – 2, 3, 14, 19, 26, 35, 36, 40, 46, 55; Bet, gl, dw; (Izydorek 2010);
C. novochlorophaea (Sipman) Brodo & Ahti – 19; gl;
C. parasitica (Hoffm.) Hoffm. – 34, dw;
C. phyllophora Hoffm. – 14, 55; gl; (Izydorek 2010);
C. pleurota (Flörke) Schaer. – 9; Fr;
C. portentosa (Dufour) Coem. – 19, 55, 67; gl; (Izydorek 2010);
C. pyxidata (L.) Hoffm. – 46; gl; (Izydorek 2010);
C. ramulosa (With.) J.R. Laundon – 32; Fr; (Izydorek 2010);
C. rangiferina (L.) Web. – 55; gl; (Izydorek 2010);
C. scabriuscula (Delise) Leight. – 53, 55, 67; gl;
C. squamosa Hoffm. – 26; humus, dw;
C. subulata (L.) Weber ex F.H. Wigg. – 23, 55; gl; (Izydorek 2010);
C. uncialis (L.) Weber ex F.H. Wigg. – 2, 39; gl; (Izydorek 2010);
Cliostomum griffithii (Sm.) Coppins – 18, 53, 64, 65, 67; Apl, Fr, Q;
Coenogonium pineti (Schrad. ex Ach.) Lücking & Lumbsch – 3, 18, 21, 26, 31, 33, 35, 36, 42, 64; Al, Bet, Fs, Pa, Ps, dw;
Evernia prunastri (L.) Ach. – 2, 3, 7, 9–11, 13, 15, 18–20, 22, 31, 32, 34–40, 42, 49, 51–55, 58, 60, 61, 62, 64–67; Al, Apl, Aps, Bet, Cav, Cr, Fr, Fs, La, Pa, Pm, Po, Ps, Q, Sor, Tc, U, dw, stones, metal; (Izydorek 2010);

- Fellhanera bouteillei* (Desm.) Vězda – 37–39, 46; twigs of La and Pa;
- F. subtilis* (Vězda) Diederich & Sérus. – 3, 6, 29, 37, 38, 45; twigs of *Vaccinium myrtillus* and Pa;
- Fellhaneropsis myrtillicola* (Erichsen) Hafellner – 29, 37, 38, 45; twigs of Pa;
- Flavoparmelia caperata* (L.) Hale – 15, 18, 67; Fr, Fs;
- Flavoplaca citrina* (Hoffm.) Arup, Frödén & Søchting – 15, 52, 57, 61, 67, 69; walls, eternit; (Izydorek 2010);
- F. flavocitrina* (Nyl.) Arup, Frödén & Søchting – 15, 51, 52; stone, walls;
- Fuscidea arboricola* Coppins & Tønsberg – 3, 8, 35; Bet, Al;
- F. pusilla* Tønsberg – 2, 3, 31, 33, 35, 36, 38, 39, 43, 44, 46, 53, 64; Bet, Cav, Cb, Pa, Ps;
- Graphis betulina* (Pers.) Ach. – 4, 8 (cf); Fs;
- G. pulverulenta* (Pers.) Ach. – 4, 5, 8–10, 17, 21, 26–28, 30–35, 38, 40, 44; Cb, Al, Cav, Fs, Q, Aps, Sor;
- G. scripta* (L.) Ach. – 28, 30, 32–35, 54; Apl, Cav, Fr, Fs, Pav;
- Halecania viridescens* Coppins & P. James – 33; Pav;
- Hypocenomyce caradocensis* (Leight. ex Nyl.) P. James & Gotth. Schneid. – 22, 53; Pm, Q;
- H. scalaris* (Ach.) M. Choisy – 2, 3, 7, 13, 16, 19, 20, 22, 25, 26, 29–31, 35–37, 39, 40, 42–44, 46, 47, 52–55, 62, 64, 66; Al, Apl, Bet, Cav, La, Pa, Ps, Q, Tc; dw, stone; (Izydorek 2010);
- Hypogymnia physodes* (L.) Nyl. – 3, 9, 10, 13, 14, 16, 18–22, 24, 25, 27, 29, 31, 32, 34–40, 42, 43, 44, 46, 47, 51–55, 58, 61, 62, 64, 67, 68; Al, Aps, Bet, Cav, Fr, Fs, La, Pt, Ps, Pa, Q, Tc; dw, stones, sand; (Izydorek 2010);
- H. tubulosa* (Schaer.) Havaas – 3, 6, 9, 13, 14, 19, 20, 22, 25, 36, 37–40, 42, 46, 52, 53, 61, 62, 68; Bet, Cav, Fr, Fs, La, Pa, Po, Ps, Q, Sal, dw, stone; (Izydorek 2010);
- Imshaugia aleurites* (Ach.) S.L.F. Meyer – 2, 3, 19, 31, 36, 39, 40, 43, 44, 47, 53, 55, 62; Bet, Ps, Pm, dw;
- Lecania cyrtella* (Ach.) Th. – 36, 40; Aps, dw;
- L. hyalina* (Fr.) R. Sant. – (Izydorek 2010);
- L. naegelii* (Hepp) Diederich & P. Boom – 9, 32, 36, 41, 49, 54, 56; Apl, Aps, Fr, Fs, Pt, U; (Izydorek 2010);
- Lecanora albella* (Pers.) Ach. – (Izydorek 2010);
- L. albescens* (Hoffm.) Flörke – 3, 12, 15, 25, 51, 52, 55, 67; walls; (Izydorek 2010);
- L. allophana* Nyl. – 41, 46, 48, 53; Apl, Fr, Cr, Ps;
- L. argentata* (Ach.) Malme – 8, 9, 18, 29, 33, 35, 40, 45, 48, 54–56, 58; Al, Apl, Aps, Cb, Fr, Fs, Q, dw; (Izydorek 2010);
- L. carpinea* (L.) Vain. – 1, 9, 33, 35, 36, 41, 49, 53–56, 64; Apl, Cb, Cr, Fr, Pt, Q, U, dw; (Izydorek 2010);
- L. chlarotera* Nyl. – 2, 9, 11, 15, 22, 32, 33, 36, 42, 45, 48, 49, 52–56, 64; Apl, Aps, Bet, Cr, Fr, Pt, Q, U, dw; (Izydorek 2010);
- L. compallens* van Herk & Aptroot – 11, 35, 36; Aps, Cb;
- L. conizaeoides* Nyl. ex Cromb. – 2, 3, 16, 19, 20, 22, 25, 27, 29, 31, 35, 40, 43, 46, 47, 52–55, 58, 62, 64; Al, Aps, Bet, Fr, La, Pa, Ps, Q, Sor, dw; (Izydorek 2010);
- L. dispersa* (Pers.) Sommerf. – 15, 51, 52, 55, 57, 61, 63, 66–69; walls; (Izydorek 2010);
- L. expallens* Ach. – 9, 11, 15, 16, 18, 20, 22, 29–32, 37, 38, 45, 51–55; Apl, Aps, Al, Bet, Fr, Fs, Pa, Q; (Izydorek 2010);
- L. filamentosa* (Stirt.) Elix & Palice – 2, 3, 13, 14, 18, 19, 24, 25, 27, 31, 38, 40, 43, 44, 47; Bet, Fr, Ps, dw;
- L. glabrata* (Ach.) Malme – 22; Fr;
- L. hagenii* (Ach.) Ach. – 11, 15, 51; Aps, walls;
- L. intumescens* (Rebent.) Rabenh. – 5, 35, 54; Cb, Fs;
- L. pannonica* Szat. – 15; wall;
- L. persimilis* (Th. Fr.) Nyl. – 45, 52; Apl, Pa;
- L. polytropa* (Ehrh.) Rabenh. – 12, 41, 50, 51, 61; stones;
- L. populicola* (DC.) Duby – 48, 53; Cr, Fr;
- L. pulicaris* (Pers.) Ach. subsp. *pulicaris* – 3, 11, 18, 20, 22, 24, 31, 35–40, 42, 43, 47, 53, 54, 65, 68; Al, Aps, Bet, Cav, Cr, Fr, Fs, La, Pa, Ps, Pt, Pav, Q, dw; (Izydorek 2010);
- L. rupicola* (L.) A. Zahlbr. – 15; stone;
- L. saligna* (Schrad.) A. Zahlbr. – 52; dw;
- L. sarcopidoides* (A. Massal.) A.L. Smith – 62; dw;
- L. saxicola* (Pollich) Ach. – 15, 51, 52, 61, 69; walls, stone; (Izydorek 2010);
- L. semipallida* H. Magn. – 3, 12, 51; wall;
- L. symmicta* (Ach.) Ach. – 14, 19, 22, 36, 39, 52, 53; Cr, Ps, Q, dw;
- L. thysanophora* R.C. Harris – 33; Fs;
- L. varia* (Hoffm.) Ach. – 17, 29, 39, 40, 52; Pa, dw;
- Lecidea fuscoatra* (L.) Ach. – 50, 51; stones;
- L. nylanderi* (Anzi) Th. Fr. – 2, 3, 13, 19, 25, 26, 36–39, 43, 44, 46, 47, 49, 53, 68; Bet, Pa, Ps, Q, U, dw;
- Lecidella elaeochroma* (Ach.) M. Choisy – 1, 9, 15, 33, 35, 36, 40, 41, 48, 49, 52–56, 61, 64, 65, 68, 69; Ah, Apl, Cav, Cb, Fr, Fs, Pav, Po, Pt, Rr, Q, U; (Izydorek 2010);
- L. flavosorediata* (Vezda) Hertel & Leuckert – 11, 28, 29, 48; Aps, Fr, Q;
- L. stigmatea* (Ach.) Hertel & Leuckert – 12, 15, 51, 52, 67–69; walls, stone; (Izydorek 2010);
- Lepraria eburnea* J.R. Laundon – 35; Bet;
- L. elobata* Tønsberg – 2, 3, 15, 20, 22, 24, 26, 28, 31, 33, 35, 64; Bet, Cb, Fs, Ps, Q, stone;
- L. finkii* (B. de Lesd.) R.C. Harris – 3, 4, 8–10, 17, 18, 21, 24, 26, 27, 28, 29, 30, 33, 34, 35, 38, 61, 65; Al, Apl, Cb, Fr, Fs, Pt, Q, Sor;
- L. incana* (L.) Ach. – 2–4, 7, 8, 16–18, 20, 22, 24–33, 35–39, 42, 46–48, 50, 53, 61, 64, 68; Al, Bet, Cb, Fr, Fs, La, Pa, Ps, Q, Sor, Tc, dw, stone;
- L. jackii* Tønsberg – 2, 3, 13, 26, 29, 39; Bet, Pa, Ps, Q;
- L. neglecta* (Nyl.) Erichsen – 50; stone;
- L. rigidula* (B. de Lesd.) Tønsberg – 3, 13, 20, 24, 26, 32; Aps, Bet, Fs, Pt, Q;

- L. vousauxii* (Hue) R.C. Harris – 8, 15, 35, 45, 51; Al, Apl, Fr, stone;
+*Leptorhaphis epidermidis* (Ach.) Th. Fr. – 40; Bet;
Lichenomphalia umbellifera (L.) Redhead, Lutzoni, Moncalvo & Vilgalys – 17, 24, 46; gl, dw;
Melanelia fuliginosa (Fr. ex Duby) O. Blanco et al. s.str. – 11, 12, 61; stones;
M. glabratula (Lamy) Sandler & Arup – 1, 3, 4, 7–9, 17–28, 30–33, 35, 37–40, 42, 45, 48, 51–55, 62, 64–67; Al, Apl, Aps, Bet, Cav, Cb, Cr, Fr, Fs, Pa, Po, Ps, Pt, Q, Sal, Sor, Tc, dw, stones; (Izydorek 2010);
M. subaurifera (Nyl.) O. Blanco & al. – 9, 11, 18, 22, 33, 40, 53, 61, 65, 68; Aps, Bet, Cav, Fr, Fs, Ps, Tc, dw;
Melanohalea elegantula (A. Zahlbr.) O. Blanco & al. – 9, 11, 22, 62; Aps, Fr, Pt, Tc; (Izydorek 2010);
M. exasperatula (Nyl.) O. Blanco & al. – 6, 7, 9, 11, 15, 40, 42, 48, 49, 51, 52, 61; Apl, Aps, Bet, Fr, La, Q, Tc, U, dw, metal, stone, eternit, wall;
M. olivacea (L.) Blanco & al. – 55 (Tobolewski 1981).
Micarea byssacea (Th. Fr.) Czarnota, Guzow-Krzemińska & Coppins – 35; Cb, Q;
M. denigrata (Fr.) Hedl. – 38, 40, 43, 47, 52, 55; Ps, dw;
M. lithinella (Nyl.) Hedl. – 2, 35; pebbles;
M. micrococca (Körb.) Gams ex Coppins – 25, 26, 31, 37, 38, 39, 44, 46; Al, Bet, Ps, dw;
M. misella (Nyl.) Hedl. – 2, 3, 34, 35, 38, 39, 44, 46, 47; dw;
M. nitschkeana (J. Lahm ex Rabenh.) Harm. – 38, 43, 46; Pa, Ps;
M. prasina Fr. – 2, 33, 39, 40; Aps, Ps, dw;
M. viridileprosa Coppins & van den Boom – 39, 46; Ps, dw;
Microcalicium disseminatum (Ach.) Vainio – 28, 29; Q;
Ochrolechia androgyna (Hoffm.) Arnold – (Izydorek 2010) (most probably this record belongs to *O. bahusienensis*);
O. bahusienensis H. Magn. – 37, 39; Q, dw;
O. microstictoides Räsänen – 3, 6, 31, 36–40, 42, 43; Bet, La, Pt, Q, dw;
O. turneri (Sm.) Hasselrot – 15, 45; Aps, Fr;
Opegrapha atra Pers. – 54; Fr;
O. rufescens Pers. – 32; Apl;
O. vulgata (Ach.) Ach. – 30–32; Apl, Q;
Parmelia saxatilis (L.) Ach. s.l. – 8, 9, 32, 36, 37; Bet, Cb, Fr; (Izydorek 2010);
P. serrana A. Crespo, M.C. Molina & D. Hawksw. – 43; Q;
P. submontana Nádv. ex Hale – 24, Apl;
P. sulcata Taylor – 1, 3, 6, 7, 9, 10, 11 (c.ap.), 13, 15, 18–20, 22, 24, 25, 31–33, 35–42, 45, 46, 48, 49, 51–58, 60, 62, 64–69; Al, Apl, Aps, Bet, Cav, Fr, Fs, La, Pa, Ps, Q, Sal, Tc, U, metal, dw, stones, wall; (Izydorek 2010);
Parmelina tiliacea (Hoffm.) Hale – 15, 45, 63; Apl, Fr, Tc;
Parmeliopsis ambigua (Wulfen) Nyl. – 2, 3, 19–21, 25, 26, 30, 31, 36, 37, 39, 42, 46, 47, 52, 53, 55, 62; Bet, Fs, Pa, Ps, dw; (Izydorek 2010);
Peltigera canina (L.) Willd. – 55; gl;
P. didactyla (With.) J.R. Laundon – 14; gl,
P. hymenina (Ach.) Delise ex Duby – 64; on sand on dune;
P. membranacea (Ach.) Nyl. – 8; stone;
P. polydactylon (Neck.) Hoffm. – 54, 55; gl;
P. praetextata (Flörke) Zopf – 10, 28, stone, dw;
P. rufescens (Weiss) Humb. – 29, 52, 55; dw, gl;
Pertusaria albescens (Huds.) M.Choisy & Werner – 24, 34, 40, 41, 45, 48, 51, 55, 56, 67; Apl, Aps, Cb, Fr, Fs, Pt; (Izydorek 2010);
P. amara (Ach.) Nyl. – 2, 4, 6–9, 11, 15, 17, 18, 20–22, 25–27, 29–36, 38, 48, 53, 54, 58, 61, 66, 67; Al, Apl, Aps, Cb, Fr, Fs, Q, Sor, Tc; (Izydorek 2010);
P. coccodes (Ach.) Nyl. – 6, 9, 18, 22, 29, 38, 53, 56, 61; Al, Apl, Fr, Fs, Pav, Q; (Izydorek 2010);
P. flavidia (DC.) J.R. Laundon – 6, 29, 37, 38, 45; Apl, Fs, Q; (Izydorek 2010);
P. hymenea (Ach.) Schaer. – 4; Fs;
P. leioplaca DC. in Lam. & DC. – 4, 8, 18, 33–35, 54; Cb, Fr, Fs, Sor; (Izydorek 2010);
P. pertusa (L.) Tuck. – 4, 8, 9, 11, 17, 18, 20, 22, 26, 28, 31, 34, 35, 38, 41, 45; Apl, Aps, Cb, Fr, Fs, Q; (Izydorek 2010);
P. pupillaris (Nyl.) Th. Fr. – 31; Q;
Phaeophyscia nigricans (Flörke) Moberg – 12, 51, 55; walls; (Izydorek 2010);
Ph. orbicularis (Necker) Moberg – 9, 11, 15, 18, 40, 41, 45, 49, 51, 52, 55, 58, 60, 61, 63, 67, 69; Apl, Aps, Fr, Fs, Po, Tc, U, walls, stone; (Izydorek 2010);
Phlyctis argena (Ach.) Flot. – 1, 3, 4, 6–11, 15, 17, 18, 21, 22, 25–28, 32, 34–36, 40, 41, 48, 51–55, 60, 63, 65–67; Al, Apl, Aps, Bet, Cav, Cb, Fr, Fs, Ps, Pt, Q, Sal, Sor, Tc, dw; (Izydorek 2010);
Physcia adscendens (Fr.) H. Olivier – 1, 9, 11, 12, 15, 18, 22, 25, 40, 42, 45, 48, 49, 52, 53, 55, 56, 58, 60, 63–68; Apl, Aps, Bet, Cav, Fr, Fs, La, Po, Ps, Q, Rr, Sal, U, walls; (Izydorek 2010);
Ph. aipolia (Ehrh. ex Humb.) Fürnr. – 9, 32; Apl, Fr;
Ph. caesia (Hoffm.) Fürnr. – 7, 15, 51, 52; metal, stone, wall; (Izydorek 2010);
Ph. dubia (Hoffm.) Lettau – 11, 51, 52, 66, 67; Apl, Aps, walls;
Ph. stellaris (L.) Nyl. – 9, 36, 48, 49, 65; Fr, Q, U;
Ph. tenella (Scop.) DC. – 1, 6, 8, 9, 11, 18, 25, 26, 35, 36, 37, 40, 41, 42, 45, 48, 49, 52, 53, 54, 55, 56, 60, 63–68; Apl, Aps, Bet, Cav, Cr, Fr, Fs, Pa, Po, Ps, Q, Sal, Tc, U, dw, wall; (Izydorek 2010);
Physconia distorta (With.) J.R. Laundon – 1, 11, 40, 45, 56, 61; Apl, Aps, Fr, Po, Pt; (Izydorek 2010);
Ph. enteroxantha (Nyl.) Poelt – 9, 11, 25, 48; Apl, Aps, Fr; (Izydorek 2010);
Ph. grisea (Lam.) Poelt – 11, 48, 60, 61; Apl, Aps, Fr;
Ph. perisidiosa (Erichsen) Moberg – 11, 61; Apl, Aps;

- Placynthiella dasaea* (Stirt.) Tønsberg – 2, 3, 14, 15, 19, 24–26, 30, 31, 36, 40, 46, 62, 68; Bet, Fs, Q, dw, humus;
- P. icmalea* (Ach.) Coppins & P. James – 3, 20, 21, 27, 37, 46, 62; Bet, dw;
- P. oligotropha* (J.R. Laundon) Coppins & P. James – 37, 46; gl, humus; (Izydorek 2010);
- P. uliginosa* (Schrader) Coppins & P. James – 37, 43, 54, 55; Bet, gl, dw; (Izydorek 2010);
- Platismatia glauca* (L.) W.L. Culb. & C.F. Culb. – 2, 3, 6, 10, 13, 20, 21, 24–26, 29, 31, 33, 35, 36, 38–40, 42, 46, 52, 54, 55, 62; Al, Bet, Cav, Fs, La, Pa, Ps, Q, Sor, dw; (Izydorek 2010);
- Pleurosticta acetabulum* (Necker.) Elix & Lumbsch – 11, 15, 40–42, 45, 48, 55, 56, 60, 61; Apl, Aps, Fr, Po, Q, Tc; (Izydorek 2010);
- Polycauliona candelaria* (L.) Frödén, Arup & Söchting – 11, 22, 26, 45, 58, 66; Apl, Aps, Fs, La; (Izydorek 2010);
- P. polycarpa* (Hoffm.) Frödén, Arup & Söchting – 1, 17, 19, 35, 37, 41, 42, 45, 48, 49, 52, 54, 60, 63, 65; Apl, Aps, Bet, Cav, Fr, Pa, Pt, Q, Sal, Tc, U, wall, dw, polypore;
- Polysporina simplex* (Davies) Vězda – 12, 51; stones;
- Porina aenea* (Wallr.) A. Zahlbr. – 4, 31, 32, 33, 54, 67; Aps, Cb, Fr, Fs, Pav;
- P. chlorotica* (Ach.) Hafellner & Kalb – 10, 35; stones;
- Porpidia crustulata* (Ach.) Hertel & Knoph – 26, 51, 52; wall, stones;
- P. macrocarpa* (DC.) Hertel & A.J. Schwab – 31; stone;
- P. soredizodes* (Lamy ex Nyl.) J. R. Laundon – 10, 15; stone;
- P. tuberculosa* (Sm.) Hertel & Knoph – 10, 51, 52; stones, wall;
- Pseudevernia furfuracea* (L.) Zopf – 2, 3, 6, 7, 13, 14, 16, 19–21, 29, 31, 36, 37, 39, 40, 42, 44, 46, 52, 62, 66; Apl, Bet, Fs, La, Pa, Pm, Ps, metal, dw; (Izydorek 2010);
- Psilolechia lucida* (Ach.) M. Choisy – 51, 63, 67; stone, walls;
- Punctelia jeckeri* (Roum.) Kalb – 15, 51, 60; Apl, Fr;
- P. subrudecta* (Nyl.) Krog – 15; Apl, Fr;
- Pycnora sorophora* (Vain.) Hafellner – 40, 43, 44, 47; Ps, dw;
- Pyrenula nitida* (Weigel) Ach. – 8; Cb;
- Pyrrhospora quernea* (Dicks.) Körb. – 18; Fr;
- Ramalina baltica* Lettau – 61, Apl;
- R. farinacea* (L.) Ach. – 2, 3, 6, 7, 9, 11 (c.ap.), 15, 18, 22, 29, 36, 38, 41, 42, 45, 49, 51–53, 55, 56, 58, 60, 61, 64–67; Apl, Aps, Cav, Fr, Fs, Pm, Ps, Pt, Q, Tc, U, dw, stone; (Izydorek 2010);
- R. fastigiata* (Pers.) Ach. – 11, 15, 38, 41, 45, 48, 51, 55, 56, 58, 59, 60, 61, 65–67; Apl, Aps, Cav, Fr, Q; (Izydorek 2010);
- R. fraxinea* (L.) Ach. – 11, 40, 41, 45, 48, 51, 56, 58, 59, 61, 65–67; Apl, Aps, Fr, Po, Q; (Izydorek 2010);
- Reichlingia leopoldii* Diederich & Scheid. – 20; Q;
- Rhizocarpon geographicum* (L.) DC. – 51; głaz, wall;
- Rinodina oleae* Bagl. – 15, 67; walls;
- R. pyrina* (Ach.) Arnold – 45; Aps;
- Ropalospora viridis* (Tønsberg) Tønsberg – 3, 4, 17, 20–22, 26–28, 30–35, 38; Aps, Bet, Cav, Cb, Fs, Q;
- Rusavskia elegans* (Link) S.Y. Kondr. & Karnefelt – 24, 52; Apl, concrete;
- Scoliciosporum chlorococcum* (Graewe ex Stenh.) Vězda – 22, 31, 42, 43, 46, 52, 53, 62; Bet, Pa, Pm, Ps, dw; (Izydorek 2010);
- S. sarothamnii* (Vain.) Vězda – 37; Pa;
- S. umbrinum* (Ach.) Arnold – 15, 23, 37, 42, 51, 52; stones, wall;
- Strangospora pinicola* (A. Massal.) Koerb. – 39, 52; dw;
- Tephromela atra* (Huds.) Hafellner – 15; stone;
- Trapelia coarctata* (Sm.) Choisy – 25, 29, 31, 55, 67; stone, walls;
- T. obtegens* (Th. Fr.) Hertel – 15; stone;
- T. placodioides* Coppins & P. James – 10, 52; stones;
- Trapeliopsis aeneofusca* (Flörke) Coppins & P. James – (acc. Izydorek 2010);
- T. flexuosa* (Fr.) Coppins & P. James – 2, 3, 14, 21, 26, 31, 36, 40, 42–44, 46, 52; Bet, Ps, dw; (Izydorek 2010);
- T. granulosa* (Hoffm.) Lumbsch – 2, 3, 13, 14, 19, 20, 22, 31, 37, 39, 40, 43, 46, 55, 62, 68; Bet, Ps, Q, dw; (Izydorek 2010);
- T. pseudogranulosa* Coppins & P. James – 20, 24, 26, 29, 31, 34, 35; Bet, gl, dw;
- Usnea dasopoga* (Ach.) Röhl. – 3, 6, 32, 36, 38, 52; Bet, Fs, La, Pa, Q; (Izydorek 2010);
- U. florida* (L.) Weber ex F.H. Wigg. – 3, 6, 9, 13, 32, 37, 39, 40, 43, 44, 47, 52, 62; Bet, Fr, La, Pa, Ps, Q, dw; (Izydorek 2010);
- U. hirta* (L.) Weber ex F.H. Wigg. – 19, 37, 39, 40, 43, 44, 47, 52, 62; Bet, La, Ps, Q, dw; (Izydorek 2010);
- Varicellaria hemisphaerica* (Flörke) Schmitt & Lumbsch – 9, 17, 26–29, 38; Al, Fs, Q; (Izydorek 2010);
- Verrucaria nigrescens* Pers. – 47, 51, 52; wall, stone; (Izydorek 2010);
- V. praetermissa* (Trevis.) Anzi – 8; stone;
- Violella fucata* (Stirt.) T. Sprib. – 2, 3, 8, 13, 16, 18–20, 24–27, 30, 31, 33–36, 39, 40, 42–44, 47; Al, Bet, Cav, Cb, Pa, Fs, Ps, Q, dw, twigs of *Ledum palustre*;
- Vulpicida pinastri* (Scop.) J.E. Mattsson & Lai – 6, 22, 46; La, Pa; (Izydorek 2010);
- Xanthoparmelia conspersa* (Ach.) Hale – 12, 26, 48, 51; stones;
- X. delisei* (Duby) Essl. – 50, 51; stones;
- X. loxodes* (Nyl.) O. Blanco et al. – 15, 27; stones;
- Xanthoria parietina* (L.) Th. Fr. – 1, 7, 9, 11, 15, 18, 19, 36, 40, 42, 45, 48, 49, 51, 52, 54–56, 60–69; Apl, Aps,

Bet, Fr, Po, Ps, Pt, Ra, Q, Sal, Tc, U, dw, metal, wall, eternit; (Izydorek 2010);
Zwackhia viridis (Ach.) Poetsch & Schied. – 8, 18, 20, 22; Cb, Fr, Fs, Q, Tc; (Izydorek 2010).

5. Conclusions

The Karnieszewice Forest Division, despite forest management carried out since many years, is relatively in a good condition and characterized by high average of the tree stand age. Additionally, in this area a strong air pollution were never detected. Simultaneously, habitat conditions (among others high air and substrates humidity) are very convenient for the lichen vegetation, especially for epiphytic species. Presented results can prove that in this case impact of forest management on lichen biota is not as drastic as it has been described in the literature. However, there are not many lichens typical for primary forests or they are rare.

In our opinion, rational forest culture based on ecological rules and preservation of forest continuity on larger areas enable the occurrence of numerous stenotopic lichen species, sensitive to changes of habitat conditions.

Acknowledgements

The authors would like to express their gratitude to workers of the Karnieszewice Forest Inspectorate, especially Dagny Nowak-Staszewska, Jarosław Staniszewski and Tomasz Kapustyński, for their help during the fieldwork and financial support of this work. We are thankful to Reviewers for their favorable review.

References

- Adamska E., 2014, Biota porostów Torunia na tle warunków siedliskowych miasta. [Lichen biota in Toruń in relation to habitat conditions of the city], Wyd. Naukowe UMK, Toruń.
- Arcadia L., 2013, *Usnea dasopoga*, a name to be reinstated for *U. filipendula*, and its orthography, *Taxon* 62(3): 604–605.
- Arcadia L. & Knudsen K., 2012, The name *Myriospora* is available for the *Acarospora smaragdula* group, *Opuscula Philolichenum* 11: 19–25.
- Arup U. & Sandler Berlin E., 2011, A taxonomic study of *Melanelixia fuliginosa* in Europe, *Lichenologist* 43(2): 89–97.
- Arup U., Söchting U. & Frödén P., 2013, A new taxonomy of the family Teloschistaceae, *Nord. J. Bot.* 31: 16–83.
- Bezmiān, A., 1998, Porosty Koszalina na tle warunków klimatyczno-antropogenicznych [Lichens of Koszalin in climatic and anthropogenic terms], [in:] J. Miadlikowska (ed.) *Botanika polska u progu XXI wieku, Materiały sympozjum i obrad sekcji 51 Zjazdu Polskiego Towarzystwa Botanicznego*, Gdańsk, 15–19 września 1998, [Polish botany on the threshold of the XXI century, Proceedings of the 51 Congress of the Polish Botanical Society, Gdańsk, 15–19 September 1998], Wyd. Nauk. Boguccy, Poznań: 36.
- Brodo I. M., Owe-Larsson B. & Lumbsch H. T., 1994, The sorediate, saxicolous species of the *Lecanora subfuscata* group in Europe, *Nord. J. Bot.* 14(4): 451–461.
- Cieśliński S., 2003, *Atlas rozmieszczenia porostów (Lichenes) w Polsce północno-wschodniej* [Distribution atlas of lichens (Lichenes) in north-eastern Poland], *Phytocoenosis* 15(N.S.), Suppl. *Cartographiae Geobotanicae* 15: 1–430.
- Cieśliński S., Czyżewska K. & Fabiszewski J., 2006, Red list of the lichens in Poland, [in:] Z. Mirek, K. Zarzycki, W. Wojewoda, Z. Szeląg (eds) *Red list of plants and fungi in Poland*, W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków: 71–89.
- Czyżewska K., 1976, Zanikanie porostów epifitycznych pod wpływem antropogenicznej degeneracji lasów liściastych Puszczy Pilickiej [Dissapearance of epiphytic lichens under the influence of antropogenic degenerations of broadleaved forests in the Pilicka Forest complex], *Phytocoenosis* 5(3–4): 363–375.
- Ekman S. & Svensson M., 2014, *Brianaria*, a new genus to accommodate the *Micarea sylvicola* group, *The Lichenologist* 46: 285–294.
- Erichsen C.F.E., 1936, Pertusiaceae, [in:] Rabenhorst's *Kryptogamen-Flora von Deutschland, Österreich und der Schweiz* 9.5(1): 321–728, Leipzig.
- Erichsen C. F. E., 1940, *Lichenologische Beiträge*. III, *Ann. Mycol.* 38.2–4: 303–331.
- Ertz D. & Tehler A., 2011, The phylogeny of Arthoniales (Pezizomycotina) inferred from nucLSU and RPB2 sequences, *Fung. Diversity* 49(1): 47–71.
- Fałtynowicz W., 1992, The lichens of Western Pomerania (NW Poland), An ecogeographical study, *Polish Bot. Stud.* 4: 1–182.
- Fałtynowicz W., 1995, Wykorzystanie porostów do oceny zanieczyszczenia powietrza [Use of lichens for estimating of air pollution], Fundacja CEEW, Krośno.
- Fałtynowicz W. & Kukwa M., 2006, Lista porostów i grzybów naporostowych Pomorza Gdańskiego [List of lichens and lichenicolous fungi of Gdańsk Pomerania], *Acta Bot. Cassubica, Monographiae* 2: 1–98.
- Hawksworth D. L., 1971, Lichens as litmus for air pollution: a historical review, *Intern. J. Environmental Studies* 1: 281–296.

- Herzig R. & Urech M., 1991, Flechten als Bioindikatoren. Integriertes biologisches Messystem der luftverschmutzung für Schweizer Mitteland, Biblioth. Lichenol. 43: 1–283.
- Izydorek I., 2010, Porosty Wysoczyzny Polanowskiej [The lichens of Wysoczyzna Polanowska district], Słupskie Prace Biol. 7: 51–78.
- Jóźwiak M.A., 2014, Wykorzystanie organizmów wskaźnikowych w bioindykacji środowisk lądowych i wodnych na wybranych przykładach [The use of indicative organisms in bioindication of land and water environments with the chosen examples], Kieleckie Tow. Nauk., Kielce.
- Kalb K., 2007, New or otherwise interesting lichens, Biblioth. Lichenol. 95: 297–316.
- Krawiec F., 1934, Flora epityczna lasów bukowych Wielkopolski [Die Epiphyten Flora der Buchenwälder Grosspolens], Acta Soc. Bot. Pol. 11 (Suppl.): 317–327.
- Kubiak D., Czarnota P., Zduńczyk A., Dynowska M., Leśnianski G., Grabowska A., Olszewska S., Sadowska-Deś A. & Wojdal P., 2014, The preservation status of the lichen biota in the designed Special Area of Conservation NATURA 2000 „Middle Łyna River Valley – Smolajny” (the Forest Division of Wichrowo), Acta Mycol. 49(1): 135–146.
- Kukwa M., Kowalewska A., Śliwa L., Czarnota P., Czyżewska K., Flakus A., Kubiak D., Wilk K., Dimos-Zych M., Kolanko K., Szymczyk R., Lipnicki L., Adamska E., Bielec D., Guzow-Krzemińska B., Gruszka W., Hachułka M., Jabłońska A., Oset M., Kiszka J., Kozik J., Leśnianski G. & Lazarus M., 2012, Porosty i grzyby naporostowe Wdzydzkiego Parku Krajobrazowego (Pomorze Gdańskie, N Polska) [Lichens and lichenicolous fungi of the Wdzydzki Landscape Park (Pomerze Gdańskie, N Poland)], Acta Bot. Cassubica 11: 75–103.
- Kukwa M., Fałtynowicz W., Kowalewska A., Szymczyk R., Adamska E. & Pietrzykowska K., 2013, Materiały do biaty porostów i grzybów naporostowych nadleśnictwa Karnieszewice i okolic (Pomorze Zachodnie), Piąte Dni Różnorodności Biologicznej w Leśnym Kompleksie Promocyjnym Lasy Środkowopomorskie [Materials to the lichen biota and lichenicolous fungi of the Karnieszewice Forest Division (Western Pomerania)], Wyd. EKWITA, Gdańsk: 9–17.
- Laundon J. R., 2010, *Lecanora antiqua*, a new saxicolous species from Great Britain, and the nomenclature and authorship of *L. albescens*, *L. conferta* and *L. wallalis*, Lichenologist 42(6): 631–636.
- Lendemer J. C., 2011, A standardized morphological terminology and descriptive scheme for *Lepraria* (Stereocaulaceae), Lichenologist 43(5): 379–399.
- Łubek A., 2007, Antropogeniczne przemiany biaty porostów Świętokrzyskiego Parku Narodowego i otuliny [Anthropogenic changes of lichen biota of the Świętokrzyski National Park and its protective zone], Fragm. Flor. Geobot., Ser. Polonica, Supplementum 10: 3–94.
- Motyka J., 1934, W sprawie ochrony porostów [On the protection of lichens], Ochr. Przyr. 14: 50–56.
- Neuwirth G. & Aptroot A., 2011, Recognition of four morphologically distinct species in the *Graphis scripta* complex in Europe, Herzogia 24(2): 207–230.
- Nordin A., Savić S. & Tibell L., 2010, Phylogeny and taxonomy of *Aspicilia* and Megasporaceae, Mycologia 102(6): 1339–1349.
- Nylander W., 1866, Les lichens du Jardin du Luxembourg, Bull. Soc. Bot. Fr. 13: 364–372.
- Orange A., James P. W. & White F. J., 2001, Microchemical methods for the identification of lichens, British Lichen Society, London.
- Palice Z., Printzen C., Spribille T. & Elix J. A., 2011, Notes on the synonyms of *Lecanora filamentosa*, *Graphis Scripta* 23(1): 1–7.
- Rozporządzenie Ministra Środowiska z dnia 16 października 2014 r. w sprawie ochrony gatunkowej grzybów, Dz. U., poz. 1408 [Regulation of the Minister of the Environment of 16 October 2014 on the protection of fungi species, Journal of Laws no. 1408], 2014.
- Santesson R., Moberg R., Nordin A., Tønberg T. & Vitikainen O., 2004, Lichen-forming and lichenicolous fungi of Fennoscandia, Museum of Evolution, Uppsala University, Uppsala.
- Schiefelbein U., Czarnota P., Thüs H. & Kukwa M., 2012, The lichen biota of the Drawieński National Park (NW Poland, Western Pomerania), Folia Cryptog. Estonica 49: 59–71.
- Schmitt I., Otte J., Parnmen S., Sadowska-Deś A.D., Lücking R. & Lumbsch H. T., 2012, A new circumscription of the genus *Varicellaria* (Pertusariales, Ascomycota), MycoKeys 4: 23–36.
- Spribille T., Goffinet B., Klug B., Muggia L., Obermayer W. & Mayrhofer H., 2011, Molecular support for the recognition of the *Mycoblastus fucatus* group as the new genus *Violella* (Tephromelataceae, Lecanorales), Lichenologist 43(5): 445–466.
- Tobolewski Z., 1964, Lichenotheca Polonica, Fasc. XVII, No. 376–400, Lichenes Poloniae Septentrionali-occidentalis, Wyd. PAN, Poznań.
- Tobolewski Z., 1966, Rodzina Caliciaceae (Lichenes) w Polsce [The family Caliciaceae (Lichenes) in Poland], Prace Komis. Biol. Pozn. Tow. Przyj. Nauk 24(5): 1–101.
- Tobolewski Z., 1979, Porosty (Lichenes) 5, Atlas rozmięzczenia roślin zarodnikowych w Polsce, ser. III [Atlas of geographical distribution of spore-plants in Poland, ser. III.], PWN, Warszawa-Poznań.

- Tobolewski Z., 1980, Porosty (Lichenes) 6, Atlas roz-
mieszczenia roślin zarodnikowych w Polsce, ser. III
[Atlas of geographical distribution of spore-plants in
Poland, ser. III], PWN, Warszawa-Poznań.
- Tobolewski Z., 1981, Porosty (Lichenes) 7, Atlas roz-
mieszczenia roślin zarodnikowych w Polsce, ser. III
[Atlas of geographical distribution of spore-plants in
Poland, ser. III], PWN, Warszawa-Poznań.
- van Herk C. M., 2001, Bark pH and susceptibility to toxic
air pollutants as independent causes of changes in epi-
phytic lichen composition in space and time, *Lichenolo-
gist* 33(5): 419–441.
- Wieczorek A., 2005, Biota porostów Szczecińskiego Parku
Krajobrazowego [Lichen flora of the Szczecin Land-
scape Park], *Fragm. Flor. Geobot. Polonica* 12(1):
143–156.
- Wieczorek A., 2014, Operat ochrony porostów Wolińskie-
go Parku Narodowego [Statement of lichens protection
in the Woliński National Park], Mscr., Szczecin.
- Zalewska A., 2012, Ecology of lichens of the Puszcza
Borecka Forest (N Poland), W. Szafer Institute of Bot-
any, Polish Academy of Sciences, Kraków.
- Zalewska A., Fałtynowicz W., Krzysztofiak A., Krzysztofi-
ak L. & Picińska-Fałtynowicz J., 2004a, Lichens of the
Suwalski Landscape Park, [in:] A. Zalewska, W. Fał-
tynowicz (eds) *Lichens of the protected areas in the
Euroregion Niemen, Stowarzyszenie „Człowiek i Przy-
roda”*, Suwałki: 5–50.
- Zalewska A., Fałtynowicz W., Krzysztofiak A., Krzysztofi-
ak L. & Picińska-Fałtynowicz J., 2004b, Lichens of
Romincka Primeval Forest, [in:] A. Zalewska, W. Fał-
tynowicz (eds) *Lichens of the protected areas in the
Euroregion Niemen, Stowarzyszenie „Człowiek i Przy-
roda”*, Suwałki: 51–109.