

**RZADKOWSKI, Kamil and MROZKOWIAK, Mirosław. Sexual and environmental dimorphism of the frequency of significant correlations of body mass and height with feet features among 7-13-year-old adolescents. Journal of Education, Health and Sport. 2023;33(1):78-90. eISSN 2391-8306. DOI <http://dx.doi.org/10.12775/JEHS.2023.33.01.009>
<https://apcz.umk.pl/JEHS/article/view/42090>
<https://zenodo.org/record/8200476>**

The journal has had 40 points in Ministry of Education and Science of Poland parametric evaluation. Annex to the announcement of the Minister of Education and Science of 17.07.2023 No. 32318. Has a Journal's Unique Identifier: 201159. Scientific disciplines assigned: Physical Culture Sciences (Field of Medical sciences and health sciences); Health Sciences (Field of Medical Sciences and Health Sciences). Punkty Ministerialne z 2019 - aktualny rok 40 punktów. Załącznik do komunikatu Ministra Edukacji i Nauki z dnia 17.07.2023 Lp. 32318. Posiada Unikatowy Identyfikator Czasopisma: 201159. Przypisane dyscypliny naukowe: Nauki o kulturze fizycznej (Dziedzina nauk medycznych i nauk o zdrowiu); Nauki o zdrowiu (Dziedzina nauk medycznych i nauk o zdrowiu).
© The Authors 2023;
This article is published with open access at License Open Journal Systems of Nicolaus Copernicus University in Torun, Poland
Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author (s) and source are credited. This is an open access article licensed under the terms of the Creative Commons Attribution Non commercial license Share alike. (<http://creativecommons.org/licenses/by-nc-sa/4.0/>) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.
The authors declare that there is no conflict of interests regarding the publication of this paper.
Received: 05.06.2023. Revised:30.07.2023. Accepted: 31.07.2023. Published: 08.08.2023.

Sexual and environmental dimorphism of the frequency of significant correlations of body mass and height with feet features among 7-13-year-old adolescents

Author: ¹Kamil Rzadkowski^{CDEF}, ²Mirosław Mrozkowiak^{ABG}

Place of work: ¹Chiropractic School of Poland, Świecie, Poland

²Physiotherapy Clinic AKTON, Warsaw, Poland

A – preparation of research project

B – data collection

C – statistical analysis

D – data reading

E – preparation of manuscript

F – literature elaboration

G – funds obtaining

Summary

The aim of the study is to demonstrate the sexual and environmental dimorphism of the frequency of significant correlations between body weight and height with selected feet features among 7–13-year-old children.

Material and method. The research among 7-13-year-old children was being carried out over a period of 3 years in 6 semi-annual editions and allowed to register 16,462 observations of the value of 48 features describing feet, weight, and height of the body. The photogrammetric method was used to measure selected features.

Conclusions

1. Among 7-13-year-old children, the body height is more often associated with the features of the feet than the body weight.
2. Among girls, the proportions of the frequency of body height correlations are similar to the percentage of boys, where body weights are more frequent. The mass and body height shows more frequent associations with feet features among children living in the city than among children from the countryside.
3. The most frequent significant correlations between body weight and height and the features of the feet occur among boys living in the city. Whereby the percentage of body weight correlations in the rural environment is very low. The frequency of body weight and height correlations exceeds that percentage of correlations among girls, who live in the city than in those from rural environment.

Keywords: body weight and height, feet features.

1. Introduction

The literature review of the topic shows that authors often focus on the correlation between body mass and feet architecture in various aspects [1, 2, 3]. Feet are a significant static-dynamic link in the musculoskeletal system, as they keep the body weight within the actual limit of stability, being the propulsion at the same time [4]. The issue of the interdependence of feet and somatic features most often concerns the relationship of longitudinal arch of the feet [5, 6]. The observed high percentage of postural static disorders is often associated not only with the sedentary lifestyle, but also with body weight and height. The student's environment significantly influences the type and size of disorders in body posture, especially during the school period [7, 8]. A study by Drzał-Grabiec et al. on body posture, weight, and height in a group of 563 students from the first three grades of a primary school showed statistically significant differences in the characteristics describing body posture, body weight and height [9]. Research by Pauk et al. among 450 primary school students on the verification of the influence of selected epidemiological factors on the occurrence of flat feet in children, which was to lead to the development of a mathematical model of the dependence of the height of the longitudinal arch of the foot on such factors as age, gender, Cole's weight-height index and place of residence showed that age and Cole's index had the greatest influence on the height of the longitudinal arch of the foot. It has also been proved that the arch height increases with age whereas the Cole's index decreases [10]. Research by Puszczalowska-Lizis et al. in a group of 50 second graders of a primary school showed that girls in the early school period have better arched feet than boys, while in terms of heel angle, both groups are similar to each other. It has also been proved that in both girls and boys, body weight is a feature that significantly influences the shape of the longitudinal arch of the foot, as well as the presence of correlations between the Clarke's angle and body height in boys, and their absence in girls, probably results from sexual dimorphism. Moreover,

it was observed that body weight and height did not affect the transverse arch of the foot in early school children [11]. Studies by Mikołajczyk and Jankowicz-Szymańska in a group of 90 children aged 7 have shown that static flat feet are a defect often occurring in 7-year-olds, especially boys, and the fat condition and position of the lower limbs do not significantly affect the degree of arching of the feet. Additionally, it has been shown that the excessive amount of adipose tissue, visible in an increased body weight and in the thickness of the abdominal skin-fat fold, generates the probability of valgoid knee positioning in children of both sexes [3].

The aim of the research is to show the frequency of significant correlations between body weight and height with selected feet features among 7-13-year-old children. The analysis of the study results was heading in four directions. The first is the answer to the question: which feet features have got the most frequently significant correlation with weight and height at the age of 7-13? The second and third are the answer to the question: which features do most often show a significant correlation in terms of sexual and environmental dimorphism? The fourth is the answer to the question: what is the frequency of significant correlations of weight and height among girls and boys within each of the environments?

2. Material and method

The research among children aged 7-13 was conducted over a 3-year period in 6 semi-annual editions and allowed to register 16,462 observations including 8953 girls and 7509 boys. For statistical analysis, body weight and height as well as 48 angular and linear feet parameters were selected for individual age categories, gender and environment, Tab. 1. Due to the limited volume of the work, a detailed description of the somatic features of the research material and the research results obtained can be found in the author's monograph [12].

The habitual posture was basically assumed in the research to be always assessed as a relatively permanent individual characteristic of a human being. This posture reflects the individual emotional, mental, and social state of the respondent. It is the most accurate in describing the silhouette in time and place. The performed diagnostics does not determine whether the individual's posture is correct, they only present the state of its ontogenetic realization. Objectivized and comparable test results will make it possible to register the parameters adopted for the analysis with possible to define compensations. The combination of a torso and feet examination makes it possible to objectively determine the quality of the posture pattern realized in a particular environment, gender, and age category. The applied measuring device determines several dozens of parameters describing the feet. It is possible to obtain a spatial image thanks to displaying lines with strictly defined parameters. The lines falling on the skin are distorted depending on the configuration of the surface. The use of a lens makes the image of the feet possible to be picked up by a special optical system with a camera, and then transferred to a computer monitor. Line image distortions recorded in the computer memory are processed by a numerical algorithm into a contour map of the tested surface. When performing the examination, one should know the photo records an image on the skin of the plantar surface of the child's feet [12].

Empirical data appeared to be quantitative and qualitative features (gender, place of residence, etc.). The calculations of the values of positional statistics (arithmetic mean, quartiles), dispersion parameter (standard deviation) and symmetry indices (asymmetry coefficient, cluster coefficient) give a full overview of the distribution of the examined features considering age groups, gender, and environment. Correlations and significance were determined using p-value, and the frequency as a percentage value.

Tab. 1. List of registered foot features, body weight and height

No	Symbol	Parametres					
		Label	Name	Description			
1	DL p	mm	Length of right foot (p), and left foot(l)	The distance between akropodion and pterion points on the platnogram			
2	DL l						
3	Sz p						
4	Sz l						
5	W p						
6	W l						
7	Alfa p m	degrees	Angle of right hallux valgus : Alfa p p, left: Alfa l p. Angle of right hallux varus Alfa p m, left: Alfa l m.	The angle between the straight line passing through the metatarsale tibiale and the innermost points at the medial edge of the heel and the straight line passing through the metatarsale tibiale and the innermost points at the medial edge of the toe			
8	Alfa p p						
9	Alfa l m						
10	Alfa l p						
11	Beta p m						
12	Beta p p						
13	Beta l m						
14	Beta l p						
15	Gamma p (Gam.P)				mm ²	Surface of the right foot(p), and left foot(l)	Foot plantogram surface
16	Gamma l (Gam.L)						
17	PS p	mm	Length of longitudinal arch of the right foot 1, 2, 3, 4, i 5 (P),and the left foot (L)	The length of the arch from the 1st, 2nd, 3rd, 4th and 5th metatarsal bones to the pterion point			
18	PS l						
19	DP 1						
20	DP 2						
21	DP 3						
22	DP 4						
23	DP 5						
24	DL 1						
25	DL 2						
26	DL 3						
27	DL 4						

28	DL 5			
29	WP 1		Height of arch 1, 2, 3, 4 i 5 of the right foot (P), and left foot (L)	Distance from the ground to the highest point of arch 1, 2, 3, 4 and 5.
30	WP 2			
31	WP 3			
32	WP 4			
33	WP 5			
34	WL 1			
35	WL 2			
36	WL 3			
37	WL 4			
38	WL 5			
39	SP 1		Width of arch 1, 2, 3, 4 i 5 of the right foot (P), and left foot (L)	Bowstring of the arch length 1, 2, 3, 4 i 5.
40	SP 2			
41	SP 3			
42	SP 4			
43	SP 5			
44	SL 1			
45	SL 2			
46	SL 3			
47	SL 4			
48	SL 5			
49	Mc	kg	Body weight	Measurements were made on a medical scale with an accuracy of 0.5 cm and 100 grams
50	Wc	cm	Body height	

Source: own research

3. Obtained results

Body weight is significantly related to the following features like length, width and plantar surface of the feet, the size of the transverse arch (Gamma angle) and the angle of the fifth hallux varus of the left foot, the length of the five longitudinal arches of the right foot and the first and fourth left feet, the width of the five longitudinal arches of the right and left foot, the height of the first, third, fourth and fifth longitudinal arch of the right foot and the height of the first, second, third and fourth of the left foot. The height of the body is significantly related to the length and width of the feet, the plantar surface, the angle of the fifth hallux valgus, and the length and width of the five longitudinal arches, the height of the five longitudinal arches of the feet, the size of the transverse arch (Gamma angle) and hallux valgus of the right foot.

The statistical analysis shows that among 7-13-year-old children, the frequency of significant correlations between body height and feet features is 83.33%, and body weight is

70.83%. The inference of the frequency of significant correlations only within gender shows that among boys the body height is 79.16%, and body weight is 45.83%, whereas among girls it is 81.25% and 62.5%, respectively. The profile of the analysis in terms of the environment showed that among individuals living in the city, the frequency of significant correlations between body height and feet features was 83.33%, and body weight was 60.41%, whereas among those living in the countryside it was 81.25% and 45.83%, respectively, Table 2, Fig. 1.

A more precise conception of the percentage of significant correlations within the environment showed that among boys from the city the body height was associated with feet features at the level of 75.0%, and body weight only at 41.66%, among those living in the countryside it was 70.83% and 6.25%, respectively. The frequency of significant correlations of body height among girls from urban areas is 81.25%, and body weight at 60.41%, whereas from rural areas it is at 77.08% and at 41.66%, respectively. Table 3, Fig. 2.

Tab. 2. Percentage of significant correlations between height and weight of the body with the features of the feet depending on age, sex, and environment (n) K = 8953, M = 7509

Analysis profile	Body height and weight	Percentage of correlation
Age from 4 to 6	Wc	83,33
	Mc	70,83
Sex	M Wc	79,16
	M Mc	45,83
	K Wc	81,25
	K Mc	62,5
Environment	M Wc	83,33
	M Mc	60,41
	W Wc	81,25
	W Mc	45,83

Source: own research

Legend of tab. 2, fig. 1: Wc - body height, Mc - body weight, versicle "Gender": M - male, F - female, versicle "Environment": M - city, W - village.

Tab. 3. Percentage of significant correlations between body height and weight with the feet features of male and female sex of both environments (n) K = 8953, M = 7509

Analysis profile	Body height and weight	Percentage of correlation
M M	Wc	75,0
	Mc	41,66
M W	Wc	70,83
	Mc	6,25
K M	Wc	81,25
	Mc	60,41
K W	Wc	77,08
	Mc	41,66

Source: own research

Legend of tab. 3, fig. 2: MM – male sex, city, MW – male sex, village, KM – female sex, city, KW – female sex, village.

4. Discussion

The presented correlations are important from the point of view of the ongoing posturogenesis in the selected age range. They should draw the attention of parents and teachers to the undoubtedly large influence of height and weight on posture features not only considering feet. Mrozkowiak's research [13] results in showing more frequent correlation between body height and feet than body weight and feet among 4-18-year-old children. Whereas the most frequent correlations are with body height independently on sex and environment among 4-13-year-old children, but body weight is among 7-13-year-old children. Among 4-13-year-old boys the body weight and height show significantly more frequent correlation with feet features than among 14-18-year-old boys. The correlation of body height and weight with feet features are more frequent among 7-13-year-old girls.

5. Conclusions

1. There is more frequent association between the height of the body and feet features than with body weight among 7-13-year-old children.

2. Body height is more frequently associated with features of the feet than body weight among boys and girls. The proportions of the frequency of correlations with body height among girls are similar to the percentage of boys, but body weight is more frequent. Moreover, there is more frequent association of body weight and height with feet features among children living in a city than those living in the countryside.

3. The most common significant correlations between body weight and height and the features of the feet occur among boys living in the city. Whereby the percentage of body weight correlations in the rural environment is very low. The frequency of body weight and height correlations among girls living in the city exceeds the percentage of those of the rural environment.

Literatura

1. E. Puszczalowska-Lizis, Związki pomiędzy wysklepieniem poprzecznym stóp a wybranymi cechami morfologicznymi u młodych osób dorosłych, *Fizjoterapia*, 2011, 19, 1, 3-9.
2. E. Barańska, E. Gajewska, M. Sobieska, Otyłość i wynikające z niej problemy zarządu ruchu a sprawność motoryczna dziewcząt i chłopców z nadwagą i otyłością prostą, *Nowiny Lekarskie*, 2012, 81, 4, 337-341.
3. E. Mikołajczyk, A. Jankowicz-Szymańska, Wpływ otłuszczenia na wysklepienie stóp i ukształtowanie kończyn dolnych u 7-latków, *Fizjoterapia* 2010, 18, 2, 10-20.
4. J. W. Błaszczak, *Biomechanika kliniczna*, Wyd. Lek. PZWL, Warszawa, 2004.
5. Yang S. M., Kayamo J., Noriamatsu T., Fujita M., Matsusaka N., Suzuki R., Okamura H. Dynamic changes of the arch of the foot during walking. *Biomechanics IX-A*. J. Hum. Kinet., 1985, 417-422.
6. Lizis P., Kasperczyk T., Szmigiel Cz., Całka-Lizis T., Emmerich W., Szczygieł P. Postawa ciała i jej związki z cechami morfologicznymi dzieci otyłych, [w:] J. Ślężyński

- (red.) Postawa ciała człowieka i metody jej oceny. AWF, Katowice 1992, 99-107.
7. Mrozkowiak M. Próba określenia znaczenia DOBREGO KRZESŁA w profilaktyce zaburzeń postawy ciała = Attempt to determine the importance of GOOD CHAIR in the prevention of body posture disorder's. Journal of Health Sciences. 2014;4(4):195-214.
 8. M. Mrozkowiak, H. Żukowska, Znaczenie Dobrego Krzesła, jako elementu szkolnego i domowego środowiska ucznia, w profilaktyce zaburzeń statyki postawy ciała = The significance of Good Chair as part of children's school and home environment in the preventive treatment of body statistics distortions. Journal of Education, Health and
 9. J. Drzał-Grabiec, S. Snela, B. Szczepanowska-Wołowicz, K. Bibrowicz, Badanie zależności pomiędzy wybranymi parametrami charakteryzującymi postawę ciała oraz masą i wysokością ciała a miejscem zamieszkania, Przegląd Medyczny Uniwersytetu Rzeszowskiego i Narodowego Instytutu Leków w Warszawie, Rzeszów 2012, 4, 410-417.
 10. J. Pauk, V. Ezerskiy, M. Rogalski, Wpływ czynników epidemiologicznych na wystąpienie stopy płaskiej u dzieci, Fizjoterapia, 2010, 18, 2.
 11. E. Puszczałowska-Lizis, T. Ridan, M. Ogarek, Charakterystyka parametrów wysklepienia podłużnego i poprzecznego stóp dziewcząt i chłopców w okresie wczesnoszkolnym, Młoda Sportiwna Nauka, Ukraina, 2011, t. 3, 234-239.
 12. M. Mrozkowiak, Modulacja, wpływ i związki wybranych parametrów postawy ciała dzieci i młodzieży w wieku od 4 do 18 lat w świetle metody projekcyjnej, Wydawnictwo Uniwersytetu Kazimierza Wielkiego, Bydgoszcz. I, t. II.
 13. Mrozkowiak M., Jakże związki mają masa i wysokość ciała z cechami stop wśród dzieci i młodzieży w wieku 4-18 lat, Fizjoterapia Funkcjonalna, 2018, 3(12), 5-12, ISSN 2543-9545.




Fig.1 The percentage of significant correlations of the body weight and height w feet features depending on the range of age, sex, and environment (n) K=8953, M=7509




Fig.2 The percentage of the body weight and height of females and males depending on environment (n) K=8953, M=7509