Wilczyński Jacek, Habik Natalia, Wypych Żaneta. The effects of sensory integration techniques on muscle tone in children after prenatal exposure to alcohol. Journal of Education, Health and Sport. 2018;8(8):347-358. eISNN 2391-8306. DOI http://dx.doi.org/10.5281/zenodo.1312425

http://ojs.ukw.edu.pl/index.php/johs/article/view/5660

The journal has had 7 points in Ministry of Science and Higher Education parametric evaluation. Part B item 1223 (26/01/2017). 1223 Journal of Education, Health and Sport eISSN 2391-8306 7

© The Authors 2018;

This article is published with open access at Licenses Open Journal Systems of Kazimierz Wielki University in Bydgoszcz, Poland Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author (s) and source are credited. This is an open access article licensed under the terms of the Creative Commons.org/licenses/by-nc-sa/4.0/) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.

The authors declare that there is no conflict of interests regarding the publication of this paper.

Received: 25.06.2018. Revised: 28.06.2018. Accepted: 15.07.2018.

The effects of sensory integration techniques on muscle tone in children after prenatal exposure to alcohol

Jacek Wilczyński¹, Natalia Habik², Żaneta Wypych²

- 1. Department Posturology, Hearing and Balance Rehabilitation, Institute of Physiotherapy, Faculty of Medicine and Health Sciences, Jan Kochanowski University in Kielce, Poland;
- 2. Żaneta Wypych, student, Institute of Physiotherapy, Faculty of Medicine and Health Sciences, Jan Kochanowski University in Kielce, Poland
- 3. Natalia Habik: Ph.D. student, Faculty of Medicine and Health Sciences, Jan Kochanowski University in Kielce, Poland

Correspondence: Assoc. Prof. UJK Jacek Wilczyński, Ph.D, Head of Posturology Department, Hearing and Balance Rehabilitation, Institute of Physiotherapy, Faculty of Medicine and Health Sciences, Jan Kochanowski University, Kielce, Al. IX Wieków Kielc 19, 25-317 Kielce, Poland. Tel: 48-603-703-926. E-mail: jwilczyński@onet.pl

ABSTRACT

Background: Prenatal exposure to alcohol is the cause of Foetal Alcohol Syndrome (FAS). Alcohol exerts an especially negative effect on the central nervous system of an unborn child. Children with FAS frequently have problems with sensory integration. The objective of the study was an evaluation of the effectiveness of SI therapy in children with Foetal Alcohol Syndrome on the example of muscle tension. **Methods:** The study was conducted in 2012 at the Centre of Advanced Physiotherapy in Kielce, and covered a group of 20 children aged 4-5 who had a history of prenatal exposure to alcohol. The children systematically participated in a one-hour SI therapy session, once a week for the period of 6 months. **Results:** A high effectiveness of SI therapy was confirmed in children after prenatal exposure to alcohol based on the example of muscle tension (p<0.001). The positive effects of therapy were proved for the following tests: in prone extension position (p<0.001), in supine, flexed position (p<0.05), ATOS (p<0.001), STOS (p<0.001), the Schilder test (p<0.001) and trunk stabilization (p<0.001). **Conclusions:** The sensory integration technique exerts a beneficial effect on the normalization of muscle tone in children after prenatal exposure to alcohol.

Key words: therapy using sensory integration technique, muscle tone, prenatal exposure to alcohol

INTRODUCTION

Foetal Alcohol Syndrome (FAS) is the consequence of the effects of alcohol on the foetus during the prenatal development period [1]. Alcohol has an especially negative effect on the central nervous system of an unborn child [2,3]. It also results in abnormalities in physical development (face, lips, eyes, body size), difficulties in cognitive development, behavioural disorders, memory deficits and ADHD syndrome [4,5]. FAS is the sum of abnormalities in physical and mental development, resulting in challenges for the future development of a child, as well as for parents and society [6,7]. It is characterized by psychophysical disorders in the form of mental retardation, brain dysfunction, abnormalities in general development, psychomotor and physical abnormalities, disorders in learning and shaping personality [8,9]. Children with FAS often experience problems with sensory integration [10,11]. A child with SI disorders may behave in a strange way, incomprehensible for the environment [12,13]. If the child incorrectly receives and processes tactile stimuli, s/he may hate hair-washing, may pat or scratch without evident reason, fidget in a chair, avoid close contact or demand this contact more often than peers [14,15]. If any abnormalities occur in the vestibular system responsible for balance reactions, the child may be afraid of stairs, dislike swings, seek assistance when the ground changes, or to the contrary, perform locomotor manoeuvres which make others dizzy or sick by just looking at them [16]. With abnormal proprioceptive sensations, the child may maintain in constant motion: running instead of walking, jumping instead of sitting or may assume very strange positions [17]. Muscle tone is a constantly maintained state of contraction of the muscles, which is affected by many factors - mainly one's own response of the muscle to stretching [18]. The physiological mechanisms exerting an effect on the regulation of the tone are very complex [19]. The most important receptors

related with muscle tone are proprioceptors. They include a neuromuscular spindle (in muscles) and Golgi organs (in tendons). Contraction of a muscle causes an increase in its tone, which stimulates the spindles and Golgi organs. In this way, a closed circuit is created, spontaneously evoked by impulses from the nerve endings [20]. Apart from the abovementioned feedback, muscle tone is also affected by impulses constantly flowing from the cerebral cortex and subcortical nuclei. Their action consists mainly of the inhibition and regulation of muscle excitability. Normal muscle tone should be high enough to be able to resist the force of gravity, and low enough that the movements can be freely performed [21]. Physiological muscle tone may differ and assume lower or higher values. Abnormal muscle tone impairs the psychophysical development of a child [22]. The objective of this study was an evaluation of the effectiveness of SI therapy in children after prenatal exposure to alcohol, based on the example of muscle tone.

MATERIAL AND METHOD

The study was conducted in a group of 20 children (6 girls and 14 boys), aged 4-5 years, with a history of prenatal exposure to alcohol. The majority of children came from adoptive and foster families or orphanages. The study was conducted in 2012 at the Centre of Advanced Physiotherapy in Kielce (Poland). All procedures performed in tests involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards. The guardians of the children were informed about the purpose of the study and expressed written consent for their children's participation in the study. The study was non-invasive and free of charge. The patients willingly participated in the study and perceived it as a concern about their state of health. All children were diagnosed with SI disorders and dysfunctions typical of children with FAS spectrum, including: hyperactivity, pertinacity, passiveness, impulsivity, sleep problems, malice, hypersensitivity to touch and sound, adaptation difficulties, problems with organization, low self-esteem, difficulty with self-control and ease of falling into depressive states. The examination of sensory integration disorders consisted of three 50-minute diagnostic meetings. The diagnosis of SI included: an interview with a parent/caregiver, observation of controlled and spontaneous movements of the child in the therapeutic room, response to the basic sensory stimuli administered, as well as clinical observation. The evaluation included gross and fine motor skills, muscle tone and balance responses of the body. The presented study comprehensively discusses the effect of SI on muscle tone. While evaluating muscle tone, 8 trials with clinical observations were performed according to Ayres: cocontraction, prone extension position, supine flexed position,

ATOS, STOS, muscle tone, the Schilder test and trunk stabilisation. Each child systematically participated in a one-hour SI therapeutic session once a week for 6 months. The therapeutic programme covered: normalization of the vestibular, proprioceptive and tactile systems, normalization of muscle tone, development of motor planning, oculomotor performance and motor coordination, hand therapy, integration of ATOS, STOS, development of locomotion and balance functions, as well as improvement of the efficacy of gross and fine motor skills. Measurement data were collected using the Microsoft MS EXCEL spreadsheet, and after preliminary processing, imported to Statistica StatSoft software. Data analysis included anthropometric characteristics (height, body mass, BMI) and tests for SI. Basic statistical parameters were calculated, such as: arithmetic mean (x), standard deviation (s), median value (Me), skewness (Sk), and kurtosis (Ku), as well as extreme values and confidence intervals for the level of 95%. The variables were verified from the aspect of normality of distribution using the Shapiro-Wilk test. In addition, Cronbach's Alpha analysis of reliability was also performed. In order to reveal the differences between studies prior to and after SI therapy, the Student's *t*-test and the Wilcoxon matched pairs test were applied.

RESULTS

The mean age of children was (x = 4.45), standard deviation (s = 0.51); mean body height (x = 4.45)1.01), standard deviation (s = 0.03); body mass (x = 16.35), standard deviation (s = 1.34); mean BMI (x = 15.73), standard deviation (s = 0.52) (Tab. 1). The mean result in the test for co-contraction was the same prior to and after SI therapy and was 1.000; the mean result in the prone extension position was 0.700 before therapy and 0.900 after SI therapy, the difference 0.290; the mean result in the supine flexed position was 0.888 before therapy and 0.988 after SI therapy, the difference 0.100; the mean result in the ATOS test was 0.350 prior to therapy and 0.500 after SI therapy, the difference - 0.150; the mean result in the STOS test was 0.700 before therapy and 1.000 after SI therapy, the difference -0.300; the mean result for muscle tone was 0.850 before therapy and 0.850 after SI therapy; the mean result in the Schilder test was 0.4170 before therapy and 0.717 after SI therapy, the difference - 0.300; the mean result for trunk stabilisation was 0.500 before therapy and 0.767 after SI therapy, the difference 0.267; the overall result of muscle tone was 0.676 before therapy and 0.851 after therapy, the difference - 0.176 (Tab. 2). A non-significant result of the Shapiro-Wilk test confirmed that the distribution of the observed variable (muscle tone) was similar to normal distribution both prior to (p = 0.291) and after therapy (p = 0.118). The Cronbach's Alpha coefficient assume values from 0-1. The higher the value, the higher the reliability of the scale. It is assumed that values of over 0.7 mean the acceptable reliability of the scale.

Analysis of Cronbach's Alpha coefficients showed a high reliability of the applied tests, especially prior to therapy. The high effectiveness of SI therapy was confirmed in children after prenatal exposure to alcohol, based on muscle tone. Positive outcomes of SI therapy were obtained for the following tests: in prone extension position (p<0.001), in supine flexed position (p<0.05), ATOS (p<0.001), STOS (p<0.001), the Schilder test (p<0.001) and trunk stabilisation (p<0.001). The Student's *t*-test for muscle tone in the case of normalized values showed significant differences in tests conducted prior to and after SI therapy (p<0.001) (Tab. 3).

DISCUSSION

The report by the Birth Defects Monitoring Program conducted by the Centers for Disease Control (CDC) based on data from 1,500 hospitals showed that the incidence of the cases of FAS in the American population is 0.3-0.9 per 10,000 births [23]. In turn, Abel and Sokol, after reviewing 19 epidemiological reports from the whole world, estimated the incidence of FAS at 1.9 per 1,000 live births. According to the CDC reports, the rates of occurrence of FAS per 10,000 births differ according to the ethnic origin of their mothers. The incidence of FAS worldwide is estimated at 3-9 per 1,000 live births [24]. Knuiman, Rijk, Hoksbergen et al. [25] examined a group of children adopted from Poland by Dutch parents. The study included 121 children aged 6-17. Three groups were distinguished: children with the diagnosis of FAS (31%), children whose adoptive parents suspected FAS (21%), and children whose adoptive parents did not suspect FAS (49%). In children with the diagnosis of FAS, body height insufficiency was observed. The children attended special needs schools and had difficulties with social functioning. However, the features of FAS were also observed in the other two groups. Children adopted from Poland showed a high risk of FAS. The researchers presumed that the Dutch adoptive parents must have realized the potential consequences of prenatal exposure to alcohol in children adopted from Poland. It is estimated that in Poland, there are 900 cases of fully symptomatic FAS in infants annually, whereas, in 9,000 children, the occurrence of some of its symptoms is observed [25]. The study conducted at the Institute of Mother and Child in Warsaw, consisting in the examination of urine samples in pregnant women who declared abstinence for the presence of markers of alcohol consumption (betahexosaminidase and gamma-glutamyltranspeptidase), showed that every third woman consumed alcohol knowing that she was pregnant [26]. A survey carried out at 2005 in the Sopot Laboratory for Social Studies by order of the State Agency for the Prevention of

Alcohol-Related Problems (PARPA) in a representative group of 1,038 women at the reproductive age of 18-40, showed that every third woman consumed alcohol during pregnancy [27]. Alcohol was most frequently consumed by women with secondary school education, while most rarely by those with primary school education. The highest consumption of alcohol was observed among women from small and medium-size towns, whereas the lowest - among those from large cities and rural areas. In 2005, the IBSOS International Research Group Laboratory, by order of the Childbirth with Dignity Foundation, conducted a survey in a representative group of 1,000 Poles aged 15 and older. One-third of the women (33.5%) who had given birth to one or more children admitted that they had consumed small amounts of alcohol during pregnancy. Although 83% of respondents had heard that even the smallest amount of alcohol may be dangerous, one-third of them (31.7%) mentioned that a small amount of alcohol may exert a positive effect on the mother and child. The prepared expertise, based on an all-Polish survey in 2009, showed that 29% of women consuming alcohol within the previous year were pregnant [28]. The effects of ethyl alcohol on the foetus still remain a poorly described and recognized problem. In children whose mothers consumed alcohol, problems with functioning difficult to diagnose and classify are observed, as well as sensory integration disorders [29,30]. For these children, effective SI therapy is necessary. The objective of my other studies was an analysis of the effect of SI therapy on gross motor functions in children after prenatal exposure to alcohol. Positive effects of SI therapy were observed for the following tests: finger to nose, in the prone extension position, in supine flexed position, ATOS, STOS, the Schilder test, dynamic balance, static balance, gravitational insecurity and trunk stabilisation. The effectiveness of SI therapy was not confirmed only in the tests for cocontraction and muscle tone [31]. In my other studies, the effect of SI on fine motor functions in children after prenatal exposure to alcohol was analysed. The high effectiveness of SI therapy was confirmed in children after prenatal exposure to alcohol based on fine motor functions (p < 0.001). The positive effects of SI therapy were proved for the following tests: quick rotation of the forearms (p < 0.001), eye movements (p < 0.001), fingers to a thumb and in prone extension position (p < 0.001) [31]. The author of the presented report also investigated the effect of SI on the balance system. A high effectiveness of SI therapy was confirmed in children after prenatal exposure to alcohol on the example of the balance system. Positive results of SI therapy were confirmed for the following tests: Schilder (p < 0.001), dynamic balance (p < 0.001), static balance (p < 0.001), gravitational insecurity (p < 0.001) and post-rotatory nystagmus (p < 0.05) [32,33]. Similar results of therapy were obtained in studies on children with developmental deficits from

Kielce (Poland). The study covered a group of 153 children admitted for therapy at the third year of life. The objective of the study was an analysis of the effect of sense integration: mainly proprioception, the vestibular system on vocabulary development. The presented results of the performed diagnoses of the sensor integration processes indicate that the most impaired areas are related with static and dynamic balance, as well as post-rotatory nystagmus. They also indicate considerable disorders in the development of the vestibular system Early diagnosis of sensor integration disorders in the population examined can, to a great extent, prevent more serious problems in the development of speech and language. The greatest deficits in the sphere of vocabulary development were noted in the area of creating subwords, and defining concepts in combination with dynamic and static balance disorders. In addition, a relationship was observed between vestibular system disorders and acquisition of skills with respect to vocabulary [34].

CONCLUSIONS

The high effectiveness of SI therapy was confirmed in children after prenatal exposure to alcohol on the example of muscle tone. Statistically significant results of SI therapy were proved for the following tests: in the prone extension position, in supine flexed position, ATOS, STOS, the Schilder test and trunk stabilisation. Analysis of Cronbach's Alpha positions showed a high reliability of the applied SI tests, especially prior to therapy. The sensor integration technique (SI) exerts a beneficial effect on the normalization of muscle tone in children after prenatal exposure to alcohol.

REFERENCES

- Medina AE, Wozniak JR, Klintsova AY, Hamilton DA. Proceedings of the 2016 annual meeting of the Fetal Alcohol Spectrum Disorders Study Group. Alcohol. 2017, 21, 65:19-24. doi: 10.1016/j.alcohol.2017.07.001.
- Fish EW, Wieczorek LA, Rumple A, Suttie M, Moy SS, Hammond P, Parnell SE. The Enduring Impact of Neurulation Stage Alcohol Exposure: A Combined Behavioral and Structural Neuroimaging Study in Adult Male and Female C57BL/6J Mice. Behav Brain Res. 2017, 28. pii: S0166-4328(17)31040-9. doi: 10.1016/j.bbr.2017.10.020.
- Abbott CW, Rohac DJ, Bottom RT, Patadia S, Huffman KJ. Prenatal Ethanol Exposure and Neocortical Development: A Transgenerational Model of FASD. Cereb Cortex. 2017, 6:1-14. doi: 10.1093/cercor/bhx168.

- Allen S, Casey J. Developmental coordination disorders and sensory processing and integration: Incidence, associations and co-morbidities. Br J Occup Ther. 2017, 80 (9): 49-557. doi: 10.1177/0308022617709183.
- Jarmasz JS, Basalah DA, Chudley AE, Del Bigio MR. Human Brain Abnormalities Associated With Prenatal Alcohol Exposure and Fetal Alcohol Spectrum Disorder. J Neuropathol Exp Neurol. 2017, 1, 76 (9): 813-833. doi: 10.1093/jnen/nlx064.
- 6. Nash A, Davies L. Fetal Alcohol Spectrum Disorders: What Pediatric Providers Need to Know. J Pediatr Health Care. 2017, 31 (5): 594-606. doi: 10.1016/j.pedhc.2017.04.002.
- Lange S, Probst C, Gmel G, Rehm J, Burd L, Popova S. Global Prevalence of Fetal Alcohol Spectrum Disorder Among Children and Youth: A Systematic Review and Metaanalysis. JAMA Pediatr. 2017, 1, 171 (10):948-956. doi: 10.1001/jamapediatrics. 2017. 1919.
- Ganthous G, Rossi NF, Giacheti CM. Oral narrative of individuals with Fetal Alcohol Spectrum Disorder. Codas. 2017, 10, 29 (4): e20170012. doi: 10.1590/2317-1782/201 72017012.
- Tran TD, Amin A, Jones KG, Sheffer EM, Ortega L, Dolman K. The Use of Trace Eyeblink Classical Conditioning to Assess Hippocampal Dysfunction in a Rat Model of Fetal Alcohol Spectrum Disorders. J Vis Exp. 2017, 5: 126. doi: 10.3791/55350.
- Sarman I, Rangmar J. Alkohol under graviditet kan riskera folkhälsan Alkohol under fosterlivet påverkar inte bara barnets hjärna – risk för kardiovaskulära och metabola sjukdomar på sikt. Lakartidningen. 2017, 3: 114.
- Ren Z, Wang X, Yang F, Xu M, Frank JA, Wang H, Wang S, Ke ZJ, Luo J. 35. Ethanolinduced damage to the developing spinal cord: The involvement of CCR2 signaling. Biochim Biophys Acta. 2017, 1, 1863 (11):2746-2761. doi: 10.1016/j.bbadis.2017.07.035.
- Pascual M, Montesinos J, Montagud-Romero S, Forteza J, Rodríguez-Arias M, Miñarro J, Guerri C. TLR4 response mediates ethanol-induced neurodevelopment alterations in a model of fetal alcohol spectrum disorders. J Neuroinflammation. 2017, 24; 14 (1): 145. doi:10.1186/s12974-017-0918-2.
- Fan J, Taylor PA, Jacobson SW, Molteno CD, Gohel S, Biswal BB, Jacobson JL, Meintjes EM. Localized reductions in resting-state functional connectivity in children with prenatal alcohol exposure. Hum Brain Mapp. 2017, 38 (10):5217-5233. doi: 10.1002/ hbm.23726.

- 14. Camargo Moreno M, Mooney SM, Middleton FA. Heterogeneity of p53 dependent genomic responses following ethanol exposure in a developmental mouse model of fetal alcohol spectrum disorder. PLoS One. 2017, 19; 12 (7): e0180873. doi: 10.1371/journal. pone.0180873.
- 15. Denny L, Coles S, Blitz R. Fetal Alcohol Syndrome and Fetal Alcohol Spectrum Disorders. Am Fam Physician. 2017, 15, 96 (8): 515-522.
- McLennan JD, Braunberger P. A Critique of the New Canadian Fetal Alcohol Spectrum Disorder Guideline. J Can Acad Child Adolesc Psychiatry. 2017, 26 (3): 179-183.
- Bolaños AD, Coffman BA, Candelaria-Cook FT, Kodituwakku P, Stephen JM. Altered Neural Oscillations During Multisensory Integration in Adolescents with Fetal Alcohol Spectrum Disorder. Alcohol Clin Exp Res. 2017 Sep 25. doi: 10.1111/acer.13510.
- Seguin D, Gerlai R. Fetal alcohol spectrum disorders: Zebrafish in the analysis of the milder and more prevalent form of the disease. Behav Brain Res. 2017, 5, pii: S0166-4328 (17) 31429-8. doi: 10.1016/j.bbr.2017.10.005.
- Syed YY. AbobotulinumtoxinA: A Review in Pediatric Lower Limb Spasticity. Paediatr Drugs. 2017, 19 (4): 367-373. doi: 10.1007/s40272-017-0242-4.
- 20. Paquet A, Olliac B, Golse B, Vaivre-Douret L. Evaluation of neuromuscular tone phenotypes in children with autism spectrum disorder: An exploratory study. Neurophysiol Clin. 2017, 47 (4): 261-268. doi: 10.1016/j.neucli.2017.07.001.
- Steiner N, Torres A, Reddy A, Augustyn M. Congenital Hypotonia in Toddlerhood. J Dev Behav Pediatr. 2017, 38 (7): 556-557. doi:10.1097/DBP.000000000000497.
- Myrie SB, Pinder MA. Skeletal muscle and fetal alcohol spectrum disorder. Biochem Cell Biol. 2017, 1. doi: 10.1139/bcb-2017-0118.
- Burden MJ, Jacobson SW, Sokol RJ, Jacobson JL. Effects of prenatal alcohol exposure on attention and working memory at 7.5 years of age. Alcoholism, Clinical and Experimental Research 2005, 29, 443–452.
- 24. Abel, EL, Sokol RJ. Fetal alcohol syndrome is now leading cause of mental retardation. Lancet 1986, 2: 1222.
- 25. Knuiman S, Rijk CH, Hoksbergen RA, van Baar AL Children adopted from Poland display a high risk of foetal alcohol spectrum disorders and some may go undiagnosed. Acta Paediatr 2015, 104, (2): 206-11.
- Klecka M, Janas-Kozik M, Krupka-Matuszczyk I. Rozwój diagnostyki poalkoholowego spektrum zaburzeń rozwojowych – przegląd narzędzi diagnostycznych. Psychiatria i Psychologia Kliniczna 2010, 4: 298-302.

- 27. Murawski NJ, Moore EM, Thomas JD, Riley EP. Advances in Diagnosis and Treatment of Fetal Alcohol Spectrum Disorders: From Animal Models to Human Studies. Alcohol Res 2015, 37 (1): 97-108.
- 28. Okulicz-Kozaryn K, Borkowska M, Brzózka K. FASD Prevalence among Schoolchildren in Poland. J Appl Res Intellect Disabil 2015, 1: 10.1111/12219.
- Horecka-Lewitowicz A, Lewitowicz P, Adamczyk-Gruszka O, Skawiński D, Szpringer M. Objawy, przebieg i postępowanie w alkoholowym zespole płodowym. Studia Medyczne 2013, 29, 2: 195–198.
- Horecka-Lewitowicz A, Lewitowicz P, Adamczyk-Gruszka O, Skawiński D, Szpringer M. Alkoholowy zespół płodowy – przyczyny, kryteria diagnostyczne i prewalencja. Studia Medyczne 2014, 30, 1: 48–50.
- Wilczyński J, Zawada K. The impact of sensory integration therapy on gross motor function in children after prenatal exposure to alcohol. Studia Medyczne 2015, 31 (1): 10– 17.
- 32. Wilczyński J. The effectiveness of therapy by Sensory Integration in children with Fetal Alcohol Syndrome on the example of fine motor. Studium Vilense 2015, 11, A: 86-89.
- 33. Wilczyński J, Sowińska A, Wilczyński I. Sensory integration technique and body equilibrium processes in children after prenatal exposure to alcohol. Journal of Education, Health and Sport 2017; 7, 12: 356-367 DOI http://dx.doi.org/10. 5281/ zenodo.1145826 http://ojs.ukw.edu.pl/index.php/johs/article/view/5206.
- 34. Chalik P. Wyniki badania logopedycznego. Biuletyn Pedagogiczny 2013, 3, 4: 38-54.

Variables	Ν	Mean	Median	Minimu Maximu		SD	Skewnes	Kurtosi
				m	m		s	S
Age	20	4.45	4	4	5	0.510	0.218	-2.183
Body height	20	1.0185	1.02	0.94	1.08	0.036	-0.393	-0.228
Body mass	20	16.35	16	14	19	1.348	0.283	-0.771
BMI	20	15.738	15.893	14.704	16.340	0.527	-0.671	-0.666

Table 1. Anthropometric data of children in the study.

Table 2. Muscle tone before and after SI therapy.

Muscle tone prior to SI therapy								
Tests for muscle tone		N	Mean	Median	Minimum	Maximu m	SD	
Test for cocontraction		20	1.000	1.000	1.000	1.000	0.000	
Test in prone extension position		20	0.700	0.700	0.200	1.000	0.178	
Test in supine flexed position		20	0.888	1.000	0.250	1.000	0.190	
Test for ATOS		20	0.350	0.333	0.333	0.500	0.051	
Test for STOS		20	0.700	0.667	0.667	1.000	0.103	
Muscle tone	20		0.850	1.000	0.667	1.000	0.170	
Schilder test	20		0.417	0.333	0.000	0.667	0.183	
Trunk stabilisation	20		0.500	0.667	0.000	1.000	0.296	
Muscle tone - overall result	20		0.676	0.692	0.390	0.896	0.120	
Muscle tone after SI therapy								
Tests for muscle tone	Ν	Mean	Median	Minimum	Maximum	SD	Difference	
Test for cocontraction	20	1.000	1.000	1.000	1.000	0.000	0.000	
Test in prone extension position	20	0.990	1.000	0.800	1.000	0.045	0.290	
Test in flexed supine position	20	0.988	1.000	0.750	1.000	0.056	0.100	
Test for ATOS	20	0.500	0.500	0.500	0.500	0.000	0.150	
Test for STOS	20	1.000	1.000	1.000	1.000	0.000	0.300	
Muscle tone		0.850	1.000	0.667	1.000	0.170	0.000	
Schilder test		0.717	0.667	0.667	1.000	0.122	0.300	
Trunk stabilisation		0.767	0.667	0.333	1.000	0.244	0.267	
Muscle tone - overall result		0.851	0.854	0.715	0.938	0.057	0.176	

Tests for muscle tone _		Initial	examination	Final exan	nination	Difference		Wilcoxon test	
		N	Mean	N	Mean			p	
									•
Test for cocontraction		20	3	20	3		0	-	
Test in prone extension position		20	3.5	20	4.95		1.45	< 0.001	
Test in supine flexed position		20	3.55	20	3.95		0.4	< 0.05	
Test for ATOS		20	2.1	20	3		0.9	< 0.001	
Test for STOS		20	2.1	20	3		0.9	< 0.001	
Muscle tone		20	2.55	20	2.55		0	-	
Schilder test		20	1.25	20	2.15		0.9	< 0.001	
Trunk stabilisation 2		20	1.5	20	2.3		0.8	< 0.001	
Student's <i>t</i> -test for muscle tone prior to and after SI therapy $(n = 20)$									
Mean	SD	Differ	ence	SD Difference	t	df	р	Confidence	Confidence
0.851	0.057							-95.00%	95.00%
0.676	0.120	0.1	76	0.075	10.444	19	0.001	0.141	0.211

Table 3. Wilcoxon matched pairs test for muscle tone.