Korabiusz Katarzyna, Fabian Danielewska Anna, Wawryków Agata, Stecko Monika, Torbé Andrzej. Possibilities of physiotherapy in lipodystrophy. Journal of Education, Health and Sport. 2018;8(9):1771-1777. eISNN 2391-8306. DOI http://dx.doi.org/10.5281/zenodo.1410088 http://ojs.ukw.edu.pl/index.php/johs/article/view/5920

https://pbn.nauka.gov.pl/sedno-webapp/works/876212

The journal has had 7 points in Ministry of Science and Higher Education parametric evaluation. Part b item 1223 (26/01/2017).

1223 Journal of Education, Health and Sport eissn 2391-8306 7

© The Authors 2018;

This article is published with open access at Licensee Open Journal Systems of Kazimierz Wielki University in Bydgoszcz, Poland

Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author (s) and source are credited. This is an open access article licensed under the terms of the Creative Commons Attribution Non commercial license Share alike.

(http://creativecommons.org/licenses/by-nc-sa/4.0/) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.

The authors declare that there is no conflict of interests regarding the publication of this paper.

Received: 02.08.2018. Revised: 18.08.2018. Accepted: 20.09.2018.

Possibilities of physiotherapy in lipodystrophy

mgr Katarzyna Korabiusz¹, lek. Anna Fabian- Danielewska², mgrAgataWawryków¹, mgr Monika Stecko¹,

prof.drhab.n.med.Andrzej Torbé³

¹Pomeranian Medical University in Szczecin, Doctoral Studies of the Faculty of Health Sciences, Żołnierska 54 st., 71-210 Szczecin

²Pomeranian Medical University in Szczecin, PhD Studies at the Faculty of Medicine, Żołnierska 54 st., 71-210 Szczecin

³Pomeranian Medical University in Szczecin, Obstetrics and Gynecology Clinic,

Powstańców Wlkp. 72 st., 70-111 Szczecin

Dane autora korespondencyjnego: Katarzyna Korabiusz email: korabiuszk@gmail.com

Abstract

Lipodystrophy is one of the most common and the most undesirable female dermatological conditions. Both slim and overweight people can struggle with it. It is possible to fight cellulite with various physiotherapeutic techniques and methods.

Research goal: Goal of this dissertation is a review of literature about possibilities of physiotherapy in lipodystrophy.

Conclusion: The appropriate physiotherapeutic methods are necessary to improve the flexibility, appearance and microcirculation in areas affected by lipodystrophy.

Key words: lipodystrophy, physiotherapy

Introduction

Possibilities of physiotherapy in lipodystrophy

The female type of lipodystrophy is characterized by unsightly bumps, depressions and bulges, which appear most often around the thighs, hips, buttocks, arms and abdomen. The skin in area of lipodystrophy is dry, often also rough because of trophic disorders. Lipodystrophy is often associated with the occurrence of stretch marks. Cellulite can occur at any age, most often affects mature women, but more and more often begins to appear in adolescence. It affects about 90% of women, previously combined with its overweight and lack of physical activity, while research now shows that it also applies to athletic slim women [1,2]. In cellulite - in contrast to cellulitis - there is no inflammation, only changes in the subcutaneous tissue [3-5].

There are two types of lipodystrophy: water (hard) and fatty (soft). It depends on the ground, on which it is formed. In the form of water cellulite, it is very difficult to distinguish it from strongly developed muscle tissue - it happens in women who did not have weight fluctuations, and thus their skin did not lose its elasticity [6]. Cases of water type of cellulite are often found in sportsmen, dancers and teenagers. Soft cellulite, on the other hand, usually occurs in inactive women because it is associated with poor tension and muscle flaccidity. Often, it also occurs in women who were overweight and quickly returned to the optimal weight. It is also found in people with genetically conditioned flaccid skin. Its appearance may also be caused by the use of dehydrating medications. It differs from water cellulite in that it is loose, does not stick tightly to the muscles, it usually takes up a large body surface. Changes in body position usually change the shape of the skin surface. Fatty cellulite gives the impression that it is between the skin and muscles, reminiscent of a quilted coverlet. In this form of cellulite there are often changes in the microcirculation. It should be emphasized that soft cellulite often develops from a poorly treated hard form [7].

Among the forms of lipodysystrophy we can also distinguish edema cellulite (this is the heaviest of the form of cellulite), mixed cellulite (combining soft and hard cellulite), and pseudo-lipodystrophy in which there is a limp skin in the buttocks, thighs and arms [8.9].

Physiotherapeutic treatments used in lipodystrophy

It should be emphasized that just as the reasons for the occurrence of cellulite are varied, so to combat it should be approached individually. Certainly, lipodystrophy treatment should be multi-path and based on a balanced diet, physical activity, wearing appropriate clothing (loose and airy), walking in properly selected shoes, avoiding hot baths, and avoiding long-term stress. One should also exclude coexisting diseases - cardiovascular, gynecological, nephrological, gastrointestinal, metabolic and hormonal disorders. The appropriate physiotherapeutic methods are also necessary to improve the flexibility, appearance and microcirculation in areas affected by lipodystrophy [10,11].

Lymphatic drainage

It should be used every day, using special techniques, it moves the lymph from the areas of retention. All techniques used in drainage must be pushing, they should not warm up, so they should be smooth and gentle. It is important that the upper limb is massaged in the drainage position, i.e. facilitating the outflow of the lymph. For this purpose, the limb is

placed in the facade with special wedges. Selected classical massage techniques are used, such as stroking, rubbing, kneading, and compressions [12].

Chinese cupping massage

It seems to be a very effective method. It is certainly one of the methods with the fastest effects. Before the intervention should be applied a specially composed olive, which contains active ingredients. Then create underpressure using the Chinese bulb and move the longitudinal or circular movements towards the lymph nodes. A gentle massage improves microcirculation, accelerates metabolism and lipolysis. As a result, cellulite is reduced, the skin is smoothed, the figure is modeled and its color is improved. The process itself should last 15 to 30 minutes, repeated every 2-3 days. The series of treatments should be from 10 to 20, then prophylactically one Chinese cupping massage should be performed every week [13-16].

You can also use a vibration massage (specially constructed cups filled with water) improving circulation and reducing tissue tension, or manual segmental massage. Sequential pneumatic massage allows the pressure of cuffs filled with air to drain the lymph and reduce the formation of edema. During this massage, toxins and excess fluids from the retention sites are discharged in the right direction, and the tropics improve [17].

Electrostimulation

The electrostimulation of the gluteal muscles and the back of the thighs brings tangible results in the reduction of lipodystrophy. It involves stimulating the muscles to spasms by impulses generated by the apparatus for muscle stimulation. Using two pairs of electrodes placed in the place of cellulite, we force the muscles to work, thereby reducing fat tissue and reducing the circumferences.

Ultrasounds

The basis of this therapy is the phenomenon of electro-evaporation, i.e. the opening of path in the cell membrane. This is done due to the use of an electric field and transport of active gel components inside the cells by means of impulses and phonophoresis.

Acoustic wave

Radio waves are electromagnetic waves generated by the high frequency electric current. The warming of the tissues causes various physiological and thermoregulation-related reactions. It

is a controlled tissues warming, and thus safe. The operation of this treatment allows the removal of excess fat tissue, and thus the visible reduction of cellulite [18, 19].

Kinesiotape

It involves sticking spots exposed to swelling and places with circulatory disorder with special patches. Kinesiotape facilitate drainage of lymph, reduce edema, improve trophic and accelerate regeneration. Slices are practically imperceptible, because they have thickness and weight similar to human skin, they are made of cotton, do not contain glue, are resistant to water and breathable, so the risk of skin allergy is minimal [20].

Pearl bath

It is based on a bath in water, to which air with a pressure of 2 to 4 atmospheres is introduced using special nozzles. The bath takes place in a specially adapted bathtub, which is equipped with numerous holes, through which air escapes, creating small bubbles in the water. The bath can be used spring water, mineral water or with the addition of essential oils. Water should have a temperature between 35-37 ° C. Bubbles produced during bathing have a positive effect on the condition of the skin, cardiovascular system and the human nervous system.

Scottish whips

The treatment takes place in a bath, it can be used on selected areas or whole body. It involves pouring the body with water at a pressure of 40 degrees Celsius in a series of 2-minute alternately with 30-second watering with water at a temperature of 20 degrees Celsius or lower. This treatment regenerates tissues and improves tropics.

Thalassotherapy

In these treatments are used: seawater, marine algae and healing mud from the seabed with the addition of plant extracts. These preparations are heated, because the skin in this way easily and quickly absorbs valuable mineral salts, oligoelements and vitamins. The treatment slims, relaxes and stimulates the natural renewal of the body's forces.

Paraffin treatments

Paraffin is an unsaturated hydrocarbon obtained in the distillation of crude oil. It has a large heat capacity, i.e. it absorbs large amounts of heat and gives them very slowly. Treatments with its use reduce fat tissue and cellulite.

In the prevention and treatment of lipodystrophy, graded pressure tights are also used to improve the functioning of lower limb muscles and improve circulation [21].

Summary

Literature review suggests that the problem of lipodystrophy is very common and concerns not only middle-aged and overweight women, but also young women with BMI within normal limits. It is important to choose appropriate physiotherapeutic methods in the prevention and treatment of this phenomenon. These treatments should be combined with a well-balanced diet and physical activity.

References:

1. Zegarska B. Cellulit (II). Nazewnictwo, definicja, związek występowania z wiekiem, płcią i budową ciała, Dermatol Estet 2011, 13/1: 29-32.

2. Haneke E. Cellulite – fakty i mity, Dermatol Estet 2006, 3(44): 132-138.

3. Błaszczyk-Kostanecka M, Wolska H (red.). Dermatologia w praktyce. PZWL-Wydawnictwo Lekarskie, Warszawa 2006.

4. Pugliese PT. The pathogenesis of cellulite: a new concept. J Cosmet Dermatol 2007, 6(2): 140-142.

5. Wollina U, Gildman A, Berger U. Esthetic and cos

6. [Pavicic T, Borelli C, Korting HC. Cllulite-the greatest skin problem in healthy women. J Dtsch Dermatol Ges 2008, (10): 861-870.

7. Quatresooz P, Xhaufl aire-Uhoda E, Piérard-Franchimont C et al. Cellulite histopathology and related mechanobiology. Int J Cosmet Sci 2006, 28(3): 207-210.

8. Ignaciuk A. Cellulit – przyczyny, terapie, możliwości. Beauty Forum 2003, 9: 32-33.

9. Rossi R, Andre Vergnanini L. Cellulite. Dermatol Estet 2001, 6 : 259-267 10. Avram MM. Cellulite. A review of its physiology and treatment. J Cosmet Laser Ther 2004, 6 (4): 181-185.

10. Smalls LK, Hicks M, Passeretti D et al. Eff ect of weight loss on cellulite: gynoid lypodystrophy. Reconstr Surg 2006, 118 (2): 510-516.

11. Smalls LK, Lee CY, Whitestone J et al. Quantitative model of cellulite: threedimensional skin surface topography, biophysical characterization, and relationship to human perception. J Cosmet Sci 2005, 56 (2): 105-120.

12. Fey H., Petsitis X.: Słownik kosmetyczny. Medpharm, Wrocław 2011, 40–41.

13. Pelc K., Smaga S., Zielińska K., Ocena skuteczności antycellulitowego masażu bańką chińską, Wyższa Szkoła Informatyki i Zarządzania w Rzeszowie.

14. Ortonne J-P, Queille-Roussel C, Duteil L. Treatment of cellulite: Effectiveness and sustained effect at 6 months with Endermologie demonstrated by several quantitative evaluation methods. Nouv Dermatol 2003, 22: 261-269.

15. Rona C., Carrera M., Berardesca E.: Testing anticellulite products. "International Journal of Cosmetic Science", 2006; 28(3), 169-173.

16. Cao H., Liu J., Lewith G.T.: Traditional Chinese Medicine for treatment of fibromyalgia: a systematic review of randomized controlled trials. "The Journal of Alternative and Complementary Medicine", 2010; 16(4), 397-409.

17. Kasprzak WP, Mańkowska A. Fizykoterapia, medycyna uzdrowiskowa i SPA. PZWL, Warszawa 2008.

18. de Godoy J.M., de Godoy M.F.: Treatment of cellulite based on the hypothesis of a novel physiopathology. "Clin. Cosmet. Investig Dermatol.", 2011;4:55-9

19. Wanitphakdeedecha R., Manuskiatti W.: Treatment of cellulite with a bipolar radiofrequency, infrared heat, and pulsatile suction device: a pilot study. "Journal of Cosmetic Dermatology", 2006; 5(4):284-288.

20. Kase K., Wallis J., Kase T., Clinical therapeuthic applications of Kinesio Taping Method, Japan 2003.

21. Ponikowska I., Ferson D.,: Nowoczesna medycyna uzdrowiskowa, Medi Press, Warszawa 2009.