

Fiałek Bartosz, Zukow Walery. Myocardial infraction – the history of therapy prior to reperfusion. Journal of Education, Health and Sport. 2018;8(10):57-61. eISSN 2391-8306. DOI <http://dx.doi.org/10.5281/zenodo.1447149> <http://ojs.ukw.edu.pl/index.php/johs/article/view/6157>

The journal has had 7 points in Ministry of Science and Higher Education parametric evaluation. Part b item 1223 (26/01/2017).
1223 Journal of Education, Health and Sport eISSN 2391-8306 7

© The Authors 2018;

This article is published with open access at Licensee Open Journal Systems of Kazimierz Wielki University in Bydgoszcz, Poland
Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author (s) and source are credited. This is an open access article licensed under the terms of the Creative Commons Attribution Non commercial license Share alike.
(<http://creativecommons.org/licenses/by-nc-sa/4.0/>) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.

The authors declare that there is no conflict of interests regarding the publication of this paper.

Received: 02.09.2018. Revised: 12.09.2018. Accepted: 05.10.2018.

Myocardial infraction – the history of therapy prior to reperfusion

Bartosz Fiałek¹, Walery Zukow²

1) Angiology and Internal Clinic, Dr. J. Biziel University Hospital no. 2 in Bydgoszcz, Bydgoszcz, Poland

2) Nicolaus Copernicus University, Toruń, Poland

Abstract

The description of stenocardial ailments was presented for the first time in 1772. Until the 1950s, the patients suffering from myocardial infraction were treated with the wait-and-see attitude – they were confined to bed for 6 weeks, were not allowed to eat or to take up everyday activities on their own. It was as late as in the 1960 it was discovered that being immobilised for too long may be harmful and lead to the increase of the clotting/embolism complications. The first issue of Harrison's Principles of Internal Medicine was released in

1950 which advised the oxygen therapy, as well as pharmacotherapy (nitro-glycerine, anticoagulants) used in the severe phase of myocardial infarction. One of the breaking moments on the path of the development of care considering the patient with myocardial infarction was the introduction of CCU (Desmond Julian, 1961).

Key words: myocardial infarction, stent,

In 1772 Wilim Heberden presented the description of stenocardial ailments, naming them *angina pectoris*. The term is present in the modern cardiology – in fact, its creation preceded the examination of pathogenesis and of the histopathological image of the described pathology [12]. Specialists were able to determine the thrombosis of coronary artery as the cause of death as early as the first half of the 19th century. It was proven by various examinations carried out on animals whose coronary arteries were purposely ligated, as well as posthumous observations on individual deceased people. In 1901, Krehl was the first person to notice that thrombosis of coronary artery does not have to necessarily lead to sudden death caused by heart issues, although it is a frequent cause of creation of heart aneurysm, scars or heart walls [13]. Hektoen proved in the 1990s that myocardial infarction is caused by a clot of the coronary artery, which is secondary to atherosclerotic changes [14]. Then, in 1910, Obrastzov and Straschesko presented a series of five patients who, before their death, showed physical features of myocardial infarction which also proved to be the cause of their death later on [15].

In 1912, James Herrick published his thesis, which had an enormous impact on the way of conducting with myocardial infraction until the 1950s. In his thesis, Herrick drew a conclusion that the main element of conducting with the above-mentioned patient should be them being confined to their bed for six weeks. Such patient should not be permitted to move, shave or eat on their own. An important part of the therapy was diet, which consisted of 1200 kcal – mostly fluid meals with low levels of salt and animal fats. It needs to be emphasised that only a small portion of the patients managed to recover to normal, despite continuing the sitting mode after being released from the hospital [16]. Herrick has had an even bigger impact on modern cardiology by introducing and popularising the electrocardiography examination (EKG) as a diagnostic instrument after a severe myocardial infraction, which was invented by Einthoven in 1902 [13]. In 1923, Wearn described the first series of clinical cases of patients with myocardial infraction. They were advised of absolutely no physical effort and limiting fluids in order to prevent a possible lung oedema, the glycosides of digitalis were introduced for the patients suffering from stasis in the pulmonary circulation stasis caffeine and camphor were also used in treating hypotension, fainting and atrioventricular block. The nitrates were not advised as they lowered the arterial pressure. In 1928, Parkinson and Bedford introduced their experiments using morphine used in order to relieve the pain of patients suffering from myocardial infraction [13]. In 1940, French and Dock showed the description of important changes in atherosclerosis of young patients, who had died in a result of wounds and injuries without any visible symptoms of myocardial infraction [17]. The pioneer of cardiological prophylaxis is Paul Dudley who introduced his theory in 1944, which lead to the creation of Framingham Heart Study in 1948. Two year later, the first issue of Harrison's Principles of Internal Medicine is released which advises, in the case of severe myocardial infraction, oxygen therapy, nitro-glycerine administered under the tongue or anticoagulants (such as heparin or warfarin) used in order to prevent from subsequent myocardial infarction and clotting/embolism complications. This conducting actually reduces the mortality in this particular group of patients [19].

The idea that longer immobilisation of the patient might be harmful, as it resulted in considerable increase of the clotting/embolism complications, began becoming more popular in the 1960s. Because of that in cases of myocardial infractions that are not complicated, the period, in which the patient is confined to bed, was shortened to five days and the overall time

of hospitalisations to four weeks. Additionally, the patient was gradually prepared to live normally before being released from the hospital [20].

The next phase of the development of care considering the patient with myocardial infarction was introducing the coronary care units (CCU) which reduced the mortality in the first hours following the infarction from 30% (noticed in the last decade) to 15%. This phase began in 1961, when Desmond Julian released the idea of CCU in the Royal Infirmary in Edinburgh, and when Hugh Day introduced it to the common practice of the Bethany Medical Centre. Julian analysed the course of the cardiac arrest of five patients, four of which had died. The author thought the cause of therapeutic failure was the late attempt of CPU, as well as little experience connected with infarctions within the team. CCU functioned according to four main rules: 1) the patients suffering from myocardial infarction were supposed to be constantly monitored with the EKG equipment, featuring an alert that would set in case of any serious arrhythmias, 2) patients stayed in a specially assigned room with experienced and trained staff, as well as most modern equipment and medications, 3) the staff had to be ready for an immediate start of the CPR procedure, 4) the possibility of fast CPR and external defibrillation which, in case there were not any doctors around, were the roles of special nurses. Within five years, CCUs were introduced in the majority of hospitals in developed countries [21].

References

- [1] Zipes DP, Libby P, Bonow R. Braunwald's Heart Disease: a textbook of cardiovascular medicine 7th EDITION. Dunfermline: Elsevier Saunders; 2004.
- [2] Kardiologia, red. T. Mandecki – Warszawa 2000 p. 221
- [3] Szczeklik A. Interna Szczeklika 2014. Kraków: Medycyna Praktyczna; 2014.
- [4] Kardiologia red. T. Mandecki – Warszawa 2000 p. 224-227
- [5] Thygesen K, Alpert JS, White HD. Uniwersalna definicja zawału serca. Kardiol Pol 2008;66:47–62. doi:10.1017/CBO9781107415324.004.
- [6] Rywik S., Broda G., Piotrowski W. i wsp. Epidemiologia chorób układu krążenia- Program Pol-MONICA. Kard Pol T XLIV, sup II:II-7.
- [7] Wong ND. Epidemiological studies of CHD and the evolution of preventive

- cardiology. *Nat Rev Cardiol* 2014;11:276–89.
- [8] Bandosz P, Drygas W, Rutkowski M, Koziarek J, Wyrzykowski B, Bennett K, et al. Decline in mortality from coronary heart disease in Poland after socioeconomic transformation : modelling 2012;8136:1–10.
- [9] Berg J, Björck L LG i wsp. Continuing decrease in coronary heart disease mortality in Sweden. *BMC Cardiovasc Disord* 2014;14:9.
- [10] Vartiainen E, Laatikainen T, Peltonen M i wsp. Thirty-five-year trends in cardiovascular risk factors in Finland. *Int J Epidemiol* 2010;39:504–18..
- [11] Preis SR, Pencina MJ Hwang S-J i wsp. Trends in cardiovascular disease risk factors in individuals with and without diabetes mellitus in the Framingham Heart Study. *Circulation* 2009;120:212–20
- [12] Silverman ME. William Heberden and Some Account of a Disorder of the Breast.. *Clin Cardiol.* 1987 10(3):211-213.
- [13] Braunwald E. Evolution of the management of acute myocardial infarction: a 20th century saga. *Lancet* 1998;352:1771–4.
- [14] Hektoen L. Embolism of the left coronary artery; sudden death. *Med. Newsl (Lond).* 1892; 61: 210–210.
- [15] Obrastzov WP, Straschesko ND. Zur Kenntnis der Thrombose der Koronararterien des Herzens. *Z Klin Med.* 1910; 71: 116–32..
- [16] Lown B. *The Lost art of healing*: Houghton Mifflin Company (1st ed) 1996;332.
- [17] French AJ. Fatal coronary arteriosclerosis in young soldiers. *JAMA* 1944;124:1233–1237.
- [18] Muller JE. Diagnosis of myocardial infarction: historical notes from the Soviet Union and the United States. *Am J Cardiol* 1977; 40: 269–71.
- [19] Harrison TR, Beeson PB, Thorn GW, et al. *Principles of internal medicine*. New York: Blackiston: 1950.
- [20] Braunwald E. The treatment of acute myocardial infarction: the Past, the Present, and the Future. *Eur Hear J Acute Cardiovasc Care* 2012;1:9–12. doi:10.1177/2048872612438026.
- [21] Julian DG. Treatment of cardiac arrest in acute myocardial ischemia and infarction. *Lancet* 1961:840–44.