Polyovyy V., Khorshani Bilel, Sydorchuk R., Kyfiak P., Khomko O., Plehutsa I. The clinically relevant diabetic foot syndrome morphology. Journal of Education, Health and Sport. 2021;11(05): 262-274. eISSN 2391-8306. DOI http://dx.doi.org/10.12775/JEHS.2021.11.05.027 https://apcz.umk.pl/czasopisma/index.php/JEHS/article/view/JEHS.2021.11.05.027 https://zenodo.org/record/5129576

The journal has had 5 points in Ministry of Science and Higher Education parametric evaluation. § 8. 2) and § 12. 1. 2) 22.02.2019. © The Authors 2021; This article is published with open access at Licensee Open Journal Systems of Nicolaus Copernicus University in Torun, Poland Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author (s) and source are credited. This is an open access article licensed under the terms of the Creative Commons Attribution Non commercial license Share alike. (http://creativecommons.org/license/s/bu-cs.4/4.0/) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited. The authors declare that there is no conflict of interests regarding the publication of this paper.

Received: 15.05.2021. Revised: 21.05.2021. Accepted: 28.05.2021.

THE CLINICALLY RELEVANT DIABETIC FOOT SYNDROME MORPHOLOGY

V. P. Polyovyy, Bilel Khorshani, R. I. Sydorchuk, P. V. Kyfiak, O. Y. Khomko, I. M. Plehutsa

Bukovinian State Medical University, Chernivtsi, Ukraine

Abstract

Objective: to evaluate the morphological changes in diabetic foot syndrome emphasizing the ultrastructural cellular changes.

Material and methods. Biopsies of the chronic lesions in 120 diabetic foot patients following necessary preparatory stages were studied by histological, histochemical and electron microscopy methods.

Results. The inflammatory infiltration and destruction and involving all layers of skin and soft tissue, the phenomena of vacuolation, apoptosis or karyolysis; tissue edema, diffuse infiltration by neutrophils, vascular plethora with stasis, thrombosis, and necrosis were observed. Microcirculatory tract thrombosis exacerbated tissue ischemia, contributing to the progression of pathology. The macrophages' cytoplasm characterized by the presence of phagocytosed and digested microorganisms often intact inside them. The changes in the status of nuclear and perinuclear material were accompanied by compression and dehydration of the cell, which ended in its fragmentation and the formation of tightly contacting bodies of various shapes

Conclusions. Morphological changes in diabetic foot syndrome are diverse and uneven, occurring not only on tissue and cellular level but involves subcellular and molecular structures, encouraging further research in this field. Foci of destruction and lysis of the newly

formed epithelium are characteristic for diabetic foot, which is not observed in the normal wound process.

Keywords: diabetes mellitus type 2; diabetic foot; morphological and microscopic findings; electron microscopy.

Background

The number of diabetes mellitus (DM) patients worldwide is growing sustainably, making it one of the biggest contemporary healthcare challenges worldwide, with the global prevalence of diabetic foot syndrome (DFS) varying from 3% in Oceania to 13% in North America, with a global average of about 6.4%. The annual incidence of DFS complicating DM is known to be about 2-5% with the lifetime risk ranging from 15-20%. It is worth mentioning that peripheral arterial disease (PAD) is an important risk factor associated with a twentyfold higher prevalence in diabetic patients with the probability of amputation within one year after the first ulcer or gangrene over 34% and the mortality rate over 5% [1]. Approximately 20-33% of all costs related to DM are expenditures related to treatment of DFS [2], with the cost of medical care for DFS increasing. The steady growth in the number of diabetic patients has become an important issue in the decision of the health and welfare budget worldwide. In addition, the increase in complications due to diabetes is a burden, not only for patients, but also for the national healthcare systems and economies. Furthermore, DFS determines huge portion of public health loses making it essential for public health policy makers to advocate for implementation of prevention and treatment recommendations [3].

Pathogenesis of DFS involves both systemic and local factors. Major influence on metabolic disorders, endothelial dysfunction and atherosclerosis is thought to have strong genetic background with multiple genetic polymorphisms involved [4, 5]. Local factors are generally more likely linked to microbiota, biofilm formation and impaired local resistance not to mention neurological and vascular factors, etc [6, 7].

Treatment of diabetic foot is complex and often unsuccessful [8]. Most recent IWGDF Guidelines indicate that "Multiple disciplines are involved in the management of diabetic foot disease, and interdisciplinary treatment is the cornerstone of its management and prevention" [9]. Moreover, common combination of systemic effects of diabetes and comorbidities like hypertension and target-organ damage add complexity to this task [10, 11]. While the term "chronic diabetic wound" to some extend corresponds to diabetic foot syndrome, there are several pathogenetic mechanisms involved in the condition making it distinguishable from

other chronic wounds and skin defects. Moreover, surprisingly few papers are dedicated to morphological studies of the DFS, stimulating current research interest.

The objective of the study was to evaluate the morphological changes in diabetic foot syndrome emphasizing the ultrastructural cellular changes.

Material and methods

The comprehensive pathomorphological study of biopsy material of 120 cases of surgical treatment of DFS was performed. Biopsies of the chronic skin wounds in DFS patients were taken intraoperatively or during non-operative procedures (debridement, necrectomies, tissue grafting, etc). The study was conducted in full accordance with the principles of the Council of Europe Convention on Human Rights and Biomedicine, Declaration of Helsinki on the ethical principles for medical research involving human subjects, and other acting international and national legislations in bioethics (including GCP, EU directives, etc). All patients signed an informed consent prior to participating in the study. The study protocol was fully approved by the institutional ethics committee. All patients underwent treatment conducted in accordance with the state-of-the-art guidelines and recommendations [8].

Bioptates sampling, preparation and staining

The bioptates material underwent standard care and passed all necessary preparatory stages (Fixation, Embedding, Sectioning, Staining, and Mounting). Samples were fixed for 48 hours in a neutral buffered 10% formalin solution, then dehydrated in an ascending battery of alcohols and embedded into paraffin medium. Serial histological sections 5 µm thick were produced on a sled microtome. Several approaches were combined for optical analysis: review histological methods (hematoxylin and eosin staining, van Gieson picric acid and acid fuchsia staining); histochemical methods (Gömöri trichrome stain, Slinchenko stain, and Weighert staining for detection of connective tissue components, fibrin; PAS-reaction; toluidine blue for visualization of tissue basophils and large granular lymphocytes).

Electron microscopy

For electron microscopy, bioptates samples were fixed in 2.5% phosphate buffered glutaraldehyde solution (pH 7.2-7.4) and fixed in 1% osmium tetroxide solution. The material was dehydrated in alcohols of increasing concentration and embedded in Araldite (HAM, USA). Morphological structures were contrasted in the process of dehydration of the material with a saturated solution of uranyl acetate, and in sections with lead citrate. Sections with a thickness of 40-60 nm were analyzed in the TESLA BS-500 electron microscope.

Results

Destruction and inflammatory infiltration involving all layers of skin and soft tissues of the marginal area of skin wounds were observed during morphological examination (Fig. 1-2).



Figure 1. Wound skin of the lower shin. Pyoderma. The stratum corneum is unevenly basophilic, stratified. A large number of intraepidermal and subepidermal blister defects filled with purulent exudate with a small amount of fibrin. Hematoxylin and eosin staining, ×250.



Figure 2. The skin of the foot. Pyoderma. The epidermis is in a state of severe hydropic dystrophy, cellular edema in the form of translucent perinuclear spaces. Hematoxylin and eosin staining, $\times 250$.

The epidermis (absent in ulcers) was characterized by a decrease in thickness towards the edge of the wound, the presence of a large number of cells with the phenomena of vacuolation, apoptosis or karyolysis; tissue edema, diffuse infiltration by neutrophils, vascular plethora with stasis and thrombosis, necrosis (Fig. 3).



Figure 3. Soft tissues of the shin. Skeletal muscle tissue with a predominance of fibers in a state of necrosis (fibers are swollen, unstructured, with the loss of striation), fibrous connective tissue with partial necrosis. Hematoxylin and eosin staining, ×250.

The lumen of dilated blood vessels was filled with clearly contoured blood elements. Hyalinosis of the microcirculatory tract vascular wall was present in all tissues studied. In the presence of phlegmon or abscess, the vascular wall was infiltrated with leukocytes and subjected to lysis, which led to small hemorrhages (Fig. 4).

The microcirculatory tract thrombosis exacerbated tissue ischemia, contributing to the progression of pathologic processes. Destruction of collagen fibers (fibrinoid edema and necrosis) with infiltration of the surrounding area by neutrophils and macrophages was observed in the wound edges. Against the background of necrotic changes, elements of granulation tissue in the form of short thin strands of proliferating spindle-shaped fibroblasts accompanied by capillaries were revealed; at some patients such phenomenon was absent. At a distance of 0.1-0.3 mm from the edges of the wound, we observed the regeneration of the epidermis in the form of small thin layers of newly formed keratinocytes.

Masses of chaotically located high electron density fibrillary structures, occasionally were found in the cytoplasm of macrophages. In microfibrillar intracellular masses, tufts of more or less scattered fibrils with fuzzy contours were observed. The fibrous fibrillary structures or their fragments were partially dissolved in the cytosol mass, especially near the nucleus (Fig. 5).



Figure 4. The chronic wound of the foot. Pyoderma. Intravascular leukocytosis, acute purulent vasculitis. Diffuse leukocyte infiltration in perivascular tissues. Hematoxylin and eosin staining, ×250.



Figure 5. Area of macrophage with the presence of minor intussusceptional formations of the nuclear membrane and perinuclear location of fibrous structures. Electron microscopic image, $\times 30000$.

The excrescences located on the cell surface were represented by large fields of peripheral areas of the cytoplasm. The granular endoplasmic reticulum was moderately developed and represented by intracellular channels or cisterns. The cytoplasm contained a small number of membrane-bound ribosomes. There was mainly peripheral localization of the granular endoplasmic reticulum and vacuolar formations. The latter are represented by clusters of rounded or irregularly shaped vacuoles. In addition, the endoplasmic reticulum had the appearance of oblong cisterns and accumulations of rounded or elongated vesicles (Fig. 6).



Figure 6. Fine-grained vacuolation of macrophage cytoplasm. Electron microscopic image, ×35000.

A small number of pinocytic edged vesicles, as well as homogeneous small electrondense granules in the cytoplasm were observed in some macrophages. The cytoplasmic matrix had a moderate electron density, on outside of the cell, insignificant digit-like processes or pseudopodia were observed. In some vesicles there was a fine electron density of fine-grained material. Larger vacuoles contained membrane fragments and myelin figures. The extracellular material included preserved and cleaved collagen fibrils, granular matter, electron-dense homogeneous formations (Fig. 7). The cytoplasm between the zones of the lamellar complex was occupied by small mitochondria, a small number of polysomal rosettes, tanks of granular endoplasmic reticulum and various sizes of phagolysosomes with the presence of phagocytosed and digested microorganisms (Fig. 8). Phagocytosed microorganisms were often found intact inside macrophages (Fig. 9).



Figure 7. Extracellular lysis of tissue structures. Electron microscopic image, ×40000.



Figure 8. Completed phagocytosis. There are residual bodies of microorganisms and tissue structures. Electron microscopic image, ×30000.

Uneven cell hypertrophy was often observed, and the nucleus, as a rule, acquired an intricate shape, and it possible to see accumulations of chromatin on the periphery of the nuclei. In addition to the described changes, a constant feature was the appearance of large vacuoles, diverse in structure. Often they were near the nucleus, were filled with crumbly or light homogeneous content (Fig. 10).



Figure 9. Incomplete phagocytosis. Individual microorganisms that are in a state of division. Electron microscopic image, ×45000.



Figure 10. Formation of a large electron-enlightened vacuole near the nucleus. Electron microscopic image, ×35000.

Vacuoles could contain other, often high-density structures (Fig. 11). The described changes in the state of nuclear and perinuclear material were accompanied by compression and dehydration of the cell, which ended in its fragmentation and the formation of tightly contacting bodies of various shapes (Fig. 12).



Figure 11. Aggregation of phagolysosomes of high electron density in the central regions of the macrophage. Electron microscopic image, ×25000.



Figure 12. Residual bodies and lysosomal structures with the presence of dense granular material and vacuolar formations. Electron microscopic image, ×20,000.

Discussion

Diabetic foot syndrome generally presents with diverse and often unclear morphological picture, ranging from minor skin defect to vast necrotic gangrenous and inflammatory changes. Thus, finding something more specific or clinically relevant is an uneasy task. However, it is clear that in the diabetic foot, a comprehensive morphological assessment of all ulcers and defects is required to devise a proper treatment and revascularization strategy [12]. Obviously, DFS presents with a multitude of wounds that are heterogeneous in both morphology and topography. Even one single patient may present with several wounds dispersed over more than one angiosome or manifest a large ulcer that lies on the verge of two angiosomes with different phases of wound processes and different infestation status. Furthermore, different systemic changes and peculiarities of pathogenesis may significantly variate adding to diversity in morphological picture.

Morphological studies of the DFS are usually focused on clinically relevant changes like vascular changes and deformities (e.g. Charcot's foot) formation [13] with much less attention payed to microscopic changes in skin and soft tissues. Even lesser consideration is concentrated on the intracellular changes as they are assumed to be sufficiently studied by biochemical and physiological methods [14].

Current study complements additional information to the existing database on DFS focusing on intracellular changes obtained through electronic microscopy. Such information may be useful for determining future local therapies for DFS as new types of treatment approaches are undergoing rapid development, like molecular or cellular surgery [15].

Conclusions

Morphological changes in diabetic foot syndrome are diverse and uneven. This study confirms that pathologic changes in diabetic foot syndrome occur not only on tissue and cellular level but involves subcellular and molecular structures, encouraging further research in this field. Microcirculatory disorders are expressed in the form of venular plethora, the phenomena of stasis in capillaries, hemolysis of erythrocytes and marginal standing of formed elements in venules and capillaries. The vascular network occupied $11.7\pm1.0\%$, but the vascular loops were unevenly spaced and markedly different in size. The appearance of foci of destruction and lysis of the newly formed epithelium are characteristic for DFS, which is not observed in the normal wound process.

Conflict of Interest

The authors declare no conflict of interest.

References

1. Zhang P, Lu J, Jing Y, Tang S, Zhu D, Bi Y. Global epidemiology of diabetic foot ulceration: a systematic review and meta-analysis. *Ann Med.* 2017;49(2):106-16.

2. Behrendt CA, Sigvant B, Szeberin Z, Beiles B, Eldrup N, Thomson IA, Venermo M, Altreuther M, Menyhei G, Nordanstig J, Clarke M, Rieß HC, Björck M, Debus ES. International Variations in Amputation Practice: A VASCUNET Report. *Eur J Vasc Endovasc Surg.* 2018;56(3):391-99.

272

3. Chun DI, Kim S, Kim J, Yang HJ, Kim JH, Cho JH, Yi Y, Kim WJ, Won SH. Epidemiology and Burden of Diabetic Foot Ulcer and Peripheral Arterial Disease in Korea. *J Clin Med.* 2019;8(5):748. doi:10.3390/jcm8050748.

4. Sydorchuk L, Dzhuryak V, Sydorchuk A, Levytska S, Petrynych V, Knut R, Kshanovska A, Iftoda O, Tkachuk O, Kyfiak P, Popovich, Khomko O, Sydorchuk R. The cytochrome 11B2 aldosterone synthase gene rs1799998 single nucleotide polymorphism determines elevated aldosterone, higher blood pressure, and reduced glomerular filtration, especially in diabetic female patients. *Endocrine Regulations*. 2020;54(3):217-226. doi:10.2478/enr-2020-0024.

5. Dzhuryak V, Sydorchuk L, Sydorchuk A, Kamyshnyi O, Kshanovska A, Levytska S, Knut R, Sheremet M, Ivashchuk S, Petrynych O, Kazantseva T, Nikyfor L, Melnychuk L, Sokolenko A, Yarynych Yu, Semianiv M, Repchuk Yu, Voroniuk K, Sydorchuk R, Sokolenko L, Iftoda O, Kushnir O. The cytochrome 11B2 aldosterone synthase gene CYP11B2 (RS1799998) polymorphism associates with chronic kidney disease in hypertensive patients. *Biointerface Res Applied Chem.* 2020;10(3):5406-5411. https://doi.org/10.33263/BRIAC103.406411.

6. Pavlovich KV, Sydorchuk RI. [Injecting material for skin regeneration Lacerta application in treatment of trophic ulcers in patients with diabetic foot syndrome]. *Klin Khir*. 2015;(7):44-6. Ukrainian.

7. Sydorchuk RI, Sydorchuk LP, Khorshani Bilel, Hrushko OI, Plehutsa OM, Sydorchuk I, Plehutsa IM, Vakarchuk AV, Sydorchuk AR, Ilaschuk II, Plehutsa N. Microbiota alleviates host resistance and causes septic complications in diabetic foot syndrome. *Infection*. 2019;47(Suppl. 1):S57-S58.

8. Schaper et al. Practical guidelines on the prevention and management of diabetic foot disease (IWGDF 2019 update). *Diab Metab Res Rev.* 2020. e3266.

9. Van Netten et al. Definitions and criteria for diabetes foot disease. *Diab Metab Res Rev.* 2020. e3268.

10. Sydorchuk LP, Dzhuryak VS, Sydorchuk AR, Levytska SA, Knut RP, Sokolenko MO, Iftoda OM, Kushnir OV, Popovych AI, Sydorchuk RI. Association of lipids' metabolism disorders with aldosterone synthase CYP11B2 (-344C/T) gene polymorphism in hypertensive patients depending on glomerular filtration rate. *PharmacologyOnLine*. 2020;2:230-242.

11. Sydorchuk LP, Sokolenko AA, Sydorchuk AR, Kryklyvets LG, Biryuk IG, Fliundra IG, Sokolenko MA. Insulin resistance in patients with arterial hypertension and

abdominal obesity depending on ACE and PPAR-γ2 genes polymorphism: A new opinion concerning an old problem. *New Armenian Med J.* 2015;9:43-51.

12. Aerden D, Denecker N, Gallala S, Debing E, Van den Brande P. Wound morphology and topography in the diabetic foot: hurdles in implementing angiosome-guided revascularization. *Int J Vasc Med.* 2014;2014:672897. doi: 10.1155/2014/672897.

13. Gurmeet S. Morphological Changes in Diabetic Foot. *Int J Pulm Res Sci.* 2019;410): 1-5. DOI: 10.19080/IJOPRS.2019.04.555637.

14. Huang Y, Kyriakides TR. The role of extracellular matrix in the pathophysiology of diabetic wounds. *Matrix Biol Plus*. 2020;6-7:100037. https://doi.org/10.1016/j.mbplus.2020.100037.

15. Brem H, Tomic-Canic M. Cellular and molecular basis of wound healing in diabetes. *J Clin Invest*. 2007;117(5):1219-22. doi: 10.1172/JCI32169.