

Figure 1. ** — for proofs wherein no line is used as a hypothesis of a rule of inference more than once; a is the inverse Ackermann function

cannot p-simulate truth tables and the reverse simulation is impossible too. Another important result is that tree-like Gentzen systems without cut cannot p-simulate dag-like Gentzen systems without cut.

It is also important to mention the paper by Finger [6] wherein it was proved that dag-like Gentzen systems without cut but with substitution rule linearly simulate tree-like Gentzen systems with cut.

Another important result is mentioned by Cook and Nguyen in [3] and attributed to Krajíček [10]. He shows that tree-like Gentzen systems with cut p-simulate dag-like Gentzen systems with cut.

2. A simulation of natural deduction

In this section we prove some theorems that provide speedups of one natural deduction system over another.

THEOREM 1. $\Gamma \mid \frac{nd_{\mathcal{F}}}{n} A \Rightarrow \Gamma \mid \frac{d_{\mathcal{F}}}{O(n)} A$

PROOF. The proof is straightforward, since every instance of axiom schema and every hypothesis in $nd_{\mathcal{F}}$ -derivation remain w.l.o.g. an axiom and hypothesis respectively in $d_{\mathcal{F}}$ -derivation, every instance of mp_n and dr_n becomes an instance of mp_g and dr_g respectively. \square

THEOREM 2. $\Gamma \mid \frac{d\mathcal{F}}{n} D \Rightarrow \Gamma \mid \frac{nd\mathcal{F}}{O(n^2)} D$

PROOF. We prove this theorem with the method used by Buss and Bonnet in [1] in their proof of Theorem 4.

Recall that every line in a $d\mathcal{F}$ -derivation is a sequent of the form $\Gamma \vDash D$. Suppose we have a $d\mathcal{F}$ -proof π_g of a sequent $\Gamma \vDash D$ of length n . This means that the $d\mathcal{F}$ -derivation is actually the following sequence:

$$\Gamma_1 \vDash A_1, \dots, \Gamma_n \vDash A_n = \Gamma \vDash D$$

Our simulation goes as follows. We first substitute each sequent $\Gamma_i \vDash A_i$ ($1 \leq i \leq n$) for the formula $\bigwedge \Gamma_i \supset A_i$ with $\bigwedge \Gamma_i$ denoting the conjunction of all formulas in Γ_i ordered and associated arbitrarily. This gives us the following sequence π' :

$$\bigwedge \Gamma_1 \supset A_1, \dots, \bigwedge \Gamma_n \supset A_n \tag{1}$$

It is important to note that since $d\mathcal{F}$ is complete, all formulas in (1) are tautologies. However, π' is not a valid $nd\mathcal{F}$ -proof, which means that we have to show by induction on n that we can fill in every gap in no more than $O(n)$ steps. The proof splits into four cases depending on how $\Gamma_n \vDash A_n$ was inferred.

CASE 2.1. $\Gamma_n \vDash A_n$ is an assumption or hypothesis.

In this case $\Gamma_n \vDash A_n$ has the form $A_n \vDash A_n$ and becomes $A_n \supset A_n$ in π' . One can prove $A_n \supset A_n$ in $nd\mathcal{F}$ in a constant number of steps.

CASE 2.2. $\Gamma_n \vDash A_n$ is an instance of an axiom schema.

$\Gamma_n \vDash A_n$ has the form $\vDash A_n$ and becomes A_n in an $nd\mathcal{F}$ -derivation with A_n being an axiom schema. Since $d\mathcal{F}$ and $nd\mathcal{F}$ use the same axiom schemata, we prove A_n in $nd\mathcal{F}$ in one step by simply writing it down.

CASE 2.3. $\Gamma_n \vDash A_n$ is inferred by mp_g .

This means that $\Gamma_n \vDash A_n$ has the form $\Gamma_n \vDash B$ and there are two such sequents, namely $\Gamma_{n_1} \vDash A$ and $\Gamma_{n_2} \vDash A \supset B$, that $n_1, n_2 < n$ and $\Gamma_{n_1} \cup \Gamma_{n_2} = \Gamma_n$. These sequents become $\bigwedge \Gamma_n \supset B$, $\bigwedge \Gamma_{n_1} \supset A$, and $\bigwedge \Gamma_{n_2} \supset (A \supset B)$ in π' respectively. We fill in the gap as follows.



$$\begin{array}{l}
\vdots \\
\wedge \Gamma_{n_1} \supset A \\
\vdots \\
\wedge \Gamma_{n_2} \supset (A \supset B) \\
\left[\begin{array}{l}
\wedge \Gamma_n - \text{assumption} \\
\vdots (*) \\
\wedge \Gamma_{n_1} \\
A - mp_n \\
\vdots (*) \\
\wedge \Gamma_{n_2} \\
A \supset B - mp_n \\
B - mp_n
\end{array} \right. \\
\wedge \Gamma_n \supset B - dr_n
\end{array}$$

Let Γ_n contain m formulas, $\Gamma_{n_1} - m_1$ formulas, and $\Gamma_{n_2} - m_2$ formulas. It is clear that neither of these numbers is greater than n . One can show by induction on m that both $\wedge \Gamma_{n_1}$ and $\wedge \Gamma_{n_2}$ can be inferred from $\wedge \Gamma_n$ in $O(m)$ steps.

CASE 2.4. $\Gamma_n \vDash A_n$ is inferred by dr_g .

$\Gamma_n \vDash A_n$ has the form $\Gamma_n \vDash A \supset B$ which becomes $\wedge \Gamma_n \supset (A \supset B)$ in π' . Moreover, we have $\Gamma_{n_1} \vDash B$ in π_g such that $n_1 < n$ and $\Gamma_n = \Gamma_{n_1} \setminus A$. $\Gamma_{n_1} \vDash B$ becomes $\wedge \Gamma_{n_1} \supset B$. We have two cases depending on whether A is in Γ_{n_1} .

CASE 2.4.1. $A \notin \Gamma_{n_1}$.

In this case $\Gamma_n = \Gamma_{n_1}$. We proceed as follows.

$$\begin{array}{l}
\vdots \\
\wedge \Gamma_{n_1} \supset B \\
\left[\begin{array}{l}
\wedge \Gamma_{n_1} - \text{assumption} \\
B - mp_n \\
\left[\begin{array}{l}
A - \text{assumption} \\
A \supset B - dr_n
\end{array} \right. \\
\wedge \Gamma_n \supset (A \supset B) - dr_n
\end{array} \right.
\end{array}$$

CASE 2.4.2. $A \in \Gamma_{n_1}$

The simulation goes as follows.

$$\begin{array}{l}
 \vdots \\
 \wedge \Gamma_{n_1} \supset B \\
 \left[\begin{array}{l}
 \wedge \Gamma_n - \text{assumption} \\
 \left[\begin{array}{l}
 A - \text{assumption} \\
 \vdots (*) \\
 \wedge \Gamma_{n_1} \\
 B - mp_n
 \end{array} \right. \\
 A \supset B - dr_n
 \end{array} \right. \\
 \wedge \Gamma_n \supset (A \supset B) - dr_n \text{ because } \Gamma_n = \Gamma_{n_1}
 \end{array}$$

Let Γ_n contain m formulas and $\Gamma_{n_1} - m_1$ formulas. It is clear that neither of these numbers is greater than n . One can show by induction on m that $\wedge \Gamma_{n_1}$ can be inferred from $\wedge \Gamma_n$ and A in $O(m)$ steps. \square

2.1. Fitch-style systems

THEOREM 3. $\Gamma \mid_n^{nd\mathcal{F}} D \Rightarrow \Gamma \mid_{O(n)}^{\mathbf{F}} D$

PROOF. We prove this theorem by induction on n . Our induction hypothesis is as follows. *For any $m < n$ \mathbf{F} linearly simulates $nd\mathcal{F}$ so that all formulas occurring in an $nd\mathcal{F}$ derivation also occur in the \mathbf{F} derivation.*

The proof splits into four cases depending on how D was derived.

CASE 3.1. D is either an assumption or a member of Γ .

In this case D becomes an assumption or a member of Γ in the \mathbf{F} -derivation respectively.

CASE 3.2. D is an instance of an axiom schema.

In this case D can be proved in \mathbf{F} in a constant number of steps.

CASE 3.3. D is derived by mp_n .

In this case $D = B$ and an $nd\mathcal{F}$ -derivation has one of the two following forms:

$$\left[\begin{array}{l}
 \vdots \\
 A \supset B \\
 \vdots \\
 A \\
 B - mp_n
 \end{array} \right. \quad \left[\begin{array}{l}
 \vdots \\
 A \supset B \\
 \vdots \\
 A \\
 B - mp_n
 \end{array} \right.$$



We proceed respectively as follows.

$$\begin{array}{c}
 \vdots \\
 A \supset B \quad \text{IH} \\
 \vdots \\
 A \quad \text{IH} \\
 \hline
 B \quad \supset\text{E}
 \end{array}
 \qquad
 \begin{array}{c}
 \vdots \\
 A \supset B \quad \text{IH} \\
 \hline
 \begin{array}{c}
 \vdots \\
 A \quad \text{IH} \\
 A \supset B \quad \text{R} \\
 \hline
 B \quad \supset\text{E}
 \end{array}
 \end{array}$$

In both cases we derive A and $A \supset B$ by induction hypothesis and B by $\supset\text{E}$ in a constant number of steps. In the right-hand case, however, we have to reiterate one of the hypotheses of $\supset\text{E}$ which takes us exactly one step.

CASE 3.4. D was derived by dr_n .

The proof of this case is straightforward since in both $nd\mathcal{F}$ and \mathbf{F} when dr_n (respectively $\supset\text{I}$) is applied, the last open assumption has to be closed. Hence, we simply substitute an instance of dr_n with an instance of $\supset\text{I}$. \square

THEOREM 4. $\Gamma \stackrel{\mathbf{F}}{\vdash}_n D \Rightarrow \Gamma \stackrel{nd\mathcal{F}}{\vdash}_{O(n)} D$.

PROOF. We prove the theorem by induction on n . The proof splits into cases, depending on how D was derived.

If D is a hypothesis or an assumption in \mathbf{F} -derivation, it becomes a hypothesis (resp. assumption) in an $nd\mathcal{F}$ -derivation.

If D was reiterated, this simply means that it had already been derived earlier. Hence we obtain it by the induction hypothesis.

Finally, D could be derived by one of the rules of inference. The only interesting case here would be rule $\vee\text{E}$ since we substitute all instances of $\supset\text{I}$ with dr_n and in all other cases we simply derive conclusion of a rule from its premises. This can be done in a constant number of steps because both $nd\mathcal{F}$ and \mathbf{F} are sound and complete. If D was derived by $\vee\text{E}$, \mathbf{F} -derivation has the following form.



$$\begin{array}{c}
 \vdots \\
 \hline
 A \vee B \\
 \hline
 \begin{array}{c}
 | \quad A \\
 \hline
 \vdots \\
 D \\
 | \quad B \\
 \hline
 \vdots \\
 D
 \end{array} \\
 \hline
 D \qquad \vee E
 \end{array}$$

We proceed as follows.

$$\begin{array}{c}
 \vdots \\
 A \vee B \\
 \left[\begin{array}{l}
 A - \text{assumption} \\
 \vdots \\
 D - \text{by IH} \\
 A \supset D - dr_n
 \end{array} \right. \\
 \left[\begin{array}{l}
 B - \text{assumption} \\
 \vdots \\
 D - \text{by IH} \\
 B \supset D - dr_n
 \end{array} \right. \\
 \vdots \\
 D - \text{in a constant number of steps}
 \end{array}$$

As one can see, we have derived D in a constant number of steps. \square

2.2. Gentzen's natural deduction ND

THEOREM 5. $\Gamma \mid \frac{d\mathcal{F}}{n} D \Rightarrow \Gamma \mid \frac{ND}{O(n)} D$.

PROOF. \ulcorner The proof is straightforward since each step in $d\mathcal{F}$ -derivation is either an axiom which has an ND-proof of constant length, an assumption or member of Γ which remains an assumption or member of Γ in ND-derivation, or was derived by either mp_g or dr_g . We substitute

an instance of mp_g for $\text{FB} - \frac{A \supset B}{B} -$ and dr_g for $\text{FE} - \frac{[A]}{B} \frac{A \supset B}{A \supset B}$ ⁴. We can do this since we aren't required to close the last open assumption in both $d\mathcal{F}$ and ND. \square

THEOREM 6. $\Gamma \mid \frac{\text{ND}}{n} D \Rightarrow \Gamma \mid \frac{d\mathcal{F}}{O(n)} D.$

PROOF. Recall that we assumed that ND has dag-like proofs. This means that we can, following Reckhow [13] designate each step of the derivation of C from Γ as $\Delta \vDash A$ with Δ being the set of all open assumptions and formulas from Γ .

We need this convention because $d\mathcal{F}$ proofs are sequences of sequents (cf. [1, P.691]), not trees of them so our simulation results won't be affected by the necessity of deriving one formula multiple times from same assumptions.

We prove the theorem by induction on n .

CASE 6.1 (Base case, $n = 1$).

If $n = 1$, D is either an assumption or a member of Γ . In both cases we have a $d\mathcal{F}$ -derivation of exactly one line, namely, $D \vDash D$.

CASE 6.2 (Induction step).

We suppose that for all $m < n$ if all occurrences of falsum are substituted for the fixed contradictory formula $-p_1 \wedge \neg p_1$ $d\mathcal{F}$ simulates ND linearly so that every step occurring in the ND derivation, except for occurrences of falsum is present in the $d\mathcal{F}$ derivation. Occurrences of \wedge are substituted with $p_1 \wedge \neg p_1$. We now need to show this for n . We have different cases depending on how D was derived.

The cases of all rules, except for

$$\text{OB: } \frac{\frac{[A] \quad [B]}{A \vee B} \quad C}{C}$$

and

$$\text{NE: } \frac{[A]}{\neg A}$$

are straightforward and will be omitted here.

CASE 6.2.1. D was derived by OB.

⁴ $[A]$ means that assumption A is discharged.



In this case $D = C$ and the ND-derivation has the following form.

$$\text{OB: } \frac{\begin{array}{c} \vdots \\ A \vee B \\ \vdots \end{array} \quad \frac{\begin{array}{c} [A] \\ \vdots \\ C \end{array} \quad \begin{array}{c} [B] \\ \vdots \\ C \end{array}}{C}}$$

Since ND derivations are dag-like, we can rewrite this derivation and get the following one.

$$\begin{array}{c} \vdots \\ A \vDash A \\ \vdots \\ \Gamma_1, A \vDash C \\ B \vDash B \\ \vdots \\ \Gamma_2, B \vDash C \\ \vdots \\ \Gamma_3 \vDash A \vee B \\ \vdots \\ \Gamma_1 \cup \Gamma_2 \cup \Gamma_3 \vDash C \quad \text{OB} \\ \\ \Gamma_1 \cup \Gamma_2 \cup \Gamma_3 = \Gamma \end{array}$$

Our simulation goes as follows.

$$\begin{array}{c} A \vDash A \\ \vdots \\ \Gamma_1, A \vDash C \quad \text{by IH} \\ B \vDash B \\ \vdots \\ \Gamma_2, B \vDash C \quad \text{by IH} \\ \vdots \\ \Gamma_3 \vDash A \vee B \quad \text{by IH} \\ \Gamma_1 \vDash A \supset C \quad dr_g \\ \Gamma_2 \vDash B \supset C \quad dr_g \\ \vdots \\ \Gamma_1 \cup \Gamma_2 \cup \Gamma_3 \vDash C \quad \text{in a constant number of steps} \end{array}$$

CASE 6.2.2. D was derived by NE.

In this case $D = \neg A$ and the ND-derivation has the following form.

$$\begin{array}{c} [A] \\ \vdots \\ \text{NE: } \frac{\wedge}{\neg A} \end{array}$$

Our simulation goes as follows.

$$\begin{array}{ll} A & \vDash A \\ & \vdots \\ \Gamma, A & \vDash p_1 \wedge \neg p_1 & \text{by IH} \\ & \vdots \\ & \vDash (p_1 \wedge \neg p_1) \supset p_1 & \text{in a constant number of steps} \\ & \vDash (p_1 \wedge \neg p_1) \supset \neg p_1 & \text{in a constant number of steps} \\ \Gamma, A & \vDash p_1 & mp_g \\ \Gamma, A & \vDash \neg p_1 & mp_g \\ \Gamma & \vDash A \supset p_1 & dr_n \\ \Gamma & \vDash A \supset \neg p_1 & dr_n \\ & \vdots \\ & \vDash (A \supset p_1) \supset ((A \supset \neg p_1) \supset \neg A) & \text{in a constant number of steps} \\ \Gamma & \vDash (A \supset \neg p_1) \supset \neg A & mp_g \\ \Gamma & \vDash \neg A & mp_g \end{array}$$

In both cases we have derived D in a constant number of steps. \square

3. A simulation of Gentzen-style systems

In this section we will prove theorems considering pairwise simulation of $d\mathcal{F}$, PKT and PKT*. We start with a theorem about the simulation of PKT by $d\mathcal{F}$.

Once again, recall that we assume our PKT and PKT* proofs to be dag-like, not tree-like. This lets us use one sequent many times without need to derive it multiple times. Note, also, that the additive version of the cut rule is used.

LEMMA 1. Assume, we have a sequent $A_1, \dots, A_n \rightarrow C$. Then it takes no more than $O(n^2)$ steps of PKT proof to remove all repeated formulas from the antecedent.

PROOF. This statement can be easily proved since we need no more than $2n$ permutations and then one contraction for any repeated formula to be removed. \square

LEMMA 2. Assume, we have a sequent $A_1, \dots, A_n \rightarrow C$. Then we need no more than $O(m \cdot n)$ steps of PKT proof to augment its antecedent with m formulas in any order.

PROOF. The proof is straightforward since we only need to add m formulas and put them on their places which won't take more than n permutations for each formula. \square

THEOREM 7. $\Gamma \mid \frac{d\mathcal{F}}{n} D \Rightarrow \Gamma \mid \frac{\text{PKT}}{O(n^3)} D$.

PROOF. We will now prove the theorem by induction on n . The simulation goes as follows: every sequent $\Gamma_i \vDash A_i$ from $d\mathcal{F}$ -derivation becomes a sequent $\Gamma_i \rightarrow A_i$ with formulas in the antecedent being in an arbitrary order. We need to show that the gaps can be filled in in $O(n^2)$ steps.

An assumption or a member of Γ , i.e., the line of the form $A \vDash A$ becomes $A \rightarrow A$ which is an initial sequent. An axiom $\vDash A$ becomes $\rightarrow A$ which has a constant-length PKT-proof.

Sequents constituting $mp_g \left(\frac{\Gamma_1 \vDash A \supset B \quad \Gamma_2 \vDash A}{\Gamma_1 \cup \Gamma_2 \vDash B} \right)$ rule of inference become $\Gamma_1 \rightarrow A \supset B$, $\Gamma_2 \rightarrow A$ and $\Gamma_1 \cup \Gamma_2 \rightarrow B$ in PKT derivation. We show that third sequent can be derived in PKT from first and second ones in $O(n^2)$ steps.

$$\begin{array}{c}
 \vdots \\
 W_l \frac{\Gamma_2 \rightarrow A}{\Gamma_2 \rightarrow A, B} \\
 E_l \frac{\Gamma_2 \rightarrow A, B}{\Gamma_2 \rightarrow B, A} \quad W_l \frac{B \rightarrow B}{\Gamma_2, B \rightarrow B} \\
 \vdots \\
 O(n^2) W_l \text{ and } E_l \frac{\Gamma_1 \rightarrow A \supset B}{\Gamma_1, \Gamma_2 \rightarrow A \supset B} \quad W_l \text{ and } E_l \frac{A \supset B, \Gamma_2 \rightarrow B}{A \supset B, \Gamma_1, \Gamma_2 \rightarrow B} \\
 \text{Cut} \frac{\Gamma_1, \Gamma_2 \rightarrow A \supset B \quad A \supset B, \Gamma_1, \Gamma_2 \rightarrow B}{\Gamma_1, \Gamma_2 \rightarrow B} \\
 O(n^2) E_l \text{ and } C_l \frac{\Gamma_1, \Gamma_2 \rightarrow B}{\Gamma \rightarrow B}
 \end{array}$$

We use lemmas 1 and 2 to derive $\Gamma \rightarrow B$ from $\Gamma_1, \Gamma_2 \rightarrow B$ and $\Gamma_1, \Gamma_2 \rightarrow A \supset B$ from $\Gamma_1 \rightarrow A \supset B$ respectively. It is also clear that we need $O(n)$ steps to derive $\Gamma_2, B \rightarrow B$ from $B \rightarrow B$ and $A \supset B, \Gamma_1, \Gamma_2 \rightarrow B$ from $A \supset B, \Gamma_2 \rightarrow B$.

Finally consider the case of $dr_g \left(\frac{\Gamma \vDash B}{\Gamma \setminus \{A\} \vDash A \supset B} \right)$. The sequents become $\Gamma \rightarrow B$ and $\Gamma \setminus \{A\} \rightarrow A \supset B$ in the PKT derivation. We have two cases: $A \in \Gamma$ (left) and $A \notin \Gamma$ (right).



$$\begin{array}{c} \vdots \\ \leq n E_l \frac{\Gamma \rightarrow B}{A, \Gamma \setminus \{A\} \rightarrow B} \\ \supset_r \frac{\Gamma \setminus \{A\} \rightarrow A \supset B}{} \end{array} \quad \begin{array}{c} \vdots \\ W_l \frac{\Gamma \rightarrow B}{A, \Gamma \rightarrow B} \\ \supset_r \frac{\Gamma \rightarrow A \supset B}{} \end{array}$$

□

THEOREM 8. $\Gamma \mid_{\frac{d\mathcal{F}}{n}} D \Rightarrow \Gamma \mid_{\frac{\text{PKT}^*}{O(n)}} D$.

The proof is straightforward since we can see from the proof of Theorem 7 that if we don't count steps made by weakening, permutation and contraction rules (which in the case of PKT* we don't), then we can fill every gap in a constant number of steps. □

We will finally prove that $d\mathcal{F}$ linearly simulates PKT*. Since it is clear that PKT* linearly simulates PKT, it will also entail the linear simulation of PKT by $d\mathcal{F}$.

THEOREM 9. *If $\Gamma \mid_{\frac{\text{PKT}^*}{n}} D$, then there is such a subset $\Gamma' \subseteq \Gamma$ that $\Gamma' \mid_{\frac{d\mathcal{F}}{O(n)}} D$.*

We prove this theorem the same way as Buss and Bonet proved Theorem 11 in [1]. We need the following lemma.

LEMMA 3. *If there is a PKT*-proof π_{PKT^*} of $\Gamma \rightarrow \Delta$ of length n , then there is such a subset $\Xi \subseteq (\Gamma \cup \neg\Delta)$ ($\neg\Delta$ denotes that every formula in Δ is negated), that there exists a $d\mathcal{F}$ -proof $\pi_{d\mathcal{F}}$ of a sequent $\Xi \vdash p_1 \wedge \neg p_1$ of length $O(n)$.*

CASE 3.1. *Base case.* $n = 1$. If the length of π_{PKT^*} is 1, then it consists only of an initial sequent, say, $A \rightarrow A$. It takes a constant number of steps to prove $A, \neg A \vdash p_1 \wedge \neg p_1$ in $d\mathcal{F}$.

CASE 3.2. *Induction step.* We assume that for all $m < n$ there exists such a constant c that $\mid_{\frac{d\mathcal{F}}{c \cdot m}} \Xi \vdash p_1 \wedge \neg p_1$ and prove the lemma for n . The proof splits depending on how the last line of π_{PKT^*} was inferred. We will prove the most representative cases.

CASE 3.2.1. \neg_r . The end of π_{PKT^*} has the following form.

$$\begin{array}{c} \vdots \\ \neg_r \frac{A, \Gamma \rightarrow \Delta}{\Gamma \rightarrow \Delta, \neg A} \end{array}$$

By the induction hypothesis there exist a $d\mathcal{F}$ -proof $\pi'_{d\mathcal{F}}$ of $\Xi' \vdash p_1 \wedge \neg p_1$ ($\Xi' \subseteq (\{A\} \cup \Gamma \cup \neg\Delta)$) of length $c \cdot m$. We need to construct a

$d_{\mathcal{F}}$ -proof $\pi_{d_{\mathcal{F}}}$ of $\Xi \vDash p_1 \wedge \neg p_1$ ($\Xi \subseteq (\{\neg\neg A\} \cup \Gamma \cup \neg\Delta)$) of length $O(n)$.

$$\begin{array}{lll}
\neg\neg A & \vDash & \neg\neg A & \text{assumption} \\
\Xi & \vDash & \Xi & \text{assumptions} \\
& & \vdots & \\
\neg\neg A & \vDash & A & \text{in a constant number of steps} \\
& & \vdots & \\
\Xi & \vDash & p_1 \wedge \neg p_1 & \text{from } \pi'_{d_{\mathcal{F}}}
\end{array}$$

CASE 3.2.2. \wedge_r . The end of π_{PKT^*} has the following form.

$$\wedge_r \frac{\begin{array}{c} \vdots \\ \Gamma \rightarrow \Delta, A \end{array} \quad \begin{array}{c} \vdots \\ \Gamma \rightarrow \Delta, B \end{array}}{\Gamma \rightarrow \Delta, A \wedge B}$$

By the induction hypothesis there exists a proof $\pi'_{d_{\mathcal{F}}}$ of length $c \cdot m$ containing both $\Xi_1 \vDash p_1 \wedge \neg p_1$ ($\Xi_1 \subseteq (\{\neg A\} \cup \Gamma \cup \neg\Delta)$) and $\Xi_2 \vDash p_1 \wedge \neg p_1$ ($\Xi_2 \subseteq (\{\neg B\} \cup \Gamma \cup \neg\Delta)$). We construct the $d_{\mathcal{F}}$ -proof $\pi_{d_{\mathcal{F}}}$ of $\Xi \vDash p_1 \wedge \neg p_1$ ($\Xi \subseteq (\{\neg(A \wedge B)\} \cup \Gamma \cup \neg\Delta)$) as follows.

$$\begin{array}{lll}
\neg(A \wedge B) & \vDash & \neg(A \wedge B) & \text{assumption} \\
\Xi & \vDash & \Xi & \text{assumptions} \\
& & \vdots & \\
\neg(A \wedge B) & \vDash & \neg A \vee \neg B & \text{in a constant number of steps} \\
& & \vdots & \\
\Xi_1 & \vDash & p_1 \wedge \neg p_1 & \text{from } \pi'_{d_{\mathcal{F}}} \\
\Xi_2 & \vDash & p_1 \wedge \neg p_1 & \text{from } \pi'_{d_{\mathcal{F}}} \\
\Xi_1 \setminus \{A\} & \vDash & \neg A \supset (p_1 \wedge \neg p_1) & dr_g \\
\Xi_2 \setminus \{B\} & \vDash & \neg B \supset (p_1 \wedge \neg p_1) & dr_g \\
& & \vdots & \\
\Xi \setminus \{\neg(A \wedge B)\} & \vDash & (\neg A \vee \neg B) \supset (p_1 \wedge \neg p_1) & \text{in a constant number of steps} \\
\Xi & \vDash & (p_1 \wedge \neg p_1) & mp_g
\end{array}$$

CASE 3.2.3. *Cut*. The end of π_{PKT^*} has the following form:

$$\text{Cut} \frac{\begin{array}{c} \vdots \\ \Gamma \rightarrow \Delta, A \end{array} \quad \begin{array}{c} \vdots \\ A, \Gamma \rightarrow \Delta \end{array}}{\Gamma \rightarrow \Delta}$$

By the induction hypothesis there exists a $d\mathcal{F}$ -proof $\pi'_{d\mathcal{F}}$ of length $c \cdot m$ containing both $\Xi_1 \vDash p_1 \wedge \neg p_1$ ($\Xi_1 \subseteq (\Gamma \cup \neg\Delta \cup \{\neg A\})$) and $\Xi_2 \vDash p_1 \wedge \neg p_1$ ($\Xi_2 \subseteq (\Gamma \cup \neg\Delta \cup \{A\})$). We construct the proof of $\Xi \vDash p_1 \wedge \neg p_1$ ($\Xi \subseteq (\Gamma \cup \neg\Delta)$) as follows.

$\neg A$	\vDash	$\neg A$	assumption
A	\vDash	A	assumption
$\Xi \setminus \{A, \neg A\}$	\vDash	$\Xi \setminus \{A, \neg A\}$	assumptions
	\vdots		
Ξ_1	\vDash	$p_1 \wedge \neg p_1$	from $\pi'_{d\mathcal{F}}$
Ξ_2	\vDash	$p_1 \wedge \neg p_1$	from $\pi'_{d\mathcal{F}}$
Ξ	\vDash	$A \supset (p_1 \wedge \neg p_1)$	dr_g
Ξ	\vDash	$\neg A \supset (p_1 \wedge \neg p_1)$	dr_g
	\vdots		
	\vDash	$A \vee \neg A$	in a constant number of steps
	\vdots		
Ξ	\vDash	$p_1 \wedge \neg p_1$	in a constant number of steps

The result follows taking c to be not less than any constant number of steps in any case. \dashv

Theorem 9 follows by the application of Lemma 13 from [1] and Theorem 1 to Lemma 3. \square

4. Concluding remarks

The results from this paper allow us to include \mathbf{F} in the scheme depicted on Figure 1. Combining our results with those from [1], we get the scheme depicted in Figure 2.

It is worth mentioning that the quadratic speedup of $d\mathcal{F}$ over $nd\mathcal{F}$ is due to the deduction rule that allowed for the exclusion of arbitrary assumption instead of the last open one. It turns out that the necessity of reiterating formulas does not lead to a non-linear speedup of one calculus over another since we do not need to infer a formula once more.

Acknowledgements. The author would like to thank his supervisor Vasily Shangin and Professor Dmitry V. Zaitsev for their much-appreciated help and advice during the work on this paper.




Figure 2. ** — for proofs wherein no line is used a hypothesis of a rule of inference more than once; a is inverse Ackermann function

References

- [1] Bonet, M.L., and S.R. Buss, “The deduction rule and linear and near-linear proof simulations”, *The Journal of Symbolic Logic* 58, 2 (1993): 688–709. DOI: [10.2307/2275228](https://doi.org/10.2307/2275228)
- [2] Buss, S.R. (ed.), *Handbook of Proof Theory*, volume 137 of *Studies in Logic and Foundations of Mathematic*, Elsevier Science, Amsterdam, 1st edition, 1998.
- [3] Cook, S.A., and P. Nguyen, *Logical Foundations of Proof Complexity*, Cambridge University Press, Cambridge, 2010. DOI: [10.1017/CBO9780511676277](https://doi.org/10.1017/CBO9780511676277)
- [4] Cook, S.A., and R.A. Reckhow, “The relative efficiency of propositional proof systems”, *The Journal of Symbolic Logic* 44, 1 (1979): 36–50. DOI: [10.2307/2273702](https://doi.org/10.2307/2273702)
- [5] D’Agostino, M., *Investigations into the Complexity of Some Propositional Calculi*, Oxford University Computing Laboratory, Oxford, 1990.
- [6] Finger, M., “Dag sequent proofs with a substitution rule”, pages 671–686 in *We Will Show Them: Essays in Honor of Dov Gabbay’s 60th Birthday*, London, Kings College Publications, 2005.
- [7] Fitch, F.B., *Symbolic Logic: An Introduction*, The Ronald Press Company, N.Y., 1952.

- [8] Gentzen, G., “Untersuchungen über das logische Schließen I”, *Mathematische Zeitschrift* 39 (1935): 176–210. DOI: [10.1007/BF01201353](https://doi.org/10.1007/BF01201353)
- [9] Jaśkowski, S., “On the rules of supposition in formal logic”, *Studia Logica* 1 (1934).
- [10] Krajčiček, J., “On the number of steps in proofs”, *Annals of Pure and Applied Logics* 41 (1989): 153–178.
- [11] Papadimitriou, C.H., *Computational Complexity*, Addison-Wesley Publishing Company, Inc., NY, 1995.
- [12] Pelletier, F.J., “A brief history of natural deduction”, *History and Philosophy of Logic* 20, 1 (1999): 1–31. DOI: [10.1080/014453499298165](https://doi.org/10.1080/014453499298165)
- [13] Reckhow, R. A., “On the lengths of proofs in the propositional calculus”, PhD thesis, University of Toronto, 1976.
- [14] Smullyan, R. M., *First-Order Logic*, Dover Publications, Inc., N.Y., 1995.
- [15] Tseitin, G. S., *On the Complexity of Derivation in Propositional Calculus*, pages 466–483, Springer-Verlag, NY, 1983.
- [16] Urquhart, A., “The complexity of gentzen systems for propositional logic”, *Theoretical Computer Science* 66, 1 (1989): 87–97. DOI: [10.1016/0304-3975\(89\)90147-3](https://doi.org/10.1016/0304-3975(89)90147-3)
- [17] Urquhart, A., “The relative complexity of resolution and cut-free Gentzen systems”, *Annals of Mathematics and Artificial Intelligence* 6, 1–3 (1992): 157–168. DOI: [10.1007/BF01531026](https://doi.org/10.1007/BF01531026)
- [18] Urquhart, A., “The complexity of propositional proofs”, *The Bulletin of Symbolic Logic* 1, 4 (1995): 425–467. <https://projecteuclid.org/euclid.bsl/1181154880>; DOI: [10.2307/421131](https://doi.org/10.2307/421131)

DANIIL KOZHEMIACHENKO
Department of Philosophy
Moscow State University
Lomonosovsky prospekt, 27-4, GSP-1
Moscow 119991, Russian Federation
kodaniil@yandex.ru