

LI, Tianwei, WANG, Lei and YU, Songmao. The impacts of lockdown restrictions on children's and adolescents' physical activity levels and sedentary behaviour including gender differences: A systematic review and meta-analysis. *Quality in Sport*. 2025;37:57482. eISSN 2450-3118.  
<https://doi.org/10.12775/QS.2025.37.57482>  
<https://apcz.umk.pl/QS/article/view/57482>

The journal has been 20 points in the Ministry of Higher Education and Science of Poland parametric evaluation. Annex to the announcement of the Minister of Higher Education and Science of 05.01.2024. No. 32553.

Has a Journal's Unique Identifier: 201398. Scientific disciplines assigned: Economics and finance (Field of social sciences); Management and Quality Sciences (Field of social sciences).

Punkty Ministerialne z 2019 - aktualny rok 20 punktów. Załącznik do komunikatu Ministra Szkolnictwa Wyższego i Nauki z dnia 05.01.2024 r. Lp. 32553. Posiada Unikatowy Identyfikator Czasopisma: 201398.

Przypisane dyscypliny naukowe: Ekonomia i finanse (Dziedzina nauk społecznych); Nauki o zarządzaniu i jakości (Dziedzina nauk społecznych).

© The Authors 2025;

This article is published with open access at Licensee Open Journal Systems of Nicolaus Copernicus University in Torun, Poland Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author (s) and source are credited. This is an open access article licensed under the terms of the Creative Commons Attribution Non commercial license Share alike. (<http://creativecommons.org/licenses/by-nc-sa/4.0/>) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.

The authors declare that there is no conflict of interests regarding the publication of this paper.

Received: 01.12.2024. Revised: 05.02.2025. Accepted: 05.02.2025. Published: 08.02.2025.

## **The impacts of lockdown restrictions on children's and adolescents' physical activity levels and sedentary behaviour including gender differences: A systematic review and meta-analysis**

1. Tianwei Li (senior author)

Department of Kinesiology, Southwest University, Chongqing, China

<https://orcid.org/0009-0003-8280-8467>

2. Lei Wang (Corresponding Author)

Department of Kinesiology, Southwest University, Chongqing, China

3. Songmao Yu (second author)

Department of Kinesiology, Southwest University, Chongqing, China

<https://orcid.org/0009-0004-2168-5798>

### **Abstract**

**Background:** The 2019 COVID-19 pandemic outbreak has caused a global public health emergency. In addition to preventing the spread of the disease, restrictions such as school closures may reduce children and adolescents' opportunities for physical activity and increase the prevalence of sedentary behaviour. However, the current evidence regarding the effects of COVID-19 on physical activity and sedentary behaviour in children and adolescents is insufficiently clear.

**Objectives:** This systemic review and meta-analysis aimed to examine the changes of children's and adolescents' total physical activity, moderate to vigorous physical activity, sedentary behaviour, and screen time before and during COVID-19-related restrictions. Subgroup analysis was aimed to compare the changes between boys and

girls, objective measures and subjective measures, cross-sectional studies and longitudinal studies.

**Methods:** A total of five databases were searched. In addition, grey literature, reference list and registries were also checked. A total of 26 studies were included in the systematic review. Total physical activity, moderate to vigorous physical activity, sedentary and ST were analysed separately using random effects models. Additionally, this study also used meta-analysis to investigate the differences between boys and girls, objective measures and subjective measures, cross-sectional studies and longitudinal studies.

**Results:** The results of the meta-analysis showed that children and adolescents had a significant decrease in total physical activity (SMD= -0.76, 95% CI= -1.46 to -0.06,  $I^2= 99.5\%$ ), moderate to vigorous physical activity (SMD= -0.36, 95% CI= -0.6 to -0.11,  $I^2= 94.7\%$ ), sedentary behaviour (SMD= 1.44, 95% CI= 0.31 to 2.56,  $I^2= 98.6\%$ ), and a significant increase in screen time (SMD= 0.75, 95% CI= 0.49 to 1.02,  $I^2= 97.3\%$ ). Meta-analysis results showed a significant decline in moderate to vigorous physical activity in boys, but no change was found in girls. However, results from the systematic review suggested that girls are more likely to develop more screen time than boys

**Conclusion:** Children and adolescents' physical activity and sedentary behaviour are negatively impacted by restrictions associated with Covid-19. In addition, girls' screen time is more likely to increase than boys'.

Keywords :Physical activity, Youths, Lockdown

## **1.0. Introduction :**

### *1.1. The impact of COVID-19*

In December 2019, a new coronavirus (COVID-19) was found to be the cause of a local outbreak of pneumonia in China, which resulted in the collapse of healthcare systems and millions of deaths all over the world (Tan et al., 2020; Torres-Signes et al., 2021). On January 30, 2020 (WHO, 2020), the World Health Organization (WHO) declared it a global public health emergency. COVID-19 is primarily transmitted between humans via aerosols (WHOa, 2020). Therefore, social distancing has been shown to be an important component of measures designed to reduce the spread and impact of COVID-19 (Berry et al., 2022). Children and adolescents are a population group that has been hit particularly hard by the restrictions despite the fact that they are less likely to develop severe symptoms and are mainly spared the direct health effects of COVID-19 (Kang et al., 2020; Bialek et al., 2020). The indirect effects of COVID-19-related restrictions are highly likely to have negative impacts on children's and adolescents' lifestyles, health, and wellbeing (Richardson et al., 2021).

Restrictions such as lockdowns, school closures, and social isolation have been implemented globally to contain the COVID-19 contagion. Indeed, more than half of the world's population was subject to some form of government-imposed restrictions in April 2020 (WHO, 2020). According to the United Nations Educational, Scientific, and Cultural Organization (UNESCO), 195 countries closed their schools in April 2020, and the school closures lasted an average of 3.5 months of an academic year (UNESCO, 2021). In addition, an estimated 1.5 billion children and adolescents have

become distant learners as a result of school closures (UNESCO, 2021). Schools are universally regarded as an important venue for health promotion, and their physical education classes, gymnasias, and activity facilities provide many opportunities for children and adolescents to engage in physical activity (van Sluijs et al., 2021). As a result of COVID-19-related restrictions, children and adolescents may spend more time sitting and have fewer opportunities to engage in physical activity.

### *1.2. Physical activity (PA) and sedentary behaviour (SB) among children and adolescents*

PA and SB are two of the many health-related behaviours among children and adolescents that affect their health and well-being (Chaput et al., 2020). Numerous health concerns, including obesity, coronary heart disease, and musculoskeletal health, have been shown to be positively influenced by the promotion of PA among children and adolescents (Brown et al., 2009; Janssen et al., 2010; Wu et al., 2017). The positive effects of reducing SB include a decreased risk of obesity, cardiovascular disease, and all-cause mortality, as well as a variety of mental health problems (Prentice-Dunn et al., 2012; Wu et al., 2017). Additionally, children and adolescents with more PA and less SB may benefit from an improvement in cognitive development and academic performance (Tremblay et al., 2011). It is necessary to note that physical inactivity (PI) and SB are two independent health risk factors (Ekelund et al., 2016). If long periods of daily sitting are unavoidable (such as during a COVID-19 lockdown period), it is essential to remain physically active (Ekelund et al., 2016).

Global PI is already an existing non-infectious public health problem. As the fourth leading cause of death worldwide, PI was characterised as a global pandemic by the Lancet series on PA and health (Kohl et al., 2012). The WHO guidelines suggest that children and adolescents between 5 and 17 years should meet the recommendation of at least 60 minutes of moderate-to-vigorous physical activity (MVPA) per day and limit the amount of time spent being sedentary, particularly the amount of recreational screen time (ST) (WHOc, 2020). However, three-quarters of children and adolescents worldwide do not currently meet the physical activity guidelines (Guthold et al., 2020). In addition, gender differences exist in the PA and SB of children and adolescents. Kalman et al. (2015) reported that girls have a lower PA level than boys. More specifically, 85% of girls and 78% of boys did not meet the WHO PA guidelines globally (Guthold et al., 2020). The Canadian SB guidelines reflect gender differences among children and adolescents and suggest that boys and girls may engage in different SB (Liwander et al., 2013). For example, girls are likely to spend a lot of time on communication-based SB such as phone calls, texting, and instant messaging, while boys are more likely to watch TV and videos or play computer games (Liwander et al., 2013). Consequently, there may be gender differences in the effects of COVID-19-related restrictions on children and adolescents. Considering gender issues in research and policy may permit a more accurate examination of changes in child and adolescent health behaviours and help improve the health of this group during the COVID-19 pandemic.

### *1.3. Current evidence on the impacts of COVID-19-related restrictions on PA and SB*

In 2020, some preliminary cross-sectional studies demonstrated that social restrictions reduced PA and increased SB among children and adolescents (Guerrero et al., 2020; Moore et al., 2020; Mitra et al., 2020). However, these are all cross-sectional studies that collected data by asking participants to recall previous COVID-

19 behaviours and compare them with current behaviours. Due to recall bias, the reported results may not accurately reflect the actual situation (Rudolf et al., 2015; Laeremans et al., 2017). One year later (2021), a scoping review found that four studies identified an increase in PA, while six studies found no significant differences in PA during the pandemic among children and adolescents (Rossi et al., 2021). However, Rossi et al. (2021) only examined PA and did not report on SB-related results. Paterson et al. (2021) also undertook a scoping review and found that PA declined and SB increased, especially with regard to ST (Paterson et al., 2021). Nonetheless, 67.9% of the 150 referenced studies utilised a cross-sectional design and did not assess the quality of data from individual studies (Paterson et al., 2021).

Currently, there are two systematic reviews and one meta-analysis in relation to this topic. One of the systematic reviews by Viner et al. (2021) included 36 studies. However, this study included too few PA and SB studies because it also examined changes in sleep, mental health, and diet, among others (Viner et al., 2021). Only two of six studies demonstrated a significant reduction in PA and a significant increase in ST (Viner et al., 2021). Kharel et al. (2022) also conducted a systematic review demonstrating that 32 studies (78.1%) reported a decrease in PA, and eight (19.5%) studies reported no significant change in PA. However, 11 studies have included children or adolescents with specific diseases, such as physical disabilities, ASD (autism spectrum disorder), or diabetes (Kharel et al., 2022). Some of the studies also included children or infants under 4 years old (2022). Meta-analytical results by Wunsch et al. (2022) reported a slightly negative effect on the PA of children and adolescents, but not significantly (Fisher's  $z = 0.08$ , 95% CI = -0.27 to 0.12). The current review of studies combined subjective and objective measurement methods, and some bias arising from the subjectivity of the former method may influence the results. This is due to the fact that subjective methods of measuring PA and SB tend to overestimate or underestimate results (Rudolf et al., 2015).

The current systematic review and meta-analysis of studies did not separate the analysis of cross-sectional and longitudinal data, or subjective and objective methods. Another research gap is that no studies have applied a meta-analytical approach to the differences between PA and SB. Additionally, the current systematic reviews included literature from before October 2021. However, new studies published in 2022 may provide a larger selection of studies for inclusion in a systematic review of this field. For example, Burkart et al. (2022) compared PA and SB before and during the pandemic with objectively collected data. Therefore, a current systematic review can incorporate some longitudinal studies and also apply a device-based measurement method.

#### 1.4. *Aims and potential value*

To the best of our knowledge, this is the first systematic review and meta-analysis focusing on examining the impacts of COVID-19-related restrictions on children's and adolescents' (5–17 years old) PA and SB. This study aims to review findings from primary studies and apply a meta-analytical procedure to quantitatively estimate changes in PA and SB before and during COVID-19-related restrictions, accompanied by a subgroup analysis of gender difference, study designs (cross-sectional and longitudinal), and measurement methods (subjective and objective) among 5–17-year-old youth. This study may be able to assist policymakers in developing PA and SB interventions for this age group during the COVID-19 pandemic. Investigating the difference between boys and girls could help develop different intervention plans for boys and girls, making the intervention more effective. Furthermore, it is critical to

understand the precise impact of such restrictions on PA and SB in order to better prepare for future waves of COVID-19 or other future pandemics.

## 2.0. Methodology

This review was performed according to Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines and the Cochrane Handbook for systematic reviews of interventions (Moher et al., 2009; Higgins & Thomas, 2019).

### 2.1. Eligibility criteria

This review and meta-analysis included English studies and peer-reviewed articles that addressed changes in children's and adolescents' PA before and during the COVID-19 pandemic. As our focus was on non-interventionist studies, studies were selected for eligibility based on the PECO (participants, exposure, comparison, outcome) rather than the PICO (participants, intervention, comparison, outcome) (Mintzker et al., 2022). Table 1 shows the inclusion and exclusion criteria.

**Table 1.** *The inclusion and exclusion criteria*

	<b>Inclusion</b>	<b>exclusion</b>
<b>Population</b>	Healthy children and adolescents aged 5-17 years old.	Children or adolescents with disabilities or specific health conditions such as physical disability, diabetes, and autism; Children or adolescents who are no longer students; Children or adolescents who are athletes.
<b>Exposure</b>	Changes of PA or SB under COVID-19-related restrictions.	Exposure unrelated to COVID-19 restrictions, such as participation in some kinds of PA or SB intervention plans during COVID.
<b>Comparison</b>	Studies compared the changes in PA/SB data before and during COVID-19 restrictions.	Studies did not compare the changes in PA/SB data before and during COVID-19 restrictions.
<b>Outcome</b>	Studies reported PA data including but not limited to total PA, moderate to vigorous PA (MVPA), vigorous PA, walking, and cycling. Studies reported any form of SB include but not limited to total sitting time, ST, and TV hours.	Studies investigating other health-related behaviours rather than PA/SB; for the meta-analysis, studies that did not report M(SD) data before and during COVID-19 restrictions.

M (SD): Mean (Standard deviation)

### 2.2. Information sources

An online search was conducted on July 10th, 2022 across the following databases: Medline, Embase, SPORTDiscus, Scopus, and Web of Science. Grey literature searching was done through the Policy Commons website, which provides almost 2.5 million full-text think-tank publications, including policy reports, administrative documents, books, projects and campaign documents (Olson, 2021). The search date

limit was set for after December 2019 (the start of COVID-19), which was to ensure that only articles concerning COVID-19 were included.

## *2.2. The searching procedure*

Across these databases, key terms are followed by other key terms and linked by Boolean operators OR and AND: "Child", "Children", "adolescent" "youth", "teenager", "sedentary time", "sedentary behaviour/behavior", "ST", "sitting time", "Physical activity", "Movement behaviour/behavior", "COVID-19", "SARS-CoV". Comprehensive search history records are shown in appendix 1. Furthermore, the reference lists of all primary studies, studies included in the existing review studies, and registered and pre-registered studies on PROSPERO were also examined for additional pertinent studies.

The title and abstract screening and full text screening were performed by Covidence software. Two reviewers (TW L, BY) were involved in the screening process. The primary reviewer TW L completed the title and abstract screening independently. During the full-text screening process, the two reviewers worked independently. All disagreements were resolved through discussion.

## *2.3. Data collection process and data items*

Analysis of the extracted data was informed by three existing systematic reviews in the field and the Cochrane handbook (Higgins & Thomas, 2019; Viner et al., 2021; Kharel et al., 2022; Wunsch et al., 2022). Data extracted included the following: general information (i.e. authors and years, country), participant characteristics (sample size, age, gender), information on methods (aim, sampling time, measurement tools), study design (cross-sectional or longitudinal), outcomes (PA/SB before and during COVID restriction, the changes of PA/SB, statistical results). For PA and SB data extraction, only total MVPA (moderate to vigorous physical activity), SB, sitting time, ST, and TV hours were extracted. Collection methods for the primary data are reported in the data extraction sheets which outline whether the data are from accelerators, self or proxy questionnaires (specific type of questionnaire), or other types of descriptive results (interview results).

Data from the eligible studies were extracted by one author independently (TW L) through Covidence software and exported into a Microsoft EXCEL sheet. The identification of the study type (cross-sectional or longitudinal) and the identification of the measurement methods (objective or subjective) were all done independently by one author (TW L). Points of uncertainty regarding the selection of articles for inclusion in the meta-analysis were discussed with the second reviewer BY, and another collaborator (ZY) served as a mediator until consensus was reached. After completion, the extracted data was double-checked and verified by TW L. To obtain data or additional details not reported in the six studies, one author (TW L) tried to contact the original study authors (Bingham et al., 2021, Chambonniere et al., 2021; Dubuc et al., 2021; Gilbert et al., 2021; Guo et al., 2021; Schmidt et al., 2020). However, no response was received from any of the authors. Due to time constraints, the decision was made to proceed and not wait for the authors' replies.

## *2.4. Risk of bias and quality assessment*

Begg's funnel plots for the evaluation of publication bias across studies were implemented using Stata 14.0 software. The sensitivity analysis of the results uses the trim and fill method.

To assess the quality within studies, one author (TW L) independently used the Newcastle-Ottawa Scale (NOS) for cohort studies (Wells GA et al., 2000) and the modified NOS for cross-sectional studies (Modesti et al., 2016). The NOS was compared using a quality assessment tool for observational cohort and cross-sectional studies created by the National Heart, Lung, and Blood Institute (NHLBI). The advantage of NOS is that it has a visual scoring system that rates studies based on three dimensions: selection (maximum 5 stars), comparability (maximum 2 stars) and outcome (maximum 3 stars). The NOS evaluated the appropriateness of the research design, recruitment strategy, response rate, sample representativeness, objectivity/reliability of outcome determination, provided power calculation, and statistical analyses (Modesti et al., 2016). During the quality assessment process, the primary author (TW L) conferred with a collaborator (ZY) and then made the final decision. Table 2 shows the NOS scoring criteria.

**Table 2.** *The scoring of the NOS for cohort and cross-sectional studies*

	<b>Scoring NOS for cohort studies</b>	<b>Scoring NOS for cross-sectional</b>
<b>Method:</b>	Thresholds for converting the Newcastle-Ottawa scale to the Agency for Health Research and Quality (AHRQ) standards: good, fair, and poor (Likis Fe et al., 2012, Vienna, 2015).	The NOS for a cross-sectional study will eventually classify the quality of the study into 4 levels (Modesti et al., 2016).
	<b>Good quality:</b> 3 or 4 stars in selection domain AND 1 or 2 stars in comparability domain AND 2 or 3 stars in outcome/exposure domain.	<b>Very good Studies:</b> 9-10 points
	<b>Fair quality:</b> 2 stars in selection domain AND 1 or 2 stars in comparability domain AND 2 or 3 stars in outcome/exposure domain.	<b>Good studies:</b> 7-8 points
	<b>Poor quality:</b> 0 or 1 stars in selection domain OR 0 stars in comparability domain OR 0 or 1 stars in outcome/exposure domain.	<b>Satisfactory studies:</b> 5-6 points <b>Unsatisfactory studies:</b> 0 to 4 points

### 2.5. Effect measures

Inconsistencies in study results arose from the diversity of measurement methods, which included accelerometer, YAP (Youth Activity Profile), and PAQ-C/A (Physical Activity Questionnaire for Older Children/ Adolescents). Consequently, standardised mean differences (SMD) were utilised in the meta-analysis.

### 2.6. Synthesis methods and results

#### *Systematic review*

General characteristics were tabulated including author and year, sample size, gender information; measurement tool(s), sampling time, study type, statistical significance (P value) or effect size of total PA, MVPA, ST, SB. For studies providing data during

and before COVID, the changes in PA, MVPA, SB, and ST were calculated manually (during minus before).

### Meta-analysis


The meta-analysis was conducted using Stata 14.0 software. Due to the considerable heterogeneity of the included studies ( $I > 75\%$ ), a random effect model will be used (Higgins et al., 2019). Then, PA, MVPA, SB, and ST will be analysed as outcome indicators. Subgroup analysis included gender (boys' total PA vs girls' total PA, boys' MVPA vs girls' MVPA, boys' SB vs girls' SB, boys' ST vs girl's ST), study design (Cross-sectional vs Longitudinal), and measurement method (Subjective vs Objective). Other outcomes (PA, SB, and ST) could not be analysed due to the limited number of device-based studies included. Therefore, MVPA was the only one included in the subjective and objective subgroup analysis.

## 3.0. Results

### 3.1. Study selection:

Figure 1 depicts a PRISMA flowchart of the study selection procedure. The search history and results for each database are shown in appendix 2. Studies were excluded for a variety of reasons: 50% (35) of the exclusions were due to incorrect study design, mostly from studies that included during-COVID data but not pre-COVID data or used after-COVID restrictions data (e.g., Salway et al., 2020); 24% (17) of the exclusions were due to adult studies (e.g., Ning et al., 2021); 17% (12) of excluded results reported changes in behaviour patterns but not PA/SB (e.g., Morrison et al., 2021); and 5% of the exclusions were due to the study population being 0–4-year-olds (e.g., Okely, 2021). The details of studies that were excluded can be seen in appendix 3.

**Figure 1.** PRISMA flow diagram. In this diagram, 493 studies came from Embase & Medline: 206, Web of science: 276, SPORTDiscus: 6, Scopus: 5, Policy commons: 0)



### 3.2. Study characteristics

Our meta-analysis included studies from Tunisia, Saudi Arabia, Qatar, UK (N = 2), USA (N = 3), France (N = 2), Italy (N = 2), Canada, Austria, China (N = 3), Croatia, Germany (N = 2), Spain (N = 2), Australia (N = 2), New Zealand, and the Czech Republic. The total sample is 42,418 (15,502 boys, 20,425 girls). One study did not



provide gender information (Genin et al., 2021) and one study only included boys (Peddie et al., 2021). Only three studies used accelerometers to report results of PA and SB (Burkart et al., 2022, Dallolio et al., 2022, García-Alonso et al., 2022). One study used ecological momentary assessment (EMA) (Munasinghe et al., 2020) and one study was conducted through interviews (Gilbert et al., 2021). The rest of the study results are all from self-report or proxy report (N = 20). Eleven of the included studies utilised a longitudinal design to compare data collected during the pandemic to data collected before the pandemic (Munasinghe et al., 2020; Schmidt et al., 2020; Dubuc et al., 2021; James et al., 2021; Medrano et al., 2021; Nathan et al., 2021, Peddie et al., 2021; Stverakova, 2021; Burkart et al., 2022; Dallolio et al., 2022; Garca-Alonso et al., 2022). Other studies employed a cross-sectional design, used a retrospective baseline and obtaining pre-pandemic data through memory recall (N = 15). Table 3 provides an outline of the individual studies, and their complete results are available in Appendix 4.

### 3.3. Risk of bias within studies

With the NOS of the risk of bias within longitudinal studies, four studies were of good quality, seven were of fair quality, and one was of poor quality. For the modified NOS quantification in cross-sectional studies, nine studies were satisfactory, and five studies were good studies. Details of the specific ratings are shown in Figure 2. The primary limitation of longitudinal studies was sample selection. The majority of longitudinal studies have a small sample size, which may result in some bias in the findings (e.g., Dallolio et al., 2022 (N = 77), Burkart et al., 2022 (N = 74), Garca-Alonso et al., 2022 (N = 110), Peddie et al., 2021 (N = 109), Stverakova 2021 (N = 98). In cross-sectional studies, the fact that results are reported through self-reporting methods is the most significant limitation. All accelerometer-based studies are longitudinal.

**Figure 2.** The risk of bias in individual studies. Selection (maximum 5 stars), comparability (maximum 2 stars) and outcome (maximum 3 stars).


Characteristics of included studies	Quality assessment (NOS/ Adapted NOS for cross-sectional studies)			Study type	Scores for cross-sectional studies or longitudinal studies
Author (s) Year/ Country	Selection	Comparability	Outcome		Scores
Abid et al., 2021	★★	★	★★	Cross-sectional	Satisfactory studies: 5-6 points
Alanazi et al., 2022	★★★	★	★★	Cross-sectional	Satisfactory studies: 5-6 points
Al-Mulla et al., 2022	★★	★	★★	Cross-sectional	Satisfactory studies: 5-6 points
Bingham et al., 2021	★★★	★	★★	Longitudinal	Good quality
Burkart et al., 2022	★★	★★	★★★	Longitudinal	Fair quality
Chambonniere et al., 2021	★★★	★	★★	Cross-sectional	Satisfactory studies: 5-6 points
Dallolio et al., 2022	★★	★★	★★★	Longitudinal study	Fair quality

Dunton et al., 2020	★★	★	★★	Cross-sectional	Satisfactory studies: 5-6 points
Dubuc et al., 2021	★★★★	★	★★	Longitudinal study	Good quality
Farello et al., 2022	★★★★★	★	★★	Cross-sectional	Good studies: 7-8 points
García-Alonso et al., 2022	★★	★★	★★	Longitudinal study	Fair quality
Genin et al., 2021	★★★★★	★	★★	Cross-sectional	Good studies: 7-8 points
Gilbert et al., 2021	★★★★	★	★★★★	Cross-sectional	Good studies: 7-8 points
Greier et al., 2021	★★	★	★★	Cross-sectional	Satisfactory studies: 5-6 points
Guo et al., 2021	★★★★	★★	★	Cross-sectional	Satisfactory studies: 5-6 points
James et al., 2021	★★★★	★★	★★	Longitudinal study	Good quality
Jia et al., 2021	★★★★	★★	★★	Cross-sectional	Good studies: 7-8 points
Jovanovic et al., 2021	★★	★★	★★	Cross-sectional	Satisfactory studies: 5-6 points
Marckhoff et al., 2022	★★★★★	★	★★	Cross-sectional	Good studies: 7-8 points
Medrano et al., 2021	★★	★★	★★	Longitudinal	Fair quality
Munasinghe et al., 2020	★★	★	★★	Longitudinal	Fair quality
Nathan et al., 2021	★★	★	★★	Longitudinal	Fair quality
Peddie et al., 2021	★	★★	★★	Longitudinal	Poor quality
Schmidt et al., 2020	★★★★★	★	★★	Longitudinal	Good quality
Stverakova 2021	★★	★	★★	Longitudinal	Fair quality
Yang et al., 2020	★★★★	★	★★	Cross-sectional	Satisfactory studies: 5-6 points

### 3.4. Risk of bias across studies


Publication bias across studies was assessed using funnel plots for the four outcomes (PA, MVPA, SB, ST). The symmetry of the funnel plot was judged subjectively by visual inspection. It was discovered that the funnel plot for ST exhibited asymmetrical scattering, but the funnel plot for PA had equal distribution on both sides. Therefore, there was probably publication bias in ST. No publication bias analysis was conducted due to the insufficient number (< 10 studies) of MVPA (N = 4) and SB (N = 4) studies. However, it is important to note that the funnel plot contains fewer than ten studies (PA = 9, ST = 7). Therefore, the funnel plot may not be a valid measure of publication bias and caution is needed in the interpretation of the results. The funnel plots can be seen in Figures 3 and 4.

**Figure 3. Total PA's funnel plot**



X-axis: standard error; Y-axis: standardised mean difference

**Figure 4. ST's funnel plot after filled (trim and filling analysis)**



X-axis: standard error; Y-axis: standardised mean difference: original studies: ○ filled studies. ◻

### 3.5. Results of individual studies

In primary studies where total PA, MVPA, SB, or ST were not reported, the PAQ scores, MET-min/week, and days of activity were reported as total PA. Regardless of whether PA guidelines (MVPA over 60 min per day) were met, was recorded as MVPA. Total sitting time was recorded as SB. The study includes a variety of effect measurement methods including median (25%, 75%), mean and standard deviation, percentage of PA/SB/ST guidelines met, and a comparison of percentage of SB/PA hours performed (e.g., SB<4hours/day: N (%) before COVID vs N (%) during COVID, PA>60mins: N(%) vs N(%) during COVID). P-values were used for most

outcome measures, with a few studies using odds ratios (OR) and 95% confidence intervals (CI) (N = 1) and descriptive outcomes (N = 1).

### 3.6. Results of systematic review

Overall, 12 of 17 studies reported a significant decrease in total PA during the COVID-19-related restriction period compared to the pre-COVID-19 period. Ten of 13 studies reported a significant increase in SB, and 12 of 14 studies reported a significant increase in ST. Six of nine studies reported a significant difference in MVPA reduction, and one study reported a statistically significant difference in MVPA reduction. Aside from this, all 22 studies reported that SB or ST increased during the pandemic, with 15 of these studies indicating a significant increase. The two studies that assessed television hours also revealed a significant increase.

However, three studies reported an increase in total PA (Schmidt et al., 2020, Nathan et al., 2021, James et al., 202) and one reported an increased MVPA in boys (Peddie et al., 2021). Notably, all studies demonstrating an increase in PA are longitudinal studies. Furthermore, the increase in PA in these four studies did not vary statistically from pre-pandemic levels, and only one study found a significant increase (James et al., 2021).

There were conflicting results regarding how gender differences affected changes in PA and SB. Eight of the included studies provided data on boys and girls during and before the epidemic, however, only four of the eight studies found a significant difference between boys and girls. Abid et al. (2021) reported a greater increase in ST among girls than boys. Jovanovic et al. (2021) reported that boys participated in significantly more physical activity per week than girls. On the other hand, Dallolio et al. (2022) found that during the epidemic, boys were more restricted and participated in more SB and less total PA and MVPA than girls (Dallolio et al., 2022). Moreover, Marckhoff et al. (2022) found that boys' total PA decreased more than girls', and also reported an overall increase in ST.


### 3.7. Results of the meta-analysis

Each meta-analysis outcome of total PA, MVPA, SB, and ST is presented in Table 4. The meta-analysis outcome regarding gender difference is presented in Table 5. Table 6 presents the results of the meta-analysis based on the results of the device measurements versus the subjective measurements. The results of the meta-analysis based on cross-sectional and longitudinal investigations are presented in Table 7.

**Table 4.** *Meta-analysis results of total PA, MVPA, SB, and ST*

Outcome on total PA: Nine studies assessed total PA in the meta-analysis. The results show that during the COVID-19 pandemic, total PA decreased significantly (SMD = -0.76, 95% CI = -1.46 to -0.06, I <sup>2</sup> = 99.5%).
---


**Figure 5. Total PA standardised mean difference (SMD)**



**Outcome on MVPA:**

Four studies assessed MVPA in the meta-analysis, and the result indicated a significant drop in MVPA during the COVID-19 pandemic. (SMD = -0.36, 95% CI = -0.6 to -0.11,  $I^2 = 94.7\%$ ).


**Figure 6. MVPA standardised mean difference (SMD)**



**Outcome on SB:**

The results of the meta-analysis showed that SB significantly increased during the COVID-19 pandemic (SMD = 1.44, 95% CI = 0.31 to 2.56,  $I^2 = 98.6\%$ ).


**Figure 7. SB standardised mean difference (SMD)**



**Outcome on ST**

Seven studies assessed ST in the meta-analysis. According to the meta-analysis, ST increased significantly (SMD = 0.75, 95% CI = 0.49 to 1.02,  $I^2 = 97.3\%$ ) during the COVID-19 pandemic.

**Figure 8. ST standardised mean difference (SMD)**




These four outcomes (total PA, MVPA, SB, and ST) demonstrated excessive heterogeneity ( $I^2 >75\%$ ) (Higgins & Thomas, 2019). Hence, it was necessary to perform a subgroup analysis to explore the possible causes of heterogeneity.

**Table 5. Meta-analysis results of gender difference among total PA, MVPA, SB, and ST**


Outcome on boys' total PA compared with total PA:

Figures 9 and 10 show the meta-analysis of boys' total PA and girls' total PA. The results showed that total PA decreased significantly during the pandemic for both boys (SMD = -1.10, 95%CI = -1.89 to -0.31,  $I^2 = 98.3%$ ) and girls (SMD = -0.94, 95%CI = -1.77 to -0.11,  $I^2 = 98.8%$ ).

**Figure 9. Boys' total PA standardised mean difference (SMD)**



**Figure 10. Girls' total PA standardised mean difference (SMD)**




Outcome on boys' MVPA compare with girls' MVPA:

Figures 11 and 12 show the meta-analysis of boys' MVPA and girls' MVPA. The results indicate a significant decrease in MVPA during COVID-19 in boys (SMD = -0.26, 95%CI = -0.44 to -0.08,  $I^2 = 84.5%$ ), but not in girls (SMD = -0.16, 95%CI = -0.47 to 0.16,  $I^2 = 96.8%$ ).




**Figure 11. Boys' MVPA standardised mean difference (SMD)**



NOTE: Weights are from random-effects model

**Figure 12. Girls' MVPA standardised mean difference (SMD)**




NOTE: Weights are from random-effects model


**Outcome on boys' SB compare with girls' SB:**

The meta-analysis outcome of SB for boys and girls can be seen in Figure 13 and Figure 14. The results indicate a significant drop in SB for both boys (SMD = 1.73, 95%CI = 0.10 to 3.35,  $I^2 = 98.2\%$ ) and girls (SMD = 1.43, 95%CI = 0.07 to 2.78,  $I^2 = 97.2\%$ ) during COVID-19.

**Figure 13. Boys' SB standardised mean difference (SMD)**




**Figure 14. Girls' SB standardised mean difference (SMD)**




**Outcome on boys' ST compare with girls' ST:**

The results of the ST meta-analysis for boys and girls are depicted in Figures 15 and 16. During the COVID-19, ST decreased significantly for both boys (SMD = 0.77, 95%CI = 0.36 to 1.18,  $I^2 = 97.1\%$ ) and girls (SMD = 0.93, 95%CI = 0.58 to 1.28,  $I^2 = 97.4\%$ ).

**Figure 15. Boys' ST standardised mean difference (SMD)**




**Figure 16. Girls' ST standardised mean difference (SMD)**




**Table 6.** Meta-analysis results on objective versus subjective measures of MVPA

Both objective (SMD = -0.61, 95%CI = -1.21 to -0.01,  $I^2 = 83.9\%$ ) and subjective (SMD = -0.61, 95%CI = -1.01 to -0.20,  $I^2 = 98.5\%$ ) measures showed a significant decrease in MVPA. The results of the included objective measures were based on accelerometers and the results of the subjective studies were derived from self or others' report questionnaires. As there were only two studies with subjective or objective measures for Total PA, SB, and ST, no comparative analysis was conducted.

**Figure 17.** MVPA: objective methods (SMD)



**Figure 18.** MVPA: subjective methods (SMD)




**Table 7.** Meta-analysis results on cross-sectional studies versus longitudinal studies in total PA, SB and ST. (Only two cross-sectional studies of the MVPA were available, which did not allow for meta-analysis)


Outcomes on cross-sectional and longitudinal studies of total PA:

Total PA decreased in the cross-sectional study (SMD = -1.00, 95% CI = -2 to -0.00,  $I^2 = 99.5\%$ ), but not in the longitudinal study (SMD = -0.56, 95% CI = -1.31 to 0.18,  $I^2 = 98.3\%$ ). Figure 19 shows the meta-analysis of cross-sectional studies, and Figure 20 shows the longitudinal studies.

**Figure 19.** Cross-sectional studies of total PA (SMD)




**Figure 20.** Longitudinal studies of total PA (SMD)



Outcomes on longitudinal studies of MVPA:

Results from the longitudinal study showed a significant decrease in MVPA during COVID-19 (SMD = -0.77, 95%CI = -1.37 to -0.17,  $I^2 = 96.2\%$ ). As only one cross-sectional study reported on the MVPA, a meta-analysis of cross-sectional study of MVPA could not be conducted. Figure 20 shows the meta-analysis of longitudinal studies in MVPA.


**Figure 21. Longitudinal studies of MVPA (SMD)**




**Outcomes on cross-sectional and longitudinal studies of SB:**

The cross-sectional studies found no increase in SB (SMD = -1.03, 95% CI = -0.44 to 2.31,  $I^2 = 98.6\%$ ), but the longitudinal study found an increase of SB (SMD = 1.48, 95% CI = 0.21 to 2.75,  $I^2 = 97.3\%$ ).

**Figure 22.** Cross-sectional studies of SB (SMD)



**Figure 23.** Longitudinal studies of SB (SMD)



**Outcomes on cross-sectional and longitudinal studies of ST:**

The results of the meta-analysis were consistent in both cross-sectional (SMD = 0.84, 95%CI = 0.40 to 1.29,  $I^2 = 98.3\%$ ) and longitudinal (SMD = 0.76, 95%CI = 0.37 to 1.15,  $I^2 = 88.5\%$ ) results for ST, both showing a significant increase during COVID-19. Figure 22 displays the cross-sectional meta-analysis results. Figure 23 displays the results of meta-analysis of longitudinal studies.

### 3.8. Heterogeneity and sensitivity analysis

It is noteworthy that after the subgroup analyses by gender, measurement method, and study design, there was still high heterogeneity with I-square > 75% (Higgins & Thomas, 2019). However, in the subgroup analysis based on accelerometer measurements, I-square was reduced to 83.9%. In addition, in the subgroup of boys' MVPA, I-square was reduced to 84.5%, and reduced to 88.5% in the longitudinal studies of ST. Although the reduction in heterogeneity in the subgroup analysis was not statistically significant, it appears that heterogeneity may be related to gender, study design, and data measurement methods. In addition to this, the use of a validated assessment tool, seasonal climate differences at different sampling times, failure to subdivide age groups, and ethnic and racial differences may also cause differences between studies (Ridgers et al., 2015; Lewis et al., 2016; Cross et al., 2020; Saffer et al., 2013). Besides, to further analyse the publication bias of ST in the funnel plot (Figure 4), a sensitivity analysis of the ST results revealed  $P < 0.05$ , indicating that publication bias exists in ST (Figure 24).

**Figure 24.** Sensitivity analysis on ST results

Egger's test						
Std_Eff	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
slope	.0528217	.0748421	0.71	0.512	-.1395662	.2452095
bias	9.0093	3.13267	2.88	0.035	.9565137	17.06209

### 4.0. Discussion

In the results of the meta-analysis, both total PA and MVPA decreased significantly, and both SB and ST increased significantly. The results of the systematic review associated with SB and ST were consistent with those of the meta-analysis. However, some different results of PA emerged from the few studies. There were four studies that revealed an increase in PA or MVPA (Schmidt et al., 2020, Nathan et al., 2021, Peddie et al., 2021, James et al., 2021). In addition, James et al. (2021) and Schmidt et al. (2020) found a significant increase in total PA. Schmidt et al. (2020) also confirmed an increase in habitual PA (playing outside, walking and cycling, gardening, housework), and an increase in adherence to WHO guidelines. The Schmidt et al. (2020) study was conducted during the heaviest lockdown in Germany (2020.04), but the weather in Germany was warmer during that time. In April 2020, Germany recorded a mean temperature of 10.4 °C (9.6 °C in 2019) and an average of 292.4 sunshine hours (227.9 in 2019). In warm settings, higher temperatures are related to greater PA and less ST (Lewis et al., 2016). James et al. (2021) observed an increase in PA among students with a higher socioeconomic status who had easier access to physical activity sites near their residences. Therefore, the studies that found an increase in physical activity were conducted with limited sample sizes and may not be typical of the general population.

In terms of gender differences, the results of the meta-analysis indicated a decline in total PA and MVPA, and an increase in SB and ST for both boys and girls. However, a systematic review revealed contradictory findings regarding gender differences. This study cannot conclusively determine whether there are gender differences affecting changes in PA between boys and girls. However, it appears that girls may have more ST than boys.

For both subjective and objective measures, the results were consistent in terms of changes in MVPA based on accelerometer measurements and questionnaire reports.



Due to the small number of studies included in the analysis and the lack of analysis of total PA, SB, and ST, the results of this study are not sufficient to draw conclusions about the differences between objective and subjective measures. However, in general, self-reported or other reporting methods may underestimate or overestimate true results due to the risk of recall bias (Laeremans et al., 2017, Lee et al., 2011). Therefore, in PA-related fields, there is a preference for relying on the results of objective devices. Under COVID-19-related restrictions, conducting guest observations may be subject to ethical and moral restrictions. Therefore, most of the current research on this topic comes from subjective or objective reporting methods (Paterson et al. 2021). In terms of study design, longitudinal and cross-sectional study differences were found to be inconsistent regarding SB and total PA. This may indicate that differences in study designs can impact the results.

#### 4.1. *Comparison with previous literature*

Compared to previous systematic reviews, this study reveals more alarming effects of COVID-19-related restrictions on PA and SB in children and adolescents. In the prior systematic review, Viner et al. (2021) demonstrated a substantial reduction in PA and an increase in ST. However, only two out of six studies reported significant changes in PA and ST (Lopez Bueno et al., 2020; Pietrobelli et al., 2020). The systematic review by Kharel et al. (2022) indicated a significant decline in PA among children and adolescents, whereas eight studies did not find a significant change and one observed an increase in PA. Interestingly, the study in which Kharel et al. (2022) found an increase in PA is also the same study that showed an increase in PA in our study (Schmidt et al., 2020). In addition, Kharel et al. (2022) reported an increase in ST during the pandemic, which is consistent with the findings of our study. However, Kharel et al. (2022) categorised the varying degrees of COVID-19-related restrictions into 4 different levels, which was not done in our study.

The meta-analysis result by Wunsch et al. (2022) showed a non-significant decrease in PA under the COVID restriction (Fisher's  $z = -0.08$ , 95% CI =  $-0.27$  to  $0.12$ ). The results of our meta-analysis showed significant decreases in total PA and MVPA. In addition, our study also performed a meta-analysis of the changes in SB and ST during the pandemic in children and adolescents, which has never been conducted in previous studies.

#### 4.4. Strengths and limitations of evidence

Overall, our study has several key strengths. First, to the authors' knowledge, this is the first study to use a meta-analytic approach to quantify changes in both PA and SB during COVID in healthy children and adolescents. Previous meta-analysis studies only analysed changes in PA (Wunsch et al., 2022, Neville et al., 2022 ). Previous studies have included different PA domains together in meta-analyses (e.g., light PA, MVPA, outside PA, walking, gardening, TV hours, video game hours) (Wunsch et al., 2022, Neville et al., 2022). As PA and SB are a complex and multidimensional behaviour, this may produce bias in the results (Thompson et al., 2015, Pettee et al., 2012). Separate analyses of total PA and MVPA, SB and ST were also a strength of this study. In addition, this study analysed subjective methods versus objective device-based methods for measuring PA and SB changes for boys and girls, as well as the differences in cross-sectional versus longitudinal outcomes.

However, there are limitations to this study that must be taken into account. Firstly, 53% (14/26) of the articles analysed were cross-sectional design studies. Cross-sectional studies measuring PA and SB data in a single pass and depending on

comparisons of data prior to recall of COVID may cause bias (Rudolf et al., 2015; Laeremans et al., 2017). Although our study analyses longitudinal and cross-sectional studies separately in the meta-analysis, the data's reliability is still unclear due to the limited number of studies included. Secondly, the results of meta-analysis were highly heterogeneous, resulting in less credible conclusions. One of the possible reasons for high heterogeneity may be the large age span of the sample (5-18). However, children (5-10) and adolescents (11-17) were not analysed separately in our study. Therefore, the findings of our meta-analysis must be interpreted with caution. In addition to this, exposure information (e.g., restriction levels) from the initial study was not investigated in this analysis. Our study also did not investigate when COVID-19-related restrictions started in each study, which could be significant as people may change their behaviour under the lockdown period. For example, Dubuc et al. (2021) reported that PA and SB change over the duration of restrictions.

### **5.0. Conclusion and suggestions for future research**

This systematic review and meta-analysis revealed that under COVID-19-related restrictions, healthy children and adolescents had a significant decrease in total PA and MVPA, and a significant rise in SB and ST. These behavioural changes may be different between boys and girls. To prevent future adverse physical and mental impacts on children and adolescents, it is necessary for governments and schools to intervene in the PA and SB of children and adolescents during and after COVID-19, and to consider boys' and girls' gender differences when implementing interventions for PA and SB.

Consistent with previous systematic review recommendations, future study should largely concentrate on gathering data on the same forms of PA or SB using the same assessment tool ( Paterson et al., 2021). This will strengthen the comparability between studies' outcomes. In addition, objective device-based measurements in this field remain rare and it is recommended that objective measurements like accelerometer readings can be used more often when permitted by ethical and moral standards. On the other hand, with the increase in vaccination rates and the gradual reduction or even elimination of COVID-19-related restrictions globally (WHO Coronavirus Dashboard, 2022), it is possible that future studies could focus more on PA and SB changes after COVID-19.

### **Acknowledgements**

Words cannot express my gratitude to my supervisor Dr Graham Baker for his invaluable patience and feedback. I also could not have undertaken this journey without Ms Ishbel Leggat, who generously provided knowledge and expertise for my searching strategies. Additionally, I am also grateful to my classmates, for their help with literature screen and discuss with me. I would also like to thank myself. It was a stressful process and some physical injuries presented huge challenges to me. From this process I have become more determined to continue working toward to a higher degree in the future.

### **Reference**

Abid, R., Ammar, A., Maaloul, R., Souissi, N., & Hammouda, O. (2021). Effect of COVID-19-related home confinement on sleep quality, screen time and physical activity in tunisian boys and girls: A survey. *International Journal of*

- Environmental Research and Public Health, 18(6), 1–12. <https://doi.org/10.3390/ijerph18063065>
- Alanazi, Y. A., Parrish, A.-M., & Okely, A. D. (2022). Impact of the COVID-19 virus outbreak on 24-h movement behaviours among children in Saudi Arabia: A cross-sectional survey. *Child: Care, Health & Development*. <https://doi.org/10.1111/cch.12999>
- Al-Mulla, N. A., & Mahfoud, Z. R. (2022). The Impact of the COVID-19 Lockdown “Home Quarantine” on the Physical Activity and Lifestyle of Children in Qatar. *Frontiers in Public Health*, 10, 877424–877424. <https://doi.org/10.3389/fpubh.2022.877424>
- Ayan, Suarez-Iglesias, D., Lopez-Valenciano, A., & Sanchez-Lastra, M. A. (2021). Impact of COVID-19 Pandemic on University Students’ Physical Activity Levels: An Early Systematic Review. *Frontiers in Psychology*, 11, 624567–624567. <https://doi.org/10.3389/fpsyg.2020.624567>
- Berry, E., Jenkins, C., & Allen, S. (2022). Facilitators and barriers to social distancing for young people during the COVID-19 pandemic. *BMC Public Health*, 22(1), 891–891. <https://doi.org/10.1186/s12889-022-13325-3>
- Bialek, S., Gierke, R., Hughes, M., McNamara, L. A., Pilishvili, T., & Skoff, T. (2020). Coronavirus Disease 2019 in Children - United States, February 12-April 2, 2020. *MMWR. Morbidity and Mortality Weekly Report*, 69(14), 422–426. <https://doi.org/10.15585/MMWR.MM6914E4>
- Bingham, D. D., Daly-Smith, A., Hall, J., Seims, A., Dogra, S. A., Fairclough, S. J., ... Barber, S. E. (2021). Covid-19 lockdown: Ethnic differences in children’s self-reported physical activity and the importance of leaving the home environment; a longitudinal and cross-sectional study from the Born in Bradford birth cohort study. *The International Journal of Behavioral Nutrition and Physical Activity*, 18(1), 1–117. <https://doi.org/10.1186/s12966-021-01183-y>
- Brown, T., & Summerbell, C. (2009). Systematic review of school-based interventions that focus on changing dietary intake and physical activity levels to prevent childhood obesity: an update to the obesity guidance produced by the National Institute for Health and Clinical Excellence. *Obesity reviews : an official journal of the International Association for the Study of Obesity*, 10(1), 110–141. <https://doi.org/10.1111/j.1467-789X.2008.00515.x>
- Burkart, S., Parker, H., Weaver, R. G., Beets, M. W., Jones, A., Adams, E. L., ... Armstrong, B. (2022). Impact of the COVID-19 pandemic on elementary schoolers’ physical activity, sleep, screen time and diet: A quasi-experimental interrupted time series study. *Pediatric Obesity*, 17(1), e12846–n/a. <https://doi.org/10.1111/ijpo.12846>
- Chambonniere, C., Lambert, C., Fearnbach, N., Tardieu, M., Fillon, A., Genin, P., ... Duclos, M. (2021). Effect of the COVID-19 lockdown on physical activity and sedentary behaviors in French children and adolescents: New results from the ONAPS national survey. *European Journal of Integrative Medicine*, 43, 101308–101308. <https://doi.org/10.1016/j.eujim.2021.101308>
- Chaput, J. P., Willumsen, J., Bull, F., Chou, R., Ekelund, U., Firth, J., Jago, R., Ortega, F. B., & Katzmarzyk, P. T. (2020). 2020 WHO guidelines on physical activity and sedentary behaviour for children and adolescents aged 5-17 years: summary of the evidence. *The international journal of behavioral nutrition and physical activity*, 17(1), 141. <https://doi.org/10.1186/s12966-020-01037-z>

- Crawford. (2009). Johns Hopkins Nursing Evidence-Based Practice Model and Guidelines. *AORN Journal*, 89(2), 426–426. [https://doi.org/10.1016/S0001-2092\(09\)00182-3](https://doi.org/10.1016/S0001-2092(09)00182-3)
- Cross, T. J., Isautier, J., Stamatakis, E., Morris, S. J., Johnson, B. D., Wheatley-Guy, C., & Taylor, B. J. (2021). Self-reported physical activity before a COVID-19 'lockdown': is it just a matter of opinion?. *BMJ open sport & exercise medicine*, 7(2), e001088. <https://doi.org/10.1136/bmjsem-2021-001088>
- Dallolio, L., Marini, S., Masini, A., Toselli, S., Stagni, R., Bisi, M. C., ... Ceciliani, A. (2022). The impact of COVID-19 on physical activity behaviour in Italian primary school children: a comparison before and during pandemic considering gender differences. *BMC Public Health*, 22(1), 52–52. <https://doi.org/10.1186/s12889-021-12483-0>
- Dubuc, Berrigan, F., Goudreault, M., Beaudoin, S., & Turcotte, S. (2021). Covid-19 impact on adolescent 24 h movement behaviors. *International Journal of Environmental Research and Public Health*, 18(17), 9256–. <https://doi.org/10.3390/ijerph18179256>
- Dubuc, M.-M., Berrigan, F., Goudreault, M., Beaudoin, S., & Turcotte, S. (2021). Covid-19 impact on adolescent 24 h movement behaviors. *International Journal of Environmental Research and Public Health*, 18(17), 9256–. <https://doi.org/10.3390/ijerph18179256>
- Dunton, G. F., Do, B., & Wang, S. D. (2020). Early effects of the COVID-19 pandemic on physical activity and sedentary behavior in children living in the U.S. *BMC Public Health*, 20(1), 1351–1351. <https://doi.org/10.1186/s12889-020-09429-3>
- Ekelund, U., Steene-Johannessen, J., Brown, W. J., Fagerland, M. W., Owen, N., Powell, K. E., Bauman, A., Lee, I. M., Lancet Physical Activity Series 2 Executive Committee, & Lancet Sedentary Behaviour Working Group (2016). Does physical activity attenuate, or even eliminate, the detrimental association of sitting time with mortality? A harmonised meta-analysis of data from more than 1 million men and women. *Lancet (London, England)*, 388(10051), 1302–1310. [https://doi.org/10.1016/S0140-6736\(16\)30370-1](https://doi.org/10.1016/S0140-6736(16)30370-1)
- Farello, G., D'Andrea, M., Quarta, A., Grossi, A., Pompili, D., Altobelli, E., ... Balsano, C. (2022). Children and Adolescents Dietary Habits and Lifestyle Changes during COVID-19 Lockdown in Italy. *Nutrients*, 14(10), 2135–. <https://doi.org/10.3390/nu14102135>
- Genin, P. M., Lambert, C., Larras, B., Pereira, B., Toussaint, J.-F., Baker, J. S., ... Duclos, M. (2021). How did the COVID-19 confinement period affect our physical activity level and sedentary behaviors? Methodology and first results from the French national ONAPS survey. *Journal of Physical Activity & Health*, 18(3), 296–303. <https://doi.org/10.1123/JPAH.2020-0449>
- German Weather Service. *Monthly Climate Status for Germany. April 2019*. [https://www.dwd.de/DE/leistungen/pbfb\\_verlag\\_monat\\_klimastatus/monat\\_klimastatus.html](https://www.dwd.de/DE/leistungen/pbfb_verlag_monat_klimastatus/monat_klimastatus.html) (2019).
- German Weather Service. *Monthly Climate Status for Germany. April 2020*. [https://www.dwd.de/DE/leistungen/pbfb\\_verlag\\_monat\\_klimastatus/monat\\_klimastatus.html](https://www.dwd.de/DE/leistungen/pbfb_verlag_monat_klimastatus/monat_klimastatus.html) (2020).
- Gilbert, A. S., Schmidt, L., Beck, A., Kepper, M. M., Mazzucca, S., & Eyler, A. (2021). Associations of physical activity and sedentary behaviors with child

- mental well-being during the COVID-19 pandemic. *BMC Public Health*, 21(1), 1770–1770. <https://doi.org/10.1186/s12889-021-11805-6>
- Greier, K., Drenowatz, C., Bischofer, T., Petrasch, G., Greier, C., Cocca, A., & Ruedl, G. (2021). Physical activity and sitting time prior to and during COVID-19 lockdown in Austrian high-school students. *AIMS Public Health*, 8(3), 531–540. <https://doi.org/10.3934/publichealth.2021043>
- Guerrero, M. D., Vanderloo, L. M., Rhodes, R. E., Faulkner, G., Moore, S. A., & Tremblay, M. S. (2020). Canadian children’s and youth’s adherence to the 24-h movement guidelines during the COVID-19 pandemic: A decision tree analysis. *Journal of Sport and Health Science*, 9(4), 313–321. <https://doi.org/10.1016/j.jshs.2020.06.005>
- Guo, Y., Liao, M., Cai, W., Yu, X., Li, S., Ke, X., ... Zeng, F. (2021). Physical activity, screen exposure and sleep among students during the pandemic of COVID-19. *Scientific Reports*, 11(1), 8529–8529. <https://doi.org/10.1038/s41598-021-88071-4>
- Guthold, Stevens, G. A., Riley, L. M., & Bull, F. C. (2020). Global trends in insufficient physical activity among adolescents: a pooled analysis of 298 population-based surveys with 1·6 million participants. *The Lancet Child & Adolescent Health*, 4(1), 23–35. [https://doi.org/10.1016/S2352-4642\(19\)30323-2](https://doi.org/10.1016/S2352-4642(19)30323-2)
- Guthold, Stevens, G. A., Riley, L. M., & Bull, F. C. (2020). Global trends in insufficient physical activity among adolescents: a pooled analysis of 298 population-based surveys with 1·6 million participants. *The Lancet Child & Adolescent Health*, 4(1), 23–35. [https://doi.org/10.1016/S2352-4642\(19\)30323-2](https://doi.org/10.1016/S2352-4642(19)30323-2)
- Hagstromer, M., Ainsworth, B. E., Oja, P., & Sjostrom, M. (2010). Comparison of a subjective and an objective measure of physical activity in a population sample. *Journal of physical activity & health*, 7(4), 541–550. <https://doi.org/10.1123/jpah.7.4.541>
- Higgins, & Thomas, J. (2019). *Cochrane handbook for systematic reviews of interventions*. (Second edition / edited by Julian P.T. Higgins, James Thomas, Jacqueline Chandler, Miranda Cumpston, Tianjing Li, Matthew J. Page and Vivian A. Welch.). Cochrane.
- James, M., Marchant, E., Defeyter, M. A., Woodside, J., & Brophy, S. (2021). Impact of school closures on the health and well-being of primary school children in Wales UK: a routine data linkage study using the HAPPEN Survey (2018–2020). *BMJ Open*, 11(10), e051574–e051574. <https://doi.org/10.1136/bmjopen-2021-051574>
- Janssen, I., & Leblanc, A. G. (2010). Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *The international journal of behavioral nutrition and physical activity*, 7, 40. <https://doi.org/10.1186/1479-5868-7-40>
- Jia, P., Zhang, L., Yu, W., Yu, B., Liu, M., Zhang, D., & Yang, S. (2021). Impact of COVID-19 lockdown on activity patterns and weight status among youths in China: the COVID-19 Impact on Lifestyle Change Survey (COINLICS). *INTERNATIONAL JOURNAL OF OBESITY*, 45(3), 695–699. <https://doi.org/10.1038/s41366-020-00710-4>
- Jovanović, G. K., Zubalj, N. D., Majanović, S. K., Rahelić, D., Rahelić, V., Lončar, J. V., & Žeželj, S. P. (2021). The outcome of COVID-19 lockdown on changes in body mass index and lifestyle among croatian schoolchildren: A cross-sectional study. *Nutrients*, 13(11), 3788–. <https://doi.org/10.3390/nu13113788>

- Kalman, M., Inchley, J., Sigmundova, D., Iannotti, R. J., Tynjälä, J. A., Hamrik, Z., Haug, E., & Bucksch, J. (2015). Secular trends in moderate-to-vigorous physical activity in 32 countries from 2002 to 2010: a cross-national perspective. *European journal of public health*, 25 Suppl 2, 37–40. <https://doi.org/10.1093/eurpub/ckv024>
- Kharel, M., Sakamoto, J. L., Carandang, R. R., Ulambayar, S., Shibamura, A., Yarotskaya, E., ... Jimba, M. (2022). Impact of COVID-19 pandemic lockdown on movement behaviours of children and adolescents: a systematic review. *BMJ Global Health*, 7(1), e007190–. <https://doi.org/10.1136/bmjgh-2021-007190>
- Kohl, Craig, C. L., Lambert, E. V., Inoue, S., Alkandari, J. R., Leetongin, G., & Kahlmeier, S. (2012). The pandemic of physical inactivity: global action for public health. *The Lancet (British Edition)*, 380(9838), 294–305. [https://doi.org/10.1016/S0140-6736\(12\)60898-8](https://doi.org/10.1016/S0140-6736(12)60898-8)
- Laeremans, Dons, E., Avila-Palencia, I., Carrasco-Turigas, G., Orjuela, J. P., Anaya, E., Brand, C., Cole-Hunter, T., De Nazelle, A., Götschi, T., Kahlmeier, S., Nieuwenhuijsen, M., Standaert, A., De Boever, P., & Int Panis, L. (2017). Physical activity and sedentary behaviour in daily life: A comparative analysis of the Global Physical Activity Questionnaire (GPAQ) and the SenseWear armband. *PloS One*, 12(5), e0177765–e0177765. <https://doi.org/10.1371/journal.pone.0177765>
- Lee, P. H., Macfarlane, D. J., Lam, T. H., & Stewart, S. M. (2011). Validity of the International Physical Activity Questionnaire Short Form (IPAQ-SF): a systematic review. *The international journal of behavioral nutrition and physical activity*, 8, 115. <https://doi.org/10.1186/1479-5868-8-115>
- Lewis, Maher, C., Belanger, K., Tremblay, M., Chaput, J.-P., & Olds, T. (2016). At the Mercy of the Gods: Associations between weather, physical activity, and sedentary time in children. *Pediatric Exercise Science*, 28(1), 152–163. <https://doi.org/10.1123/pes.2015-0076>
- Likis FE, Andrews JA, Collins MR, et al. Nitrous Oxide for the Management of Labor Pain. Comparative Effectiveness Review No. 67. (Prepared by the Vanderbilt Evidence-based Practice Center under Contract No. 290-2007-10065-I.) AHRQ Publication No. 12-EHC071-EF. Rockville, MD: Agency for Healthcare Research and Quality; 2012
- Liwander, Pederson, A., & Boyle, E. (2013). Why the Canadian Sedentary Behaviour Guidelines Should Reflect Sex and Gender. *Canadian Journal of Public Health*, 104(7), e479–e481. <https://doi.org/10.17269/cjph.104.4108>
- López-Bueno, López-Sánchez, G. F., Casajús, J. A., Calatayud, J., Gil-Salmerón, A., Grabovac, I., Tully, M. A., & Smith, L. (2020). Health-Related Behaviors Among School-Aged Children and Adolescents During the Spanish Covid-19 Confinement. *Frontiers in Pediatrics*, 8, 573–573. <https://doi.org/10.3389/fped.2020.00573>
- Marckhoff, M., Siebald, M., Timmesfeld, N., Janßen, M., Romer, G., & Föcker, M. (2022). COVID-19: Effects of Pandemic Related Restrictions on Physical Activity, Screen Time, and Mental Well-being in German adolescents. *Zeitschrift für Kinder- und Jugendpsychiatrie und Psychotherapie*, 50(4), 313–326. <https://doi.org/10.1024/1422-4917/a000867>
- Medrano, M., Cadenas-Sanchez, C., Osés, M., Arenaza, L., Amasene, M., & Labayen, I. (2021). Changes in lifestyle behaviours during the COVID-19 confinement in Spanish children: A longitudinal analysis from the MUGI project. *Pediatric Obesity*, 16(4), e12731–n/a. <https://doi.org/10.1111/ijpo.12731>



- Mintzker, Blum, D., & Adler, L. (2022). Replacing PICO in non-interventional studies. *BMJ Evidence-Based Medicine*, *bmjebm-2021-111889*. <https://doi.org/10.1136/bmjebm-2021-111889>
- Mitra, Moore, S. A., Gillespie, M., Faulkner, G., Vanderloo, L. M., Chulak-Bozzer, T., Rhodes, R. E., Brussoni, M., & Tremblay, M. S. (2020). Healthy movement behaviours in children and youth during the COVID-19 pandemic: Exploring the role of the neighbourhood environment. *Health & Place*, *65*, 102418–102418. <https://doi.org/10.1016/j.healthplace.2020.102418>
- Modesti, Reboldi, G., Cappuccio, F. P., Agyemang, C., Remuzzi, G., Rapi, S., Perruolo, E., & Parati, G. (2016). Panethnic Differences in Blood Pressure in Europe: A Systematic Review and Meta-Analysis. *PloS One*, *11(1)*, e0147601–e0147601. <https://doi.org/10.1371/journal.pone.0147601>
- Moher, Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Reprint—Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *Physical Therapy*, *89(9)*, 873–880. <https://doi.org/10.1093/ptj/89.9.873>
- Moore, S. A., Faulkner, G., Rhodes, R. E., Brussoni, M., Chulak-Bozzer, T., Ferguson, L. J., ... Tremblay, M. S. (2020). Impact of the COVID-19 virus outbreak on movement and play behaviours of Canadian children and youth: a national survey. *The International Journal of Behavioral Nutrition and Physical Activity*, *17(1)*, 85–85. <https://doi.org/10.1186/s12966-020-00987-8>
- Morrison, Meh, K., Sember, V., Starc, G., & Jurak, G. (2021). The Effect of Pandemic Movement Restriction Policies on Children’s Physical Fitness, Activity, Screen Time, and Sleep. *Frontiers in Public Health*, *9*, 785679–785679. <https://doi.org/10.3389/fpubh.2021.785679>
- Munasinghe, S., Sperandei, S., Freebairn, L., Conroy, E., Jani, H., Marjanovic, S., & Page, A. (2020). The Impact of Physical Distancing Policies During the COVID-19 Pandemic on Health and Well-Being Among Australian Adolescents. *Journal of Adolescent Health*, *67(5)*, 653–661. <https://doi.org/10.1016/j.jadohealth.2020.08.008>
- Munn, Peters, M. D. J., Stern, C., Tufanaru, C., McArthur, A., & Aromataris, E. (2018). Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Medical Research Methodology*, *18(1)*, 143–143. <https://doi.org/10.1186/s12874-018-0611-x>
- Nathan, A., George, P., Ng, M., Wenden, E., Bai, P., Phiri, Z., & Christian, H. (2021). Impact of covid-19 restrictions on western Australian children’s physical activity and screen time. *International Journal of Environmental Research and Public Health*, *18(5)*, 1–13. <https://doi.org/10.3390/ijerph18052583>
- Neville, Lakes, K. D., Hopkins, W. G., Tarantino, G., Draper, C. E., Beck, R., & Madigan, S. (2022). Global Changes in Child and Adolescent Physical Activity During the COVID-19 Pandemic: A Systematic Review and Meta-analysis. *JAMA Pediatrics*. <https://doi.org/10.1001/jamapediatrics.2022.2313>
- Ning, Chen, S.-T., Chi, X., & Liang, K. (2021). Changes in physical activity and sedentary behavior in chinese young adults during the covid-19 pandemic: A repeated-measure study throughout one year. *Healthcare (Basel)*, *9(11)*, 1404–. <https://doi.org/10.3390/healthcare9111404>
- Okely, Kariippanon, K. E., Guan, H., Taylor, E. K., Suesse, T., Cross, P. L., Chong, K. H., Suherman, A., Turab, A., Staiano, A. E., Ha, A. S., El Hamdouchi, A., Baig, A., Poh, B. K., Del Pozo-Cruz, B., Chan, C. H. S., Nyström, C. D., Koh, D.,

- Webster, E. K., ... Draper, C. E. (2021). Global effect of COVID-19 pandemic on physical activity, sedentary behaviour and sleep among 3- to 5-year-old children: a longitudinal study of 14 countries. *BMC Public Health*, 21(1), 940–940. <https://doi.org/10.1186/s12889-021-10852-3>
- Olson, Lauris. “Policy Commons – International Think Tank Publications Fulltext.” *Penn Libraries*, 14 June 2021, [www.library.upenn.edu/blogs/libraries-news/policy-commons-international-think-tank-publications-fulltext](http://www.library.upenn.edu/blogs/libraries-news/policy-commons-international-think-tank-publications-fulltext).
- Paterson, D. C., Ramage, K., Moore, S. A., Riazi, N., Tremblay, M. S., & Faulkner, G. (2021). Exploring the impact of COVID-19 on the movement behaviors of children and youth: A scoping review of evidence after the first year. *Journal of Sport and Health Science*, 10(6), 675–689. <https://doi.org/10.1016/j.jshs.2021.07.001>
- Peddie, M. C., Scott, T., & Haszard, J. J. (2021). Using a 24 h activity recall (Star-24) to describe activity in adolescent boys in new zealand: Comparisons between a sample collected before, and a sample collected during the covid-19 lockdown. *International Journal of Environmental Research and Public Health*, 18(15), 8035–. <https://doi.org/10.3390/ijerph18158035>
- Pettee Gabriel, Morrow Jr, J. R., & Woolsey, A.-L. T. (2012). Framework for physical activity as a complex and multidimensional behavior. *Journal of Physical Activity & Health*, 9(s1), S11–18. <https://doi.org/10.1123/jpah.9.s1.s11>
- Pietrobelli, Pecoraro, L., Ferruzzi, A., Heo, M., Faith, M., Zoller, T., Antoniazzi, F., Piacentini, G., Fearnbach, S. N., & Heymsfield, S. B. (2020). Effects of COVID-19 Lockdown on Lifestyle Behaviors in Children with Obesity Living in Verona, Italy: A Longitudinal Study. *Obesity (Silver Spring, Md.)*, 28(8), 1382–1385. <https://doi.org/10.1002/oby.22861>
- Pombo, Luz, C., Rodrigues, L. ., Ferreira, C., & Cordovil, R. (2020). Correlates of children’s physical activity during the COVID-19 confinement in Portugal. *Public Health (London)*, 189, 14–19. <https://doi.org/10.1016/j.puhe.2020.09.009>
- Prentice-Dunn, H., & Prentice-Dunn, S. (2012). Physical activity, sedentary behavior, and childhood obesity: a review of cross-sectional studies. *Psychology, health & medicine*, 17(3), 255–273. <https://doi.org/10.1080/13548506.2011.608806>
- Reilly, Penpraze, V., Hislop, J., Davies, G., Grant, S., & Paton, J. Y. (2008). Objective measurement of physical activity and sedentary behaviour: review with new data. *Archives of Disease in Childhood*, 93(7), 614–619. <https://doi.org/10.1136/adc.2007.133272>
- Richardson, D. L., Duncan, M. J., Clarke, N. D., Myers, T. D., & Tallis, J. (2021). The influence of COVID-19 measures in the United Kingdom on physical activity levels, perceived physical function and mood in older adults: A survey-based observational study. *Journal of Sports Sciences*, 39(8), 887–899. <https://doi.org/10.1080/02640414.2020.1850984>
- Rossi, Behme, N., & Breuer, C. (2021). Physical activity of children and adolescents during the COVID-19 pandemic—A scoping review. *International Journal of Environmental Research and Public Health*, 18(21), 11440–. <https://doi.org/10.3390/ijerph182111440>
- Ridgers, N. D., Salmon, J., & Timperio, A. (2015). Too hot to move? Objectively assessed seasonal changes in Australian children's physical activity. *The international journal of behavioral nutrition and physical activity*, 12, 77. <https://doi.org/10.1186/s12966-015-0245-x>



- Rudolf, Schaller, A., Frick, F., Grieben, C., & Froböse, I. (2015). Erfassung der Selbsteinschätzung körperlicher Aktivität von jungen Erwachsenen: Eine Querschnittstudie mit Berufsschülern. *Prävention und Gesundheitsförderung*, 11(1), 20–26. <https://doi.org/10.1007/s11553-015-0525-y>
- Saffer H, Dave D, Grossman M, Leung LA. Racial, Ethnic, and Gender Differences in Physical Activity. *J Hum Cap*. 2013 Winter;7(4):378-410. doi: 10.1086/671200. PMID: 25632311; PMCID: PMC4306423.
- Salway, Foster, C., de Vocht, F., Tibbitts, B., Emm-Collison, L., House, D., Williams, J. G., Breheny, K., Reid, T., Walker, R., Churchward, S., Hollingworth, W., & Jago, R. (2022). Accelerometer-measured physical activity and sedentary time among children and their parents in the UK before and after COVID-19 lockdowns: a natural experiment. *The International Journal of Behavioral Nutrition and Physical Activity*, 19(1), 51–51. <https://doi.org/10.1186/s12966-022-01290-4>
- Schmidt, S. C. E., Anedda, B., Burchartz, A., Eichsteller, A., Kolb, S., Nigg, C., ... Woll, A. (2020). Physical activity and screen time of children and adolescents before and during the COVID-19 lockdown in Germany: a natural experiment. *Scientific Reports*, 10(1), 21780–21780. <https://doi.org/10.1038/s41598-020-78438-4>
- Stockwell, S., Trott, M., Tully, M., Shin, J., Barnett, Y., Butler, L., McDermott, D., Schuch, F., & Smith, L. (2021). Changes in physical activity and sedentary behaviours from before to during the COVID19 pandemic lockdown: a systematic review. *BMJ open sport & exercise medicine*, 7(1), e000960. <https://doi.org/10.1136/bmjsem2020-00096>
- Štveráková, T., Jačisko, J., Busch, A., Šafářová, M., Kolář, P., & Kobesová, A. (2021). The impact of COVID-19 on physical activity of Czech children. *PLoS One*, 16(7), e0254244–e0254244. <https://doi.org/10.1371/journal.pone.0254244>
- Tan, W., Zhao, X., Ma, X., Wang, W., Niu, P., Xu, W., ... Wu, G. (2020). A Novel Coronavirus Genome Identified in a Cluster of Pneumonia Cases — Wuhan, China 2019–2020. *China CDC Weekly*, 2(4), 61–62. <https://doi.org/10.46234/ccdcw2020.017>
- The United Nations Educational, Scientific and Cultural Organization (UNESCO). (2021). UNESCO figures show two thirds of an academic year lost on average worldwide due to Covid-19 school closures.
- Thompson, Peacock, O., Western, M., & Batterham, A. M. (2015). Multidimensional physical activity: An opportunity, not a problem. *Exercise and Sport Sciences Reviews*, 43(2), 67–74. <https://doi.org/10.1249/JES.0000000000000039>
- Torres–Signes, A., Frías, M. P., & Ruiz-Medina, M. D. (2021). COVID-19 mortality analysis from soft-data multivariate curve regression and machine learning. *Stochastic Environmental Research and Risk Assessment*, 35(12), 2659–2678. <https://doi.org/10.1007/s00477-021-02021-0>
- Tremblay, M. S., LeBlanc, A. G., Kho, M. E., Saunders, T. J., Larouche, R., Colley, R. C., Goldfield, G., & Connor Gorber, S. (2011). Systematic review of sedentary behaviour and health indicators in school-aged children and youth. *The international journal of behavioral nutrition and physical activity*, 8, 98. <https://doi.org/10.1186/1479-5868-8-98>
- van Sluijs, Ekelund, U., Crochemore-Silva, I., Guthold, R., Ha, A., Lubans, D., Oyeyemi, A. L., Ding, D., & Katzmarzyk, P. T. (2021). Physical activity behaviours in adolescence: current evidence and opportunities for intervention.

- The Lancet (British Edition), 398(10298), 429–442.  
[https://doi.org/10.1016/S0140-6736\(21\)01259-9](https://doi.org/10.1016/S0140-6736(21)01259-9)
- Vienna. “Comparison of Newcastle Ottawa Scale (NOS) and Agency for Health Research and Quality (AHRQ) as Risk of Bias Assessment Tools for Cohort Studies | Colloquium Abstracts.” *Abstracts.cochrane.org*, 2015, [abstracts.cochrane.org/2015-vienna/comparison-newcastle-ottawa-scale-nos-and-agency-health-research-and-quality-ahrq-risk](https://abstracts.cochrane.org/2015-vienna/comparison-newcastle-ottawa-scale-nos-and-agency-health-research-and-quality-ahrq-risk). Accessed 16 Aug. 2022.
- Viner, Russell, S., Saulle, R., Croker, H., Stansfield, C., Packer, J., Nicholls, D., Goddings, A.-L., Bonell, C., Hudson, L., Hope, S., Ward, J., Schwalbe, N., Morgan, A., & Minozzi, S. (2022). School Closures During Social Lockdown and Mental Health, Health Behaviors, and Well-being Among Children and Adolescents During the First COVID-19 Wave: A Systematic Review. *JAMA Pediatrics*, 176(4), 400–409. <https://doi.org/10.1001/jamapediatrics.2021.5840>
- Violant-Holz, Gallego-Jiménez, M. G., González-González, C. S., Muñoz-Violant, S., Rodríguez, M. J., Sansano-Nadal, O., & GuerraBalic, M. (2020). Psychological health and physical activity levels during the covid-19 pandemic: A systematic review. *International Journal of Environmental Research and Public Health*, 17(24), 1–19. <https://doi.org/10.3390/ijerph17249419>
- Wells GA, Shea B, O’Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Ottawa Hospital Research Institute; 2011. Accessed December 14, 2020. [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp)
- Wells GA, Shea B, O’Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses [webpage on the Internet]. Ottawa, ON: Ottawa Hospital Research Institute; 2011. Available from: [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp). Accessed February 5, 2013.
- White, L. E., Barreira, T. V., & Norris, M. L. (2021). The Impact Of COVID-19 On Physical Activity And Sedentary Behavior In Children: A Pilot Study: 580. *Medicine and Science in Sports and Exercise*, 53(8S), 194–194. <https://doi.org/10.1249/01.mss.0000761320.46497.f9>
- WHO COVID-19 Dashboard. Geneva: World Health Organization, 2022. Available online: <https://covid19.who.int/> (last cited: [date]).
- World Health Organization (WHOa). 2020. “COVID-19 Public Health Emergency of International Concern (PHEIC) Global Research and Innovation Forum.” February 12, 2020. [https://www.who.int/publications/m/item/covid-19-public-health-emergency-of-international-concern-\(pheic\)-global-research-and-innovation-forum](https://www.who.int/publications/m/item/covid-19-public-health-emergency-of-international-concern-(pheic)-global-research-and-innovation-forum).
- World Health Organization (WHOb). 2020. "Transmission of SARS-CoV-2: implications for infection prevention precautions." *Scientific brief, 09 July 2020* (2020): 1-10. <https://www.who.int/news-room/commentaries/detail/transmission-of-sars-cov-2-implications-for-infection-prevention-precautions>.
- World Health Organization (WHOc).2020. "guidelines on physical activity and sedentary behaviour". Available at: <https://apps.who.int/iris/handle/10665/336656>
- Wu, X. Y., Han, L. H., Zhang, J. H., Luo, S., Hu, J. W., & Sun, K. (2017). The influence of physical activity, sedentary behavior on health-related quality of life among the general population of children and adolescents: A systematic review.

PloS One, 12(11), e0187668–e0187668.

<https://doi.org/10.1371/journal.pone.0187668>

Wunsch, K., Kienberger, K., & Niessner, C. (2022). Changes in Physical Activity Patterns Due to the Covid-19 Pandemic: A Systematic Review and Meta-Analysis. *International journal of environmental research and public health*, 19(4), 2250. <https://doi.org/10.3390/ijerph19042250>

Yang, S., Guo, B., Ao, L., Yang, C., Zhang, L., Zhou, J., & Jia, P. (2020). Obesity and activity patterns before and during COVID-19 lockdown among youths in China. *Clinical Obesity*, 10(6). <https://doi.org/10.1111/cob.12416>