

## SOLUTIONS FOR QUASILINEAR ELLIPTIC SYSTEMS WITH VANISHING POTENTIALS

BILLEL GHERAIBIA

---

ABSTRACT. In this paper, we study the following strongly coupled quasilinear elliptic system:

$$\begin{cases} -\Delta_p u + \lambda a(x)|u|^{p-2}u = \frac{\alpha}{\alpha + \beta}|u|^{\alpha-2}u|v|^\beta, & x \in \mathbb{R}^N, \\ -\Delta_p v + \lambda b(x)|v|^{p-2}v = \frac{\beta}{\alpha + \beta}|u|^\alpha|v|^{\beta-2}v, & x \in \mathbb{R}^N, \\ u, v \in D^{1,p}(\mathbb{R}^N), \end{cases}$$

where  $N \geq 3$ ,  $\lambda > 0$  is a parameter,  $p < \alpha + \beta < p^* := Np/(N - p)$ . Under some suitable conditions which are given in section 1, we use variational methods to obtain both the existence and multiplicity of solutions for the system on an appropriated space when the parameter  $\lambda$  is sufficiently large. Moreover, we study the asymptotic behavior of these solutions when  $\lambda \rightarrow \infty$ .

---

2010 *Mathematics Subject Classification*. Primary: 35H30, 35J20; Secondary: 35J45.

*Key words and phrases*. Positive solutions; potential well; quasilinear elliptic systems.

This work was partially supported by NSFC (No. 11301204, No. 11371159), self-determined research funds of CCNU from colleges' basic research and operation of MOE(CCNU16A05011).